

DYNC1H1 p.Val314Ile | JLF V3, JLF V2 | 24 aa

disclosed minimal epitope KRFHATISF at 2-10

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1

0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

ERAP1 / ERAMER

cytosol / ER motifs

TPP2

+12 more in CSV

MHC-I candidate ligands

C07:01 0.0021%

C01:02 0.69%

B27:05 1.9%

B08:01 1.3%

B27:05 0.92%

A01:01 1.3%

G K R F H A T I S F D T D T G L K Q A L E T K K

MHC-II candidate ligands

DP01:03/04:01 3%

DQ05:01/02:01 1.5%

DR03:01 1.6%

ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward

NetCleave II native 0-1

0.5 display

SERUM / EXTRACELLULAR - native scores and recognition motifs

DPP4 native (intact SLP)

serum / extracellular motifs

MME

MME

MME

MME

MME

ACE

CPB2+CPN1

SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; >=3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS <=2%/<=5% rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.