

BMP1 p.Ser366Thr | JLF V3 | 23 aa

disclosed minimal epitope KIILNFTTL at 9-17

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1

0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

ERAP1 / ERAMER

cytosol / ER motifs

ERAP2+NPEPPS

TPP2

PREP

+12 more in CSV

MHC-I candidate ligands

B27:05 1.3%

B08:01 0.68%

C01:02 0.26%

B27:05 1.4%

R | I | S | V | T | P | G | E | K | I | I | L | N | F | T | T | L | D | L | Y | R | S | R

1

5

10

DP01:03/04:01 4.6%

20

23

DP01:03/04:01 4.7%

DP01:03/04:01 4.5%

+3 more in CSV

ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward

NetCleave II native 0-1

0.5 display

SERUM / EXTRACELLULAR - native scores and recognition motifs

DPP4 native (intact SLP)

-0.0052

serum / extracellular motifs

MME

MME

MME

MME

MME

MME

MME

MME

ACE

CPB2+CPN1

SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; >=3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS <=2%/<=5% rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.