




GLIS3 p.Ser775fs | JLF V3 | 80 aa- segment 2/3, bonds 28-54

minimal epitope/window not disclosed for this SLP


CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export

four model-native tracks



ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay



EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure



DPP4 / qPISA N-terminal 2-residue removal **SITE OUTSIDE THIS SEGMENT | full SLP: 8% predicted loss** Intact SLP bond 2 is outside this segment

MME / neprilysin Internal hydrophobic preference **NOT ASSESSED / OUTSIDE RULE SCOPE** Rule is limited to peptides of at most 30 residues

FAP - internal Gly-Pro|non-Pro recognition 0 motif matches / 27 assessed bonds - partial recognition rule

FAP - N-terminal Remove 2 residues: X-Pro|non-Pro **SITE OUTSIDE THIS SEGMENT** Intact N-terminal bond 2 is outside this segment

ANPEP / CD13 exposed N-terminus only **SITE OUTSIDE THIS SEGMENT** Intact N-terminal bond 1 is outside this segment

ENPEP exposed N-terminus only **SITE OUTSIDE THIS SEGMENT** Intact N-terminal bond 1 is outside this segment

Triangle = motif match | hollow circle = no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted peptide-signal loss in the source 4 h assay. Native scores remain in CSV. Motifs indicate sequence matches only.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and >=3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.