


DYNC1H1 p.Val314Ile | JLF V3, JLF V2 | 24 aa


disclosed minimal epitope KRFHATISF at 2-10

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export

four model-native tracks



MHC-I: 8 shown / 18 eligible (top-10 cap)



MHC-II: 5 shown / 47 eligible (top-10 cap)

ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay



EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure



DPP4 / qPISA

N-terminal 2-residue removal

NOT SCORED / OUTSIDE MODEL SCOPE

Dataset EV2 has missing coefficients for P1=K: P1':R

5 motif matches / 23 assessed bonds - partial recognition rule

MME / neprilysin

Internal hydrophobic preference



FAP - internal

Gly-Pro|non-Pro recognition

0 motif matches / 23 assessed bonds - partial recognition rule



FAP - N-terminal

Remove 2 residues: X-Pro|non-Pro

0 motif matches / 1 assessed bond - partial recognition rule



ANPEP / CD13

exposed N-terminus only

0 motif matches / 1 assessed bond - partial recognition rule



ENPEP

exposed N-terminus only

0 motif matches / 1 assessed bond - partial recognition rule



Triangle = motif match | hollow circle = no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted peptide-signal loss in the source 4 h assay. Native scores remain in CSV. Motifs indicate sequence matches only.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and >=3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.