

EPG5 p.Ala869Thr | CeGaT | 23 aa

minimal epitope/window not disclosed for this SLP

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export

four model-native tracks

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display


NetCleave I native 0-1

0.5 display


NetChop 20S (in vitro) native 0-1

0.5 display

MHC-I: 3 shown / 4 eligible (top-10 cap)



MHC-II: 5 shown / 38 eligible (top-10 cap)



ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay

NetCleave II native 0-1

0.5 display

EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure



DPP4 / qPISA

N-terminal 2-residue removal

NOT SCORED / OUTSIDE MODEL SCOPE
Dataset EV2 has missing coefficients for P1=K; P1':S

MME / neprilysin

Internal hydrophobic preference

3 motif matches / 22 assessed bonds - partial recognition rule

FAP - internal

Gly-Pro|non-Pro recognition

0 motif matches / 22 assessed bonds - partial recognition rule

FAP - N-terminal

Remove 2 residues: X-Pro|non-Pro

0 motif matches / 1 assessed bond - partial recognition rule

ANPEP / CD13

exposed N-terminus only

0 motif matches / 1 assessed bond - partial recognition rule

ENPEP

exposed N-terminus only

0 motif matches / 1 assessed bond - partial recognition rule

Triangle = motif match | hollow circle = no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted peptide-signal loss in the source 4 h assay. Native scores remain in CSV. Motifs indicate sequence matches only.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and >=3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.