


disclosed minimal epitope RVVPFTKAL at 9-17

EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure



Triangle = motif match | hollow circle = no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted peptide-signal loss in the source 4 h assay. Native scores remain in CSV. Motifs indicate sequence matches only.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and ≥ 3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.


B

SMC5 p.Arg371Lys | JLF V3 | 21 aa


disclosed minimal epitope RQKRIGNTR at 2-10

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export

four model-native tracks



EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure



Triangle = motif match | hollow circle = no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted peptide-signal loss in the source 4 h assay. Native scores remain in CSV. Motifs indicate sequence matches only.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.


RED All four cytosolic tracks assessed the bond and ≥ 3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.

C

PIP5K1A p.Gly474fs | JLF V3 | 20 aa

disclosed minimal epitope AAPVATPAL at 7-15




ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay

NetCleave II native 0-1

0.5 display

EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure



Triangle = motif match | hollow circle = no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted peptide-signal loss in the source 4 h assay. Native scores remain in CSV. Motifs indicate sequence matches only.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and >=3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.


D

DYNC1H1 p.Val314Ile | JLF V3, JLF V2 | 24 aa


disclosed minimal epitope KRFHATISF at 2-10

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export

four model-native tracks



EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure



Triangle = motif match | hollow circle = no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted peptide-signal loss in the source 4 h assay. Native scores remain in CSV. Motifs indicate sequence matches only.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and >=3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.