

MAP2 p.Leu867fs | JLF V2, JLF V1 | 24 aa

disclosed minimal epitope RVVPFTKAL at 9-17

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1

0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

ERAP1 / ERAMER

cytosol / ER motifs

ERAP2+HPEP5

TPP2

PREP

+15 more in CSV

MHC-I candidate ligands

C07:01 0.058%

B08:01 0.13%

B08:01 0.096%

C01:02 0.04%

K | T | V | R | I | Y | Q | G | R | V | V | P | F | T | K | A | L | M | I | K | F | E | E | I

MHC-II candidate ligands

DQ04:01/04:02 1.1%

DQ04:01/04:02 1.2%

DQ04:01/04:02 1.9%

15

20

24

ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward

NetCleave II native 0-1

0.5 display

SERUM / EXTRACELLULAR - native scores and recognition motifs

DPP4 native (intact SLP)

0.49

serum / extracellular motifs

MME

MME

MME

MME

MME

MME

ACE

MME

SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; >=3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS Blue/purple tint and outlines show displayed <=2%/<=5% rank candidates; internal red ticks are pre-binding cut evidence. All candidates remain in CSV.