

KDM3B p.Pro633Ser | JLF V2, JLF V1 | 26 aa

minimal epitope/window not disclosed for this SLP

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1

0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

ERAP1 / ERAMER

cytosol / ER motifs

NPEPPS

TPP2

PREP

PREP

PREP

+7 more in CSV

MHC-I candidate ligands

C01:02 0.29%

C01:02 0.17%

C07:01 0.98%

C01:02 0.85%

C01:02 0.53%

H E N L F L Q P P K L S R E E S S N P F L A F V E K

MHC-II: none <= 5% rank

1

5

10

15

20

25

26

ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward

NetCleave II native 0-1

0.5 display

SERUM / EXTRACELLULAR - native scores and recognition motifs

DPP4 native (intact SLP)

0.028

serum / extracellular motifs

MME

MME

MME

MME

MME

MME

MME

ACE

CPB2+CPN1

SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; >=3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS Blue/purple tint and outlines show displayed <=2%/<=5% rank candidates; internal red ticks are pre-binding cut evidence. All candidates remain in CSV.