

Branch-Dependent Proper Time

Oleg Ponfilenok

Independent researcher, Phan Thiet, Vietnam

Corresponding author: ponfil@gmail.com

Abstract

We attach the Károlyházy uncertainty to *proper time* and postulate that its fluctuations are independent across quantum branches. Interference is sensitive to the difference of branch phase contributions. A fully distinguishable which-path carrier contributes its rest energy mc^2 ; internal branches contribute the transition energy. For a centre-of-mass interferometer the visibility is $V/V_0 = \exp(-\chi)$ with $\chi = (mc^2/\hbar)^2(\xi t_P^2 t)^{2/3}$, where t is the superposition time, t_P is the Planck time, and $\xi = \mathcal{O}(1)$. Attributing the entire contrast deficit in the Vienna sodium-nanoparticle interferometry at $m \approx 172$ kDa and $v \approx 160$ m/s to the mechanism gives $\xi \lesssim 1.1$. A decisive test is near: at $m \sim 1$ MDa and $v \approx 25$ m/s the model predicts practically complete loss of interference, whereas the standard quantum model expects measurable fringes.

Keywords: proper time; branch clock independence; Károlyházy uncertainty; decoherence; matter-wave interferometry; quantum gravity

1 Introduction

In quantum mechanics the status of the wave function—an ontic object or a computational bookkeeping of an ensemble of possibilities—remains unsettled. We adopt the second, ensemble reading [1, 2]: $|\psi\rangle = \sum_i c_i |i\rangle$ catalogues distinct variants of the system, and $|c_i|^2$ are their weights.

In the relational view of time, a clock is a physical subsystem and “time” is a correlation of the clock and the rest of the system within their joint state [3]. If each branch is a genuine variant of the system, then it carries its own world line and hence its own proper time. Penrose reaches a structurally identical statement from the opposite direction: each component of the superposition sources its own space-time, and “since there is no canonical way of relating the time parameters in the two space-times, a fundamental uncertainty in the time-translation operator arises” [4]. Consistently with this, proper time accumulates along a world line and is not fixed by a single common scale for distinct four-dimensional paths: in an interferometer with different trajectories one cannot, without an additional postulate, impose a single common proper-time parameter on all branches.

The idea that the inaccuracy of a physical clock can extinguish interference is older than the present model. Salecker and Wigner bounded the accuracy of a clock by its mass [7]. Károlyházy, and later Ng and van Dam, showed that gravity stops one from sending this bound to zero by making the clock arbitrarily heavy: what remains is the cubic relation $\delta t^3 \sim t_P^2 t$ [6, 8, 9]. Gambini, Porto and Pullin turned that relation into a decoherence law: in the energy basis of a single Hamiltonian the off-diagonal element decays as $\exp(-\omega_{nm}^2 t_P^{4/3} t^{2/3})$, where ω_{nm} is a difference of eigenfrequencies [40, 41]. While states of equal energy interfere, $\omega_{nm} = 0$ and there is no decay. Milburn’s intrinsic decoherence belongs with that decay; Adler’s simplest equation coincides with it [37, 38, 39]. There too the populations are conserved and the dephasing follows an energy difference, but the exponent is linear in time rather than growing as $t^{2/3}$.

Alongside these are constructions in which time is attached to a place rather than to a branch. Diósi coupled rest energy to random shifts of local time [39]. With his spatial correlator the speed of light cancels, and the master equation coincides with Diósi–Penrose collapse: the rate is set by the gravitational self-energy rather than by mc^2 , and population redistribution and spontaneous emission appear [18]. In quantum-clock interference a proper-time difference along two paths records which-path information unitarily, through internal degrees of freedom or a gravitational gradient [14, 15]; there is no stochastic Planckian noise.

The present work differs in that it takes the cubic clock bound and assigns independent proper-time shifts to the branches of one split. In place of the eigenfrequency difference ω_{nm} stands the energy of whatever is in superposition: for which-path this is mc^2 . Arms of equal total energy therefore still lose their mutual phase. A detailed comparison of the channels is given in § 8.4.

The contribution of this work to theoretical physics is one operational postulate, branch-clock independence, which converts Károlyházy proper-time uncertainty into a falsifiable rest-phase dephasing law for matter-wave interferometry, together with a branch-sensitive energy that is invariant under a global shift of the Hamiltonian.

2 Definitions and the postulate

2.1 Notation.

- t — the time the system spends in superposition (a proper-time interval along the world line);
- δt — the proper-time uncertainty over the interval t ; δt_a — the realized shift on branch a (the postulate), $\text{Var}(\delta t_a) = \delta t^2$;
- $t_P = \sqrt{\hbar G/c^5}$ — the Planck time;
- E_{br} — the characteristic energy of the *split* subsystem in the interfering branch pair;
- $E_{a|b}$ — the energy of branch a in the superposition: the energy carried on that branch by degrees of freedom that distinguish it from branch b , measured from the physical vacuum or ground state of the split sector after removing the sector common to both branches;
- $\omega_{\text{eff}} = E_{\text{br}}/\hbar$ — the corresponding effective phase-noise frequency;
- m — the carrier mass in the which-path case (then $E_{\text{br}} = mc^2$, $\omega_{\text{eff}} = \omega_C = mc^2/\hbar$);
- $\xi = \mathcal{O}(1)$ — a dimensionless numerical coefficient.

2.2 Definition of a branch.

These are new definitions. A branch is a term of a superposition, as in standard quantum mechanics. $E_{a|b}$ is the rule adopted here for the energy of that term.

A branch a is the term $c_a|a\rangle$ of one superposition $|\psi\rangle = \sum_a c_a|a\rangle$, differing in a *physical degree of freedom* that participates in interference: a spatial-motional state (which-path or current), or an internal eigenstate of H_{int} (spin, a hyperfine or optical level, a qubit). One connected and indistinguishable wave packet carries one clock. Branching occurs when evolution splits the state into distinguishable terms, and ceases when those terms recombine or the apparatus registers which-path. Which terms of this superposition count as branches is fixed by one split: the split whose interference is read out.

The energy of branch a in the superposition, $E_{a|b}$, is the energy carried on that branch by the degrees of freedom that distinguish it from branch b . The zero of energy is the physical

vacuum or the ground state of the split sector. A sector represented identically on both branches is common: it is subtracted before the remaining energy is multiplied by δt_a and δt_b . What couples to the independent clock shifts is this $E_{a|b}$.

On a motional pair both branches carry the same exclusive rest energy, $E_{a|b}^{\text{mot}} = E_{b|a}^{\text{mot}} = E_{\text{br}}^{\text{mot}}$. When the spatial packets of the branches are distinguishable, $E_{a|b}^{\text{mot}} = Mc^2$. How that energy is assembled from the particle masses is given in § 4.

For internal states $|0\rangle, |1\rangle$ of a common carrier the zero lies at the ground state: branch $|0\rangle$ carries no exclusive energy, and branch $|1\rangle$ carries the transition energy, $E_{0|1}^{\text{int}} = 0$ and $E_{1|0}^{\text{int}} = \Delta E$. If motion and an internal level both differ, the two contributions add on each branch: $E_{a|b} = E_{\text{br}}^{\text{mot}} + E_a^{\text{int}}$ and $E_{b|a} = E_{\text{br}}^{\text{mot}} + E_b^{\text{int}}$.

A global shift $H \mapsto H + C1$ adds the same number to the reference energy of the common sector or of the ground state. After that sector is subtracted, $E_{a|b}$ and $E_{b|a}$ are unchanged, and so are the visibilities.

2.3 Proper-time fluctuation.

The proper-time fluctuation δt is adopted from the Károlyházy uncertainty relation and is not a new postulate of this work.

The accumulated uncertainty of the proper-time interval t measured along a world line is

$$\boxed{\delta t^3 = \xi t_P^2 t}, \quad (1)$$

with $\xi = \mathcal{O}(1)$. Equation (1) is the temporal form of the Károlyházy uncertainty relation [6]; the same cubic scaling is consistent with the clock arguments of Salecker–Wigner [7], Ng–van Dam [8, 9], and the holographic bound on space-time uncertainty [10]. We do not fix the numerical coefficient ξ : in these works it is likewise never pinned down more precisely than to $\mathcal{O}(1)$.

We stress where gravity enters δt . The Salecker–Wigner lower bound for a clock of mass m_{cl} , $\delta t \gtrsim (\hbar t / m_{\text{cl}} c^2)^{1/2}$, vanishes as $m_{\text{cl}} \rightarrow \infty$. The “not a black hole” constraint stops that limit: the optimal m_{cl} drops out and leaves the mass-independent minimum $\delta t^3 \sim t_P^2 t$ adopted in Equation (1). The rest mass is equally essential: the uncertainty δt is defined along a world line with nonzero proper time; without rest mass there is no proper time. For a photon $m = 0$, $d\tau = 0$, and the rest-energy channel does not apply to a light-like carrier.

2.4 Postulate: branch clock independence.

This is the postulate of the model. In standard quantum mechanics one time parameter is common to the whole superposition. Here branches a and b have their own proper-time shifts, δt_a and δt_b , and those shifts are independent.

For two branches a, b (in the sense of the definitions above) over the superposition interval t (the same as in (1)), the realized shifts $\delta t_a, \delta t_b$ are independent, centered Gaussian random variables of accumulated proper time with $\text{Var}(\delta t_a) = \text{Var}(\delta t_b) = \delta t^2$ and $\text{Cov}(\delta t_a, \delta t_b) = 0$. This is a substantive postulate, motivated by the ensemble ontology (§ 1): distinct variants do not share a common clock; there is no strict derivation from that ontology, and what remains irreducible is the premise that the holographic information budget is allocated *per branch* rather than as a single shared register.

Within one branch, all degrees of freedom *not* participating in the split share that branch’s clock. They do not form a branch of their own: the part that is the same on both arms enters each term of the split, advancing with δt_a on term a and with δt_b on term b .

Crucially, δt_a is not a fluctuation of an external metric field and not environmental noise common to nearby trajectories. It is an *ontic* random shift of proper time; its magnitude over

the interval t is set by δt from (1)—the holographic estimate of such uncertainty [10]. An environmental source—for example, a difference of gravitational potentials on the interferometer arms—would give a *correlated* shift δt_a , the closer to common the nearer the branches, and it would cancel in the relative phase; but that is a systematic shift, not noise. The holographic uncertainty attaches to the split subsystem of a branch, not to spatial points, so spatial proximity of branches does not by itself correlate their clocks. Attachment to discrete branches rather than positions is essential: noise independent for arbitrarily close *positions* would mean a random phase field with unbounded gradient—hence a force and momentum diffusion. Noise independent for discrete *branches* is pure dephasing without force.

3 Decoherence functional

For a composite system we use the decomposition $H = mc^2 + P^2/(2m) + H_{\text{int}}$. For which-path ($E_{a|b} = E_{b|a} = E_{\text{br}} = mc^2$ by the definitions) the phase contribution of the split subsystem gives the relative phase

$$\Delta\varphi = \frac{E_{a|b}}{\hbar} \delta t_a - \frac{E_{b|a}}{\hbar} \delta t_b \approx \frac{mc^2}{\hbar} (\delta t_a - \delta t_b), \quad (2)$$

since $E_{a|b} = E_{b|a} = mc^2$ and interference involves the *difference* of branch-exclusive contributions, not an absolute eigenvalue of the total Hamiltonian.

For a single superposition interval, the relation (1) and the postulate fully specify the distribution of $\delta t_a, \delta t_b$ used in the derivation below; the multi-time structure of the noise is discussed in § 8.1. The general formula follows from (2) and the relation (1) and the postulate. By the postulate, with $\text{Var}(\delta t_a) = \text{Var}(\delta t_b) = \delta t^2$:

$$\text{Var}(\Delta\varphi) = \frac{E_{a|b}^2 + E_{b|a}^2}{\hbar^2} \delta t^2. \quad (3)$$

If the centre of mass branches (which-path), (3) gives

$$\text{Var}(\Delta\varphi) = 2\omega_{\text{eff}}^2 \delta t^2, \quad \omega_{\text{eff}} = \frac{mc^2}{\hbar}. \quad (4)$$

With *shared* clocks the global phase $mc^2\tau/\hbar$ would cancel in $\Delta\varphi$; with independent clocks, by the postulate, the vacuum-referenced branch-exclusive accumulations enter the difference. A constant common term in the total Hamiltonian is removed by the definitions and cannot generate dephasing. For internal branches $|0\rangle, |1\rangle$ with a common carrier $|C\rangle$ we take $E_{\text{br}} = \Delta E$ and $\omega_{\text{eff}} = \Delta E/\hbar$; since here $E_{0|1} = 0$ and $E_{1|0} = \Delta E$, dephasing is twice as weak as in the symmetric case: $\chi = \omega_{\text{eff}}^2 \delta t^2/2$.

Since $\delta t_a, \delta t_b$ are Gaussian by the postulate, the relative phase $\Delta\varphi$ is also Gaussian and $\langle e^{i\Delta\varphi} \rangle = \exp[-\text{Var}(\Delta\varphi)/2]$; the factor 2 from the two branches and the factor 1/2 from Gaussian averaging cancel. In the which-path case, for fringe visibility (V relative to the reference V_0):

$$\boxed{\frac{V}{V_0} = e^{-\chi}, \quad \chi = \omega_{\text{eff}}^2 \delta t^2 = \left(\frac{mc^2}{\hbar}\right)^2 (\xi t_P^2 t)^{2/3}.} \quad (5)$$

For one pair of orthogonal branches

$$\boxed{\chi_{ab} = \frac{E_{a|b}^2 + E_{b|a}^2}{2\hbar^2} \delta t^2.} \quad (6)$$

The branches are the arms of one split between which the apparatus reads out interference. For which-path the energy of a branch in the superposition is the same on every arm: mc^2 per arm.

Its own δt_a gives one factor $e^{-\chi}$ to every arm pair, including three or more arms. Independent splits are not combined into one: in a product of single-atom interferometers each atom has its own clock pair and $\mathcal{D} = 1$, whereas a NOON state and a rigid cluster are one split, with Nmc^2 and Mc^2 . The factor $e^{-\chi}$ is placed between orthogonal full branches; overlap of the one-particle $\gamma^{(1)}$ enters only $\mathcal{D}_s$. For $E_{a|b} = E_{b|a} = mc^2$ the equation reduces to (5). Because $E_{a|b}$ and $E_{b|a}$ are vacuum- or ground-state-referenced energies of the split sector, (6) is unchanged by $H \mapsto H + C\mathbf{1}$. The scaling form $\chi \propto m^2 t^2/3$ in a centre-of-mass interferometer follows from the relation (1) and the postulate and constitutes the central testable prediction; the overall coefficient carries an $\mathcal{O}(1)$ uncertainty (through $\xi^{2/3}$, the interpretation of δt , and the choice of the effective t ; see § 6). The mass-quadratic dependence follows from rest-phase accumulation ($\omega_{\text{eff}} \propto mc^2$) and matches the typical dephasing scaling in models of metric and conformal space-time fluctuations [11, 12]; in the high-mass interferometry programme, this mass growth of the signal is highlighted as a practical motivation for searching for hypothetical effects [13].

Relation to clock interference and the Compton phase. The conceptual framework of “proper time along each branch” is already used in quantum-clock interference. In Zych et al., an internal state acts as a clock: a deterministic proper-time difference along the two paths records which-path information in the internal degree of freedom and reduces visibility [14]. Pikovski et al. developed this mechanism for a composite object in a gravitational field: an externally imposed potential difference unitarily entangles the centre of mass with the internal-energy distribution, and dephasing appears after tracing out the internal degrees of freedom [15]. In contrast, the present model does not rely on an external potential difference or on a dedicated internal degree of freedom that unitarily encodes which-path information; it adopts the Károlyházy relation for δt and the postulate of branch-clock independence, and therefore predicts irreversible ensemble dephasing even for an object in its internal ground state.

Using $\omega_C = mc^2/\hbar$ requires a caveat: in standard quantum mechanics the Compton phase cancels in a closed interferometer [16], whereas atom interferometry has been interpreted as providing access to the mass frequency [17]. The present model adds the independent premise that the vacuum-referenced rest energy of branch-exclusive massive modes couples to the independent shifts $\delta t_a, \delta t_b$ and thereby becomes a random *relative* contribution. It does not couple an arbitrary identity term in the total Hamiltonian: The definitions first remove the common sector and its reference energy. Without this premise the central prediction $\chi \propto m^2$ does not arise (see also § 8).

Planck mass boundary. The decoherence time t_{dec} is defined by the condition $\chi = 1$. From (5):

$$t_{\text{dec}} = \frac{t_C^3}{\xi t_P^2}, \quad t_C = \frac{\hbar}{mc^2}. \quad (7)$$

Setting $t_{\text{dec}} = t_C$, we obtain $t_C = \sqrt{\xi} t_P$, i.e. the characteristic scale

$$m_* = \frac{m_P}{\sqrt{\xi}}, \quad m_P = \sqrt{\hbar c/G} \approx 2.2 \times 10^{-8} \text{ kg}. \quad (8)$$

For $\xi = \mathcal{O}(1)$ this is of the order of the Planck mass. For $m \gg m_*$ we have $t_{\text{dec}} \ll t_C$: inter-branch coherence vanishes over an interval below the uncertainty δt from (1) and in principle inaccessible to direct measurement. The mechanism remains pure dephasing without population redistribution or spontaneous emission [18]. The growth of the von Neumann entropy of the averaged state is treated in § 5. The ontological consequences of the macroscopic limit—an effective ensemble mixture in the branch basis, objective loss of inter-branch interference before readout, and distinction from dynamical collapse models—are discussed in § 8.3.

4 Examples

$E_{\text{br}}^{\text{mot}}$ is the rest energy of those particles by which the already given branches differ. The full mass of the body enters only when the spatial packets of all its particles are distinguishable between the two branches. For particles of species s —one type and one mass m_s —let $\gamma_{a,s}^{(1)}$ and $\gamma_{b,s}^{(1)}$ be the one-particle reduced density matrices of the branches, normalized by $\text{Tr} \gamma_{a,s}^{(1)} = \text{Tr} \gamma_{b,s}^{(1)} = N_s$. The number of distinguishable particles of that species and the corresponding energy are

$$\mathcal{D}_s(a, b) = \frac{1}{2} \left\| \gamma_{a,s}^{(1)} - \gamma_{b,s}^{(1)} \right\|_1, \quad E_{\text{br}}^{\text{mot}} = c^2 \sum_s m_s \mathcal{D}_s(a, b). \quad (9)$$

Identical particles of one species share one matrix. Different species are added separately: each has its own mass and its own packet overlap. The definition does not depend on the one-particle basis and applies to identical particles [5]; throughout, $0 \leq \mathcal{D}_s \leq N_s$. A state that is the same on both branches does not enter the difference of the matrices. For a motional pair this energy is the exclusive rest energy of each branch,

$$E_{a|b}^{\text{mot}} = E_{b|a}^{\text{mot}} = E_{\text{br}}^{\text{mot}} = c^2 \sum_s m_s \mathcal{D}_s(a, b). \quad (10)$$

This is the energy, relative to the vacuum, of the occupied branch-exclusive matter modes. The formula does not choose the decomposition into branches.

Full resolution. The spatial packets of every constituent are distinguishable: $\mathcal{D}_s = N_s$ and $E_{\text{br}}^{\text{mot}} = Mc^2$. This is the case of a rigid cluster whose branches are centre-of-mass shifts of the whole body.

Partial overlap. A species contributes $m_s \mathcal{D}_s c^2$. If the packets have separated only in part, the full mass of the body is replaced by the sum of $m_s \mathcal{D}_s$ over species.

Common carrier. Only an internal level differs, and the spatial-motional part $|C\rangle$ is common. The rest mass of the carrier does not enter the difference of the matrices; $E_{\text{br}}^{\text{int}} = |\Delta E|$, with the ordered values $E_{0|1}^{\text{int}} = 0$ and $E_{1|0}^{\text{int}} = \Delta E$ from the definitions.

Collective mode. A mode of a common body is displaced by much less than the atomic zero-point motion. E_{br} receives the mode energy, not $\mathcal{D}_s mc^2$.

Current. The current in the definitions is the persistent current in the superconducting loop of a flux qubit. The branches are the states with opposite sense of that current, clockwise and counterclockwise. The loop is the same in both, so the condensate electrons of the two branches overlap almost completely: the energy receives $\mathcal{D}_e$, not the mass of every electron. For the devices considered, $\mathcal{D}_e \simeq 42\text{--}5750$ against $10^6\text{--}10^{10}$ condensate electrons [27, 28]. The estimate of χ is given in § 7.1.

5 Growth of the von Neumann entropy

Dephasing changes the von Neumann entropy of the averaged state. For a fixed realization of the shifts $\delta t_a, \delta t_b$ the vector

$$|\psi\rangle = c_a |a\rangle + c_b e^{i\Delta\varphi} |b\rangle \quad (11)$$

remains pure, $S = 0$; the phase $\Delta\varphi$ is given by (2). A mixture appears after averaging over the distribution of these shifts. Since $\delta t_a, \delta t_b$ are Gaussian by the postulate, the off-diagonal element is multiplied by $\langle e^{i\Delta\varphi} \rangle = e^{-\chi}$:

$$\bar{\rho}_{ab} = \rho_{ab}(0) e^{-\chi}, \quad (12)$$

where χ is the exponent (6). The von Neumann entropy refers to this matrix,

$$S = -\text{Tr}(\bar{\rho} \ln \bar{\rho}). \quad (13)$$

For two branches of equal weight the eigenvalues of $\bar{\rho}$ are

$$\lambda_{\pm} = \frac{1 \pm e^{-\chi}}{2}, \quad S = -\lambda_+ \ln \lambda_+ - \lambda_- \ln \lambda_-. \quad (14)$$

At $\chi = 0$ the spectrum is $(1, 0)$ and $S = 0$. Since $\chi \propto t^{2/3}$, as the superposition time grows λ_+ and λ_- approach each other and S increases. As $\chi \rightarrow \infty$ the coherence has vanished, the spectrum coincides with the branch weights, and the entropy saturates at the Shannon value

$$S \rightarrow -\sum_i |c_i|^2 \ln |c_i|^2 \quad (15)$$

(for equal weights, $S \rightarrow \ln 2$). A further increase of χ adds no new eigenvalues.

The populations $|c_i|^2$ stay constant throughout. λ_+ decreases from 1 to 1/2 and λ_- increases from 0 to 1/2: the off-diagonal elements decay as $e^{-\chi}$, and the original pure vector ceases to be an eigenvector of the averaged matrix. This is the growth of the entropy of the averaged state of the split system itself, not of a reduced subsystem. In environmental decoherence the joint state of system and environment can remain pure, and entropy appears after the environment is traced out. Here there is no environment recording which-path information, and the purity of $\bar{\rho}$ cannot be restored by tracing out another quantum subsystem. The mixture and the growth of S arise by averaging over the noise of the postulate. The map on $\bar{\rho}$ is linear, $\rho_{ab} \mapsto \rho_{ab} e^{-\chi}$.

6 Confrontation with experiment

6.1 Matter-wave interferometry: the primary probe

By the definitions and the postulate the mechanism is pure rest-phase dephasing without force or momentum diffusion (see the postulate and § 8). The experimental consequence is that non-interferometric bounds on Diósi–Penrose (the underground search for spontaneous radiation [18], the Majorana Demonstrator limit [19], and X-ray tests [20]) do not apply. An atomic clock has no mc^2 channel: a superposition of internal levels proceeds with $E_{\text{br}} = \Delta E$, and for a microwave transition χ is negligible. Matter-wave interferometry of the centre of mass is therefore the most direct laboratory probe of the mechanism with $E_{\text{br}} = mc^2$.

For the same reason, purely optical interferometers provide no direct constraint ($d\tau = 0$ for a photon); LIGO and the Holometer constrain external metric and spatial models [21], but not branch-dependent proper time (BDPT; see § 8). Massive mirrors would be subject to the mechanism only in their own centre-of-mass superposition; in the usual classical optical regime they are not, and existing optical null results therefore place no direct bound on ξ .

6.2 Sodium nanoparticles: the Vienna group (2026)

Setup parameters. We confront Equation (5) with the sodium-nanoparticle interferometer of the Vienna group [22, 23]:

- clusters of 5000–10000 Na atoms with a mass-distribution centre $m \approx 172$ kDa ($m \approx 2.86 \times 10^{-22}$ kg);
- velocity $v \approx 160$ m/s;
- three gratings of period $d = 133$ nm spaced by $L = 0.983$ m;
- a measured fringe visibility $V = 0.10 \pm 0.01$ (second run 0.08 ± 0.01) at 172 kDa, described by the quantum model with a single global factor 0.78 for conventional contrast losses (alignment, gravitational and rotational phase averaging, vibrations, thermal and collisional decoherence);

- a record macroscopicity $\mu = 15.5$ [24, 22].

The relevant superposition time is the Talbot flight of one section $t = L/v \approx 6.1$ ms (justification of the choice—in § 6.3).

Under the sharpened definitions this is not a separate counting rule: the two resolved branches of a rigid cluster are centre-of-mass shifts of the whole body (without change of internal configuration). When their one-atom spatial supports have negligible overlap, $\mathcal{D}_{\text{Na}} = N$ and Equation (9) gives $E_{\text{br}} = Nm_{\text{at}}c^2 = Mc^2$, whence $\chi \propto M^2$. If the overlap is incomplete, M in the formulas below must be replaced by $m_{\text{at}}\mathcal{D}_{\text{Na}}$; the assumption of full branch distinguishability is thus explicit and can be checked against the packet geometry.

6.3 Numerical estimate and constraint on ξ

We first fix the effective superposition time. The definitions and the postulate do not uniquely define the moment at which independent branch clocks “switch on” (§ 8.2): delocalization of the wave function grows already after the first grating (full flight $2L/v \approx 12.3$ ms), but distinguishable diffraction orders—branches in the strict sense of the definitions—form only at the second grating, leaving one section until recombination at the third, $t = L/v \approx 6.1$ ms. For an upper bound we take the *minimal* interpretation $t = L/v$: the smaller t , the smaller the predicted dephasing and the *weaker* (more conservative) the resulting bound on ξ ; any broader interpretation of the superposition time would only tighten the limit. With $\omega_C = mc^2/\hbar = 2.43 \times 10^{29} \text{ s}^{-1}$ (at $m = 172 \text{ kDa}$) and $t_P^2 = 2.91 \times 10^{-87} \text{ s}^2$, Equation (5) gives

$$\chi \approx 0.41 \xi^{2/3} \quad (t = L/v \approx 6.1 \text{ ms}). \quad (16)$$

At $\xi = 1$ this is $\chi \approx 0.41$, i.e. a visibility deficit of $\approx 34\%$.

We use the published raw data and analysis code [25]. In the bin $P_2 = 15.252 \text{ mW}$ three independent scans give

$$V_i = 0.0732 \pm 0.0081, \quad 0.0975 \pm 0.0070, \quad 0.0780 \pm 0.0074, \quad (17)$$

where the errors are 1σ from the covariance matrix of a nonlinear sinusoidal fit. For the displayed rounded values, the inverse-variance mean is $\bar{V} = 0.0840$ with a formal error of 0.0043. For these three scans $\chi_{\text{red}}^2 = 3.08$; following the experiment’s procedure for overdispersed bins, we inflate the error by $\sqrt{\chi_{\text{red}}^2}$ and obtain $\sigma_{\bar{V}} = 0.0076$.

The official quantum curve at this power gives $V_{\text{QM}} = 0.1108$ (without conventional losses); it depends on the polarizability, the grating power, and the velocity. The lower bound on the observed visibility at the one-sided normal quantile is

$$\bar{V} - 1.645 \sigma_{\bar{V}} = 0.0716. \quad (18)$$

We construct the upper bound on ξ at $R_{\text{ord}} = 1$. The observed visibility decomposes as $V_{\text{obs}} = R_{\text{ord}} V_{\text{QM}} e^{-\chi}$, where $R_{\text{ord}} \leq 1$ is the total conventional contrast-loss factor (alignment, vibrations, phase averaging, gas, radiation). Since R_{ord} cannot exceed unity by its physical meaning, regardless of its true value we have $e^{-\chi} \geq V_{\text{obs}}/V_{\text{QM}}$. Attributing *the entire* visibility deficit to our mechanism ($R_{\text{ord}} = 1$), we obtain

$$e^{-\chi} \geq \frac{0.0716}{0.1108} = 0.646, \quad \chi \leq 0.44, \quad (19)$$

and from $\chi = 0.41 \xi^{2/3}$ we obtain

$$\boxed{\xi \lesssim 1.1.} \quad (20)$$

This bound does not depend on the conventional-loss factor R_{ord} : any real losses ($R_{\text{ord}} < 1$) could only tighten it. The number 1.1 corresponds to the quantile 1.645 after inflating the error on three scans, that is, two degrees of freedom. Student’s quantile $t_{0.95}(2) = 2.92$ gives $\xi \lesssim 1.7$.

If one trusts the published estimate of conventional losses $R_{\text{ord}} = 0.78$, the limit tightens: $e^{-\chi} \geq 0.0716/(0.78 \cdot 0.1108) = 0.828$, $\chi \leq 0.188$, whence $\xi \lesssim 0.32$. But this is a conditional result: at a single mass the product $R_{\text{ord}} e^{-\chi}$ is degenerate, and the data do not allow separating the new mechanism from underestimated conventional losses; the interpretation of δt , the choice of effective t , and the overall coefficient ξ also remain model systematics.

Summary for the data: at $\xi = 1$ the model predicts ($R_{\text{ord}} = 1$) visibility $0.1108 e^{-0.41} \approx 0.074$ against the observed 0.0840 ± 0.0076 —a deviation of only 1.3σ , so $\xi = \mathcal{O}(1)$ is *not excluded* by the data; the normal quantile gives $\xi \lesssim 1.1$, and the conditional result (at $R_{\text{ord}} = 0.78$) gives 0.32. A statistically rigorous limit requires a joint likelihood analysis of the raw scans, the quantum-curve parameters, and all conventional contrast-loss channels.

6.4 The decisive near-term test

The already-measured high visibility $V = 0.66 \pm 0.09$ at 0.4–1 MDa [22] does *not* test the mechanism as distinguishable quantum coherence: it was taken in the regime $\lambda_{dB} \lesssim 3$ fm, $L \leq L_T$ ($L_T \sim d^2/\lambda_{dB}$ —the *Talbot length*, the scale of grating self-imaging in the near field of a Talbot–Lau interferometer [22, 13]), where the quantum and classical predictions coincide (geometric optics). A discriminating test requires the quantum regime at high mass. The same programme targets $m \sim 1$ MDa with $v \approx 25$ m/s [22]. Then even with the conservative $t = L/v \approx 39$ ms and $\omega_{\text{eff}} \approx 1.4 \times 10^{30} \text{ s}^{-1}$, Equation (5) at $\xi = 1$ gives

$$\chi \approx 47, \quad V/V_0 = e^{-\chi} \lesssim 10^{-20}. \quad (21)$$

Practically complete loss of interference is robust against $\mathcal{O}(1)$ uncertainty in the coefficient. The standard quantum model of the setup at the same parameters expects measurable fringes with ordinary contrast. Observing fringes at ~ 1 MDa and $v \approx 25$ m/s would falsify the region $\xi = \mathcal{O}(1)$.

6.5 Distinguishing signature

A visibility deficit alone is not diagnostic, since thermal radiation, collisions with residual gas, and velocity spread also reduce the contrast. The signature of the present mechanism is its scaling $\chi \propto m^2 t^{2/3}$ together with *immunity to shielding*: it cannot be reduced by improving the vacuum or cooling the particle. The experimental discriminator is whether a residual deficit persists after the suppression of all known environmental channels and whether it follows the m^2 law. Disappearance of fringes at ~ 1 MDa and $v \approx 25$ m/s with scaling $V \propto \exp[-(mc^2/\hbar)^2(\xi t_P^2 t)^{2/3}]$ after excluding environmental losses would be evidence in favour of the model.

An additional discriminator is the dependence on the maximum spatial branch separation Δx . For momentum-transfer decoherence (collisions and photon emission or scattering), the localization exponent contains factors of the form $1 - \text{sinc}(q\Delta x/\hbar)$: it vanishes for unresolved paths, grows as the environment starts to resolve them, and saturates at the event rate once Δx exceeds the environment’s spatial resolution [26]. It would therefore be incorrect to say that *all* environmental channels grow without bound with Δx . A gravitational gradient usually produces a deterministic geometry-dependent phase; contrast is lost only after averaging over fluctuations of the gradient or trajectories.

The present model is likewise not strictly independent of Δx over the entire range. Under the sharpened definitions,

$$\chi_{\text{BDPT}}(\Delta x) = \left[\frac{c^2 \sum_s m_s \mathcal{D}_s(\Delta x)}{\hbar} \right]^2 (\xi t_P^2 t)^{2/3}. \quad (22)$$

While the packets overlap, $\mathcal{D}_s(\Delta x)$ grows from zero. The specific signature emerges once the branches are resolved: as $\mathcal{D}_s \rightarrow N_s$, increasing Δx further does not change χ_{BDPT} , producing a

plateau that depends only on m and t . This is the direct consequence of attaching the noise to distinguishable branches rather than to spatial points.

The test can be performed in one apparatus: at fixed m , v , superposition time, pressure, and temperature, vary the transverse momentum transfer (preferably the diffraction order; changing the grating period also requires compensating the change in Talbot length) and thereby scan Δx . The quantity to compare is not raw visibility but the residual exponent

$$\chi_{\text{res}}(\Delta x) = -\ln \left[\frac{V_{\text{obs}}(\Delta x)}{V_{\text{QM}}(\Delta x)} \right] - \chi_{\text{env}}^{\text{calc}}(\Delta x), \quad (23)$$

where V_{QM} includes coherent dynamics and grating interactions, while $\chi_{\text{env}}^{\text{calc}}$ contains the calculated gas and thermal contributions. The model predicts a constant addition to χ_{res} on the distinguishability plateau, whereas unsaturated environmental channels retain a characteristic Δx dependence. A flat residual alone is not unique: saturated environmental decoherence and Δx -independent technical losses can mimic it. The Δx scan should therefore be combined with changes in vacuum or temperature and with a test of the $m^2 t^{2/3}$ scaling.

7 Limits on quantum computation

7.1 Composite superpositions: flux qubits and atomic NOON states

The unqualified phrase “energy of the split subsystem” is particularly hazardous for a superconducting flux qubit. Naively counting all 10^6 – 10^{10} electrons carrying superconductivity would give, at the upper end, $E_{\text{br}} \simeq 8.2 \times 10^{-4}$ J and, for $t = 100 \mu\text{s}$ and $\xi = 1$, $\chi \simeq 26.5$; the observed coherence of current states would then exclude the model. A microscopic analysis of the BCS states, however, shows that the near-unit overlap of the one-electron states in the two current branches leaves only a small fraction of occupations changed. An estimate equivalent to the one-particle distinguishability in (9) gives $\mathcal{D}_e \simeq 42$ – 5750 for the devices considered, rather than the total number of condensate electrons [27, 28]. At $t = 100 \mu\text{s}$ this gives $\chi \lesssim 9 \times 10^{-12}$: existing flux qubits are safe for the model. The difference between naive and microscopic counts is not an $\mathcal{O}(1)$ uncertainty; this is why the distinguishability measure is part of the definitions.

The same rule separates two often conflated regimes of atom interferometry. A product state

$$\left[(|L\rangle + |R\rangle)/\sqrt{2} \right]^{\otimes N} \quad (24)$$

contains independent one-atom superpositions; observed first-order coherence compares components with $\mathcal{D} = 1$ and acquires the atomic, not collective, energy mc^2 . In contrast, a NOON state

$$(|N, 0\rangle + |0, N\rangle)/\sqrt{2} \quad (25)$$

has two collective branches with $\mathcal{D} = N$ and $E_{\text{br}} = Nmc^2$. For $N = 10^5$ ^{87}Rb atoms and $t = 10$ ms, Equation (5) predicts $\chi \simeq 1.4 \times 10^3$ at $\xi = 1$, i.e. practically complete suppression of interference for a massive motional NOON state. This additional falsifiable prediction does not yet conflict with experiment: demonstrated atomic NOON states remain few-particle; for example, Ref. [29] realized a second-order state.

7.2 Margolus–Levitin: a heuristic comparison

The decoherence time of the model is the moment when $\chi = 1$ and the visibility falls to e^{-1} ; it is given by Equation (7).

The Margolus–Levitin theorem [30] limits the speed of orthogonal evolution by the energy above the ground level $E_{\text{av}} = \langle H \rangle - E_0$: $\tau_{\perp} \geq \pi\hbar/(2E_{\text{av}})$, whence $\nu_{\text{max}} = 2E_{\text{av}}/(\pi\hbar)$. This is a

question *distinct* from branch decoherence: the Margolus–Levitin theorem concerns the unitary speed of state change, not the noise of independent clocks.

At fixed particle number, the term mc^2 is proportional to the identity and does not generate orthogonal centre-of-mass evolution; it cancels from $E_{\text{av}} = \langle H \rangle - E_0$. Thus the identification $E_{\text{av}} = E_{\text{br}} = mc^2$ follows neither from the theorem nor from the definitions and the postulate. Only as a formal dimensional illustration, under the additional and nonstandard assumption of that identification, one may write the number of “ticks” over the window t_{dec} as

$$N_{\text{max}} = \nu_{\text{max}} t_{\text{dec}} = \frac{2\omega_{\text{eff}}}{\pi} \cdot \frac{t_C^3}{\xi t_P^2} = \frac{2}{\pi\xi} \left(\frac{t_C}{t_P} \right)^2. \quad (26)$$

Since $t_C/t_P = m_P/m$, for which-path

$$N_{\text{max}}^{\text{formal}} = \frac{2}{\pi\xi} \left(\frac{m_P}{m} \right)^2. \quad (27)$$

At $m = m_* = m_P/\sqrt{\xi}$ this formal quantity is $2/\pi \approx 0.64$. It is not a computational-power bound derived from the model. For internal branches, where the transition energy actually generates evolution, the theorem should instead be applied directly using the physical E_{av} , without importing the formula based on mc^2 .

7.3 Formal coincidence with holographic scaling

The same additional identification gives the formal rate $N = (2/\pi) t/t_C$. Combining it with the coherence condition $t_{\text{dec}} \geq t$, i.e. $t_C \geq (\xi t_P^2 t)^{1/3}$, and taking equality at the boundary gives

$$N_{\text{formal}}(t) = \frac{2}{\pi \xi^{1/3}} \left(\frac{t}{t_P} \right)^{2/3}. \quad (28)$$

Over one second this is $\approx 4.5 \times 10^{28}$ at $\xi = 1$. The exponent $2/3$ coincides with the scaling $N \propto (t/t_P)^{2/3}$ obtained by Ng for a “space-time-foam computer” and a black-hole quantum computer [10], because both estimates use the cubic law $\delta t^3 \sim t_P^2 t$. This is a meaningful structural analogy, but not an independent derivation of an operation-count bound: without additional physical justification for $E_{\text{av}} = mc^2$, matching exponents does not turn N_{formal} into a computational limit of the model.

7.4 Phase noise and the qubit register

The definitions and the postulate act on any superposition, including a computational one. For strictly internal levels $|0\rangle, |1\rangle$ with a common one-particle spatial-motional density matrix, the definitions give $E_{\text{br}} = \hbar\omega_q = \Delta E$; the mass of the chip and resonator enters the common $|C\rangle$ and does not enter ω_{eff} . This statement cannot automatically be transferred to a superposition of opposite persistent currents, for which Equation (9) applies as discussed in § 7.1. Superconducting resonators storing a single photon for ~ 34 ms with $\gtrsim 10^{23}$ electrons in their walls [31] illustrate the internal case: with $E_{\text{br}} = \Delta E$ of the photon, not the rest energy of the common walls, long-lived coherence is consistent with the model. The random relative phase between a pair of internal branches over time t :

$$\sigma(t) = \frac{\sqrt{E_{a|b}^2 + E_{b|a}^2}}{\hbar} \delta t = \omega_q (\xi t_P^2 t)^{1/3} \quad (E_{0|1} = 0, E_{1|0} = \hbar\omega_q). \quad (29)$$

For a superconducting qubit ($\omega_q \approx 2\pi \times 5$ GHz) and $t = 1$ s: $\delta t \approx 1.4 \times 10^{-29}$ s, $\sigma \approx 4.5 \times 10^{-19}$ rad (the corresponding dephasing parameter $\chi = \sigma^2/2$). Crucially, the variance (29) is *pairwise* and independent of the number of branches: interference involves only differences

of independent δt_a , and the extremal phase spread over 2^n branches grows only as $\sigma\sqrt{2\ln 2^n} \propto \sigma\sqrt{n}$. If d excitations form one collective branch against the ground state, $E = d\hbar\omega_q$ and $\chi \propto d^2$; for independently split qubits the exponents add, $\chi \propto d$. In the collective case, for the branch with all n excitations the formal limit $n\omega_q\delta t \lesssim 1$ gives $n \lesssim 10^{18}$ —far beyond practical horizons.

7.5 Phase-noise scale: exact and approximate QFT

The quantity σ is the scale of the model’s own relative phase over time t . An angle $\theta \lesssim \sigma$ lies below this scale on a single realization; over many runs such a shift is statistically distinguishable, and the comparison $2\pi/2^n \sim \sigma$ does not forbid an exact QFT. An exact quantum Fourier transform on n qubits contains controlled rotations down to $2\pi/2^n$, and already at $n \gtrsim 64$ these angles fall below $\sim 10^{-19}$ rad. Exact QFT is not used in implementations: by Coppersmith [32] all rotations with $k > \mathcal{O}(\log(n/\varepsilon))$ can be *omitted from the circuit* entirely, losing only ε success probability; the minimum remaining angle $\sim 2\pi\varepsilon/n$ is polynomially, not exponentially, small. For Shor’s algorithm [33] at $n = 10^4$, $\varepsilon = 10^{-2}$ this is $\sim 10^{-5}$ rad—thirteen orders of magnitude above the scale (29). Phase-estimation algorithms that read the answer from the position of an interference peak are not limited by this scale: the angle remaining after the omission lies many orders of magnitude above σ .

7.6 Grover’s algorithm: a heuristic estimate

The case of amplitude amplification [34] is contrasting. Here there is no QFT and nothing to truncate: each iteration rotates the state vector by a fixed angle $2\theta \approx 2/\sqrt{N}$, set by the geometry of the problem, and all $\sim (\pi/4)\sqrt{N}$ iterations form one continuous coherent accumulation without intermediate readouts. The small rotation per step is irremovable: it *is* the algorithm. The match $2/\sqrt{N} \sim \sigma$, with σ taken from (29) at $t = 1$ s, is an order-of-magnitude estimate. The Grover rotation angle and the relative branch phase are different quantities, and the noise accumulates over the full run of $\sim \sqrt{N}$ oracle calls. This threshold does not follow from the definitions and the postulate. Comparability with the scale (29) arises when $2/\sqrt{N} \sim \sigma$, i.e. at $N \sim (2/\sigma)^2 \sim 2 \times 10^{37} \approx 2^{124}$; there the signal step is $2^{-61} \approx 4 \times 10^{-19}$ rad. At smaller N (e.g. $N = 2^{80}$, step $\sim 10^{-12}$ rad) the rotation is orders of magnitude above σ and the mechanism is imperceptible on a single iteration; at larger N the step falls below σ , and dephasing accumulated over $\sim \sqrt{N}$ iterations becomes critical—although in practice the computation time kills it first. This is consistent with the known structural fragility of the algorithm: with constant noise speed on the oracle, Grover’s quadratic speedup is destroyed completely [35].

8 Discussion

8.1 Temporal structure of proper-time noise

The relation (1) and the postulate specify only the single-interval variance δt ; multi-time covariance, the noise spectrum, and the spin-echo response are not fixed by them. The law $\text{Var}[\delta t(t)] \propto t^{2/3}$ follows from (1), but does *not* by itself determine the full stochastic process. If Gaussianity of the full process, self-similarity, and stationary increments are additionally imposed, the natural completion is fractional Brownian motion with Hurst exponent $H = 1/3$ [36]. In that completion, disjoint increments are anticorrelated, the low-frequency spectrum of fractional Gaussian noise behaves as $S(f) \propto |f|^{1/3}$, and echo requires an explicit calculation with the corresponding filter function rather than an automatic cancellation of a quasistatic phase. These spectral properties follow from an additional temporal postulate, not from the relation (1) and the postulate as presently stated.

8.2 Limitations and model systematics

The central prediction $\chi \propto m^2 t^{2/3}$ follows from the relation (1) and the postulate, but the overall coefficient remains incompletely fixed. The number ξ in (1) is specified only as $\mathcal{O}(1)$; interpreting δt as a single-interval variance and choosing the effective superposition time t (e.g. L/v versus $2L/v$ in a Talbot–Lau interferometer; see § 6.3) enter the $\mathcal{O}(1)$ uncertainty of the predicted visibility on equal footing with ξ . When confronting a single mass, the observed visibility decomposes as $V_{\text{obs}} = R_{\text{ord}} V_{\text{QM}} e^{-\chi}$; the product $R_{\text{ord}} e^{-\chi}$ is degenerate, and without varying m or t one cannot uniquely separate the BDPT contribution from underestimated conventional contrast losses. The conditional bound at fixed $R_{\text{ord}} = 0.78$ is therefore weaker than the estimate at $R_{\text{ord}} = 1$, which attributes the entire deficit to the mechanism. A statistically robust limit on ξ will require a joint likelihood analysis of the raw scans, the quantum-curve parameters, and all conventional loss channels, not aggregated visibility values alone.

8.3 Relation to objective collapse

A natural question is how BDPT relates to objective collapse if it extinguishes the relative phase between branches. Two levels must be kept apart. For *fixed* realizations $\delta t_a, \delta t_b$ the state

$$|\psi\rangle = c_a|a\rangle + c_b e^{i\Delta\varphi}|b\rangle \quad (30)$$

remains *pure*; $\Delta\varphi$ is given by (2) and depends on the particular realization of the branch clocks. A mixed state appears only after ensemble averaging over independent $\delta t_a, \delta t_b$:

$$\bar{\rho}_{ab} = \rho_{ab}(0) e^{-\chi}. \quad (31)$$

At $\chi \gg 1$ this yields the effective state

$$\rho \approx |c_a|^2 |a\rangle\langle a| + |c_b|^2 |b\rangle\langle b|, \quad (32)$$

without coherent terms $c_a c_b^* |a\rangle\langle b| + \text{h.c.}$ The loss of phase and passage to a mixed state is not reducible to the observer’s incomplete knowledge of the phase: by the postulate the noise is an *ontic* independent shift of proper time per branch, not the recording of which-path information in an environment.

In the ensemble reading (§ 1) each branch is a genuine variant of the system and $|c_i|^2$ are the variant weights. For $\chi \gg 1$ there is no observable interference between variants; the ensemble mixture with the same weights is operationally indistinguishable from a statistical description of outcomes with probabilities $|c_i|^2$. The definite outcome remains unknown until registration, but coherent superposition of alternatives as a physical resource disappears *objectively*—before and independently of readout—over a time $\lesssim t_{\text{dec}}$. In this sense the macroscopic limit of BDPT yields objective loss of inter-branch interference and effective branch superselection at $\chi \gg 1$, but does *not* select a single realization and does not replace dynamical collapse.

The mechanism differs from dynamical collapse models (Diósi–Penrose and others) in its channel: the populations $|c_i|^2$ are not redistributed by a jump and there is no spontaneous emission [18]; it is pure rest-phase dephasing. The destruction of inter-branch coherence and the passage to an ensemble mixture is the link that *resembles* objectivist programmes, but formally follows only from the ensemble ontology, independent branch clocks, and the Károlyházy relation for δt , without an additional nonlinear term in the state equation. Schrödinger’s cat at $m \gg m_*$ illustrates the limit: the amplitudinal record “alive/dead” remains, but interference between branches vanishes over $t_{\text{dec}} \ll t_C$; what remains is a statistical mixture of variants with weights $|c_i|^2$, not the selection of a single outcome before measurement.

8.4 Distinction from other decoherence mechanisms

The mechanism must be distinguished from nearby sources of interference loss. The closest formal relative is Milburn’s intrinsic decoherence [37]; Adler’s simplest equation coincides with it [38, 39]: stochastic unitary phase steps generated by the Hamiltonian yield fundamental dephasing specifically in the energy eigenbasis of H , with populations conserved and a rate that grows with $|E - E'|$. BDPT shares with that programme the idea of an internal (not environment-induced) loss of coherence from uncertain time evolution, but is not reducible to it. In Milburn’s model decoherence is tied to the eigenstates of the full H and for a free particle produces anomalous dispersion (an effective channel with an operator $\propto p^2$); at $E = E'$ there is no dephasing, so equal-rest-energy which-path arms alone yield no signal. The same vanishing at equal energies, and the factor $t_P^{4/3} t^{2/3}$ in front of the square of a frequency, were already obtained by Gambini, Porto and Pullin from the Salecker–Wigner and Ng–van Dam clock bound: the off-diagonal element decays as $\exp(-\omega_{nm}^2 t_P^{4/3} t^{2/3})$, where ω_{nm} is the difference of eigenvalues of a single Hamiltonian [40, 41]. In Equation (5), $mc^2/\hbar$ stands in place of ω_{nm} by the definitions and the postulate; the numerical coefficient differs by the factor $\xi^{2/3}$. In BDPT the noise attaches to discrete branches by the definitions, and δt is set by the Károlyházy relation with $\chi \propto m^2 t^{2/3}$, and the postulate of independent clocks turn the common Compton phase into a random *relative* contribution without force or momentum diffusion. In quantum-clock interference, a deterministic or potential-induced proper-time difference records which-path information through internal degrees of freedom or an external gravitational gradient [14, 15]; BDPT is not reducible to such unitary readout and does not require a potential difference, but introduces a stochastic Planckian uncertainty δt with independent $\delta t_a, \delta t_b$. An environment with momentum transfer (gas, thermal radiation, photon scattering) creates position-dependent noise, force, and momentum diffusion and is usually suppressed by improving the vacuum or cooling; the characteristic dependence on Δx is described by localization factors [26] (see § 6.5). The closest construction of this kind is Diósi’s local-time uncertainty [39]: the rest energy $c^2 f(r)$ couples to random shifts of local time with correlator $\propto G\hbar/(|r - r'|c^4)$. With that correlator c cancels, and the master equation coincides with the Diósi–Penrose equation, whose rate is set by the gravitational self-energy. In the present model the shifts are independent across branches and their variance is fixed by the relation (1), so c remains and the rate is set by mc^2 . Dynamical Diósi–Penrose collapse models predict population redistribution and spontaneous emission [18], to which the present model is *transparent*: it is pure rest-phase dephasing without additional channels. Spatially attached implementations of Planckian uncertainty (correlated shear noise, external metric fluctuations) are constrained by devices such as the Holometer and LIGO-like detectors [21]; BDPT attaches to discrete branches, not to points, and does not act on a massless carrier. Finally, in standard quantum mechanics the Compton phase cancels in a closed interferometer [16]; the present model does not settle that debate but postulates that, with independent branch clocks, the rest phase mc^2 becomes a random *relative* contribution (see § 3)—a separate and falsifiable premise.

9 Conclusion

Tying the Károlyházy fluctuation to proper time and postulating independent branch clocks yields a decoherence law (for which-path, $\xi = \mathcal{O}(1)$)

$$V/V_0 = \exp \left[- \left(\frac{mc^2}{\hbar} \right)^2 (\xi t_P^2 t)^{2/3} \right], \quad (33)$$

where interference involves the *difference* of branch phase contributions and mc^2 is the energy of the split carrier. The scaling $m^2 t^{2/3}$ is immune to electromagnetic shielding; sodium interferometry [22], attributing the entire contrast deficit to the mechanism, gives $\xi \lesssim 1.1$ at the

one-sided normal quantile; $\xi = \mathcal{O}(1)$ is not excluded (conditionally, with the conventional-loss normalization 0.78 , $\xi \lesssim 0.32$). For $\xi = \mathcal{O}(1)$ the characteristic scale $m_* = m_P/\sqrt{\xi}$ is of the order of the Planck mass; for $m \gg m_*$ decoherence occurs over $t_{\text{dec}} \ll t_C$, below the Károlyházy distinguishability limit for proper time. The prediction at $\xi = \mathcal{O}(1)$ is falsifiable by a test at ~ 1 MDa and $v \approx 25$ m/s: either fringes of the standard quantum model, or suppression with the $m^2 t^{2/3}$ scaling.

References

- [1] L. E. Ballentine. The statistical interpretation of quantum mechanics. *Reviews of Modern Physics*, 42(4):358–381, 1970.
- [2] D. Home and M. A. B. Whitaker. Ensemble interpretations of quantum mechanics. A modern perspective. *Physics Reports*, 210(4):223–317, 1992.
- [3] D. N. Page and W. K. Wootters. Evolution without evolution: Dynamics described by stationary observables. *Physical Review D*, 27:2885–2892, 1983.
- [4] R. Penrose. On gravity’s role in quantum state reduction. *General Relativity and Gravitation*, 28(5):581–600, 1996.
- [5] J. I. Korsbakken, K. B. Whaley, J. DuBois, and J. I. Cirac. Measurement-based measure of the size of macroscopic quantum superpositions. *Physical Review A*, 75:042106, 2007.
- [6] F. Károlyházy. Gravitation and quantum mechanics of macroscopic objects. *Il Nuovo Cimento A*, 42:390–402, 1966.
- [7] H. Salecker and E. P. Wigner. Quantum limitations of the measurement of space-time distances. *Physical Review*, 109:571–577, 1958.
- [8] Y. J. Ng and H. van Dam. Limitation to quantum measurements of space-time distances. *Annals of the New York Academy of Sciences*, 755:579–584, 1995.
- [9] Y. J. Ng and H. van Dam. Limit to space-time measurement. *Modern Physics Letters A*, 9:335–340, 1994.
- [10] Y. J. Ng. Spacetime foam, holographic principle, and black hole quantum computers. *International Journal of Modern Physics A*, 20:1328–1335, 2005.
- [11] Paolo M. Bonifacio, Charles H.-T. Wang, J. Tito Mendonça, and Robert Bingham. De-phasing of a non-relativistic quantum particle due to a conformally fluctuating spacetime. *Classical and Quantum Gravity*, 26:145013, 2009.
- [12] Heinz-Peter Breuer, Ertan Göklü, and Claus Lämmerzahl. Metric fluctuations and decoherence. *Classical and Quantum Gravity*, 26:105012, 2009.
- [13] F. Kiałka, Y. Y. Fein, S. Pedalino, S. Gerlich, and M. Arndt. A roadmap for universal high-mass matter-wave interferometry. *AVS Quantum Science*, 4:020502, 2022.
- [14] M. Zych, F. Costa, I. Pikovski, and Č. Brukner. Quantum interferometric visibility as a witness of general relativistic proper time. *Nature Communications*, 2:505, 2011.
- [15] I. Pikovski, M. Zych, F. Costa, and Č. Brukner. Universal decoherence due to gravitational time dilation. *Nature Physics*, 11:668–672, 2015.

- [16] P. Wolf, L. Blanchet, C. J. Bordé, S. Reynaud, C. Salomon, and C. Cohen-Tannoudji. Does an atom interferometer test the gravitational redshift at the Compton frequency? *Classical and Quantum Gravity*, 28:145017, 2011.
- [17] S.-Y. Lan, P.-C. Kuan, B. Estey, D. English, J. M. Brown, M. A. Hohensee, and H. Müller. A clock directly linking time to a particle’s mass. *Science*, 339(6119):554–557, 2013.
- [18] S. Donadi et al. Underground test of gravity-related wave-function collapse. *Nature Physics*, 17:74–78, 2021.
- [19] I. J. Arnquist et al. Search for spontaneous radiation from wave-function collapse in the Majorana Demonstrator. *Physical Review Letters*, 129:080401, 2022.
- [20] K. Piscicchia et al. X-ray emission from atomic systems can distinguish between prevailing dynamical wave-function collapse models. *Physical Review Letters*, 132:250203, 2024.
- [21] A. S. Chou et al. Interferometric constraints on quantum geometrical shear noise correlations. *Classical and Quantum Gravity*, 34:165005, 2017.
- [22] S. Pedalino, B. E. Ramírez-Galindo, R. Ferstl, K. Hornberger, M. Arndt, and S. Gerlich. Probing quantum mechanics with nanoparticle matter-wave interferometry. *Nature*, 649:866–870, 2026.
- [23] S. Pedalino, B. E. Ramírez-Galindo, R. Ferstl, K. Hornberger, M. Arndt, and S. Gerlich. Probing quantum mechanics using nanoparticle Schrödinger cats, 2025.
- [24] S. Nimmrichter and K. Hornberger. Macroscopicity of mechanical quantum superposition states. *Physical Review Letters*, 110:160403, 2013.
- [25] Sebastian Pedalino, Bruno Eduardo Ramírez Galindo, Richard Ferstl, Klaus Hornberger, Markus Arndt, and Stefan Gerlich. Probing quantum mechanics with nanoparticle matter-wave interferometry - dataset and code, 2025.
- [26] K. Hornberger, J. E. Sipe, and M. Arndt. Theory of decoherence in a matter wave Talbot–Lau interferometer. *Physical Review A*, 70:053608, 2004.
- [27] J. I. Korsbakken, F. K. Wilhelm, and K. B. Whaley. Electronic structure of superposition states in flux qubits. *Physica Scripta*, T137:014022, 2009.
- [28] J. I. Korsbakken, F. K. Wilhelm, and K. B. Whaley. The size of macroscopic superposition states in flux qubits. *EPL*, 89:30003, 2010.
- [29] Y.-A. Chen, X.-H. Bao, Z.-S. Yuan, S. Chen, B. Zhao, and J.-W. Pan. Heralded generation of an atomic NOON state. *Physical Review Letters*, 104:043601, 2010.
- [30] N. Margolus and L. B. Levitin. The maximum speed of dynamical evolution. *Physica D*, 120:188–195, 1998.
- [31] O. Milul, B. Guttel, U. Goldblatt, S. Hazanov, L. M. Joshi, D. Chausovsky, N. Kahn, E. Çiftyürek, F. Lafont, and S. Rosenblum. Superconducting cavity qubit with tens of milliseconds single-photon coherence time. *PRX Quantum*, 4:030336, 2023.
- [32] D. Coppersmith. An approximate Fourier transform useful in quantum factoring, 1994. IBM Research Report RC 19642.
- [33] P. W. Shor. Polynomial-time algorithms for prime factorization and discrete logarithms on a quantum computer. *SIAM Journal on Computing*, 26(5):1484–1509, 1997.

- [34] L. K. Grover. A fast quantum mechanical algorithm for database search. In *Proceedings of the 28th Annual ACM Symposium on Theory of Computing (STOC)*, pages 212–219, 1996.
- [35] O. Regev and L. Schiff. Impossibility of a quantum speed-up with a faulty oracle. In *Automata, Languages and Programming (ICALP 2008), Lecture Notes in Computer Science*, volume 5125, pages 773–781, 2008.
- [36] B. B. Mandelbrot and J. W. Van Ness. Fractional Brownian motions, fractional noises and applications. *SIAM Review*, 10(4):422–437, 1968.
- [37] G. J. Milburn. Intrinsic decoherence in quantum mechanics. *Physical Review A*, 44(9):5401–5411, 1991.
- [38] S. L. Adler. *Quantum Theory as an Emergent Phenomenon*. Cambridge University Press, Cambridge, 2004.
- [39] L. Diósi. Intrinsic time-uncertainties and decoherence: comparison of 4 models. *Brazilian Journal of Physics*, 35(2A):260–265, 2005.
- [40] R. Gambini, R. A. Porto, and J. Pullin. Realistic clocks, universal decoherence, and the black hole information paradox. *Physical Review Letters*, 93:240401, 2004.
- [41] R. Gambini, R. A. Porto, and J. Pullin. Fundamental decoherence from quantum gravity: a pedagogical review. *General Relativity and Gravitation*, 39:1143–1156, 2007.

Statements and Declarations

Funding. The author declares that no funds, grants, or other support were received during the preparation of this manuscript.

Competing interests. The author has no relevant financial or non-financial interests to disclose.

Author contributions. Oleg Ponfilenok conceived the model, performed the calculations, and wrote the manuscript.

Data availability. The numerical bound reported in this article uses the publicly archived scans and analysis code of Pedalino et al. [25], available at <https://doi.org/10.5281/zenodo.17502163>. No new datasets were generated.

Use of generative AI. A large language model was used to assist in checking algebraic and numerical calculations. The author verified all results and takes full responsibility for the content of the manuscript.