Collapse Threshold from a Computational Limit

Oleg Ponfilenok
Independent researcher
Corresponding author: ponfil@gmail.com

Abstract

We propose a modernization of quantum mechanics through the introduction of the Collapse Threshold Rule: a branch is removed if it cannot be distinguished experimentally from absence. Objectivation occurs when residual coherence falls below the same threshold $\varepsilon = 1/I_{\text{ost}}$. The threshold is derived from the finite number of binary distinctions within a causal patch: $I_{\text{ost}} = 3{,}5 \times 10^{120}$. The scale of equiprobable branching and the RCS limit are about 400 qubits. The model forbids factorization of RSA-2048. Non-unitarity of trimming and selection permits non-conservation of energy as well as superluminal information transfer. An experimental test on a single 404-qubit processor is proposed.

Keywords: objective collapse; objectivation; operationalism; branch removal; residual coherence; BDPT; einselection; computational limit; Shor's algorithm; superluminal telephone; CMB frame

1 Introduction

Standard quantum mechanics permits unitary decoherence: the reduced entropy of a subsystem grows, global entropy equals zero, the phase is recorded in correlations with the environment and is recoverable (Loschmidt echo [1]). The classical bit is thereby not born *objectively*—interpretation remains virtual.

The Copenhagen interpretation splits a microscopic system that lives in superposition from a classical apparatus, but does not say where the Heisenberg cut lies and does not fix a numerical boundary. Decoherence and einselection select a pointer basis and explain the stability of macroscopic pointers [2], yet by themselves they do not specify at what branching width a superposition remains physically distinguishable. The aim of this work is to locate that cut quantitatively and to work out the consequences.

Standard theory does not set a minimal nonzero value for density-matrix elements and probability weights. Decoherence can decrease coherences without bound, and sequential branching and spatial tails of the wave function create arbitrarily small nonzero weights. If such quantities are treated as independently distinguishable physical data and stored with fixed absolute resolution, the required bit depth for exponential decay grows linearly with time, and for a Gaussian tail quadratically with distance. Without a lower threshold it is unbounded and, given sufficiently long evolution, exceeds any finite information limit of the region, in particular the Bekenstein holographic bound [3]. Mathematically arbitrarily small coefficients are admissible, but their physical storage and distinction at finite information capacity is impossible. Hence arises the idea of an information threshold below which values are physically indistinguishable.

Information that can be obtained and verified experimentally is finite in two independent senses. The first limit is set by quantum uncertainty: incompatible physical quantities cannot be measured simultaneously with arbitrarily high precision. The second sets the maximum number of experiments—questions that nature answers. It is tied to finiteness of the causal patch—the

maximal causally closed region accessible to one observer. The cosmological budget I_{ost} of that region permits only a finite number of operations over its entire future [4, 5, 6, 7]. Therefore infinite information is accessible neither by measurement precision nor by the number of physical operations with which it can be obtained.

Ideas of finite precision and threshold were discussed earlier. Buniy, Hsu, and Zee linked the finiteness of distinguishable states to discreteness of Hilbert space and quantization of its coefficients [8], then allowed exclusion of wave-function components with norm below a discreteness threshold to eliminate "maverick branches" and restore Born's rule [9]. This threshold was not fixed: it depended on characteristic energy or system size and number of subsystems. Ozhigov, in an algorithmic approach to quantum physics, cut off small amplitudes: given an accessible computational resource (algorithm steps or memory cells), amplitudes below the inverse square root of that resource are set to zero, i.e. probabilities below its reciprocal; a fixed amplitude quantum leaves only finitely many nonzero components and is used to describe measurement and recover Born's rule [10]. Srikanth, at finite amplitude precision μ, obtained a memory overflow $\mathcal{M}_{\max} = \mu \cdot 2^{\mu/2}$ bits as entanglement grows: unitarity becomes computationally unstable, and an information transition jumps the system into a less complex factorized state by the Born rule, with the basis from decoherence [11, 12]. Davies compared the cosmological information bound $I_{\text{universe}} \sim 10^{122}$ bits with the number of complex coefficients of a generic n-qubit state and obtained the scale $n \approx \log_2 I_{\text{universe}} \approx 400$ [13]. The number 400 in Davies follows from the hypothesis that the cost of a classical description of the state limits its physical existence. Del Santo and Gisin introduced finite informational precision of physical quantities and discussed collapse-like actualization of undefined digits, but in the context of classical indeterminism rather than quantum branch weights [14]. Gambini, García-Pintos, and Pullin, in the Montevideo interpretation, include the gravitational limit on clock and rod precision: an event occurs when residual interference can no longer be distinguished from a reduction (undecidability, not FAPP) [15]. Harlow and Hayden showed that quantum gedanken experiments with black holes may be computationally inaccessible within available time [16].

The new proposal of this work is the Collapse Threshold Rule, implementing the postulate of limited description: within a causal patch no more than I_{ost} binary distinctions are executable, whence follows the absolute threshold of experimental distinguishability $\varepsilon = 1/I_{\text{ost}}$. The rule acts on the full density matrix of the entangled cluster in the causal patch, in the einselection basis, without a system/environment split. The general rule is removal by indistinguishability: a branch with $p_i < \varepsilon$ is indistinguishable from absence and is removed; a coherence with $C \leq \varepsilon$ is indistinguishable from absence—this is objectivation, and the full matrix is left with a single outcome. The threshold ε sets the resolution limit of the entire region and is not divided among individual carriers as a quota. First a mnemonic estimate of the threshold from causal-patch memory capacity is given, then it is rejected in favor of counting executable distinctions (§ 5).

2 Holographic Estimate of the Threshold

Consider a thought experiment: at the center—an atom emitting a photon of wavelength λ. The photon departs in all directions at once: a spherical superposition of directions, without slits. Around it—a spherical screen of radius R with detectors over the entire surface. The number of distinguishable detection outcomes is the sphere area divided by cell area λ^2 (as the number of microstates):

$$N_{\text{cell}}(R, \lambda) \;=\; \frac{4\pi R^2}{\lambda^2}. \tag{1}$$

Each cell is one individual outcome "photon detected here." In the cell basis the diagonal of the reduced ρ gives probabilities of these outcomes; off-diagonal elements give coherence between directions. Count only outcomes: how many diagonal addresses the sphere has.

Limiting sphere. The maximal radius is the de Sitter horizon R_{dS}, to which the cosmological event horizon asymptotically tends in the future; the minimal wavelength is the thermal length at Planck temperature T_P:

$$R_{\max} \;=\; R_{\mathrm{dS}}, \qquad \lambda_{\min} \;=\; \frac{\hbar c}{k_B T_P} \;=\; \ell_P, \tag{2}$$

where $\ell_P = \sqrt{\hbar G/c^3}$ is the Planck length. The area of the limiting sphere equals $A_{\mathrm{dS}} = 4\pi R_{\mathrm{dS}}^2$. Therefore at resolution area ℓ_P^2 the number of binary cells on it is

$$N_{\mathrm{cell,max}} \;=\; \frac{A_{\mathrm{dS}}}{\ell_P^2} \;=\; \frac{4\pi R_{\mathrm{dS}}^2}{\ell_P^2} \;\approx\; 1{,}3 \times 10^{123}, \tag{3}$$

Assigning one binary detection fact "yes" or "no" to each cell corresponds to $1{,}3\times10^{123}$ bits. The resulting quantity is of the order of holographic capacity [3]: de Sitter entropy [17], expressed in bits, equals

$$\frac{S_{\mathrm{dS}}}{\ln 2} \;=\; \frac{A_{\mathrm{dS}}}{4\ell_P^2 \ln 2} \;=\; \frac{N_{\mathrm{cell,max}}}{4\ln 2} \;\approx\; 4{,}8 \times 10^{122} \text{ bits}. \tag{4}$$

Here S_{dS} denotes dimensionless entropy in nats. The two estimates differ only by the factor $4\ln 2$. The cells form an einselection basis in which the entire reduced ρ lives. The quantity $N_{\mathrm{cell,max}}$ estimates *memory capacity* of the causal patch: how many binary facts can be stored in it. This estimate will be rejected in § 4 in favor of a different count—the number of *executable* binary distinctions I_{ost}. Before that transition it is useful to remember the order of magnitude $1/N_{\mathrm{cell,max}} \sim 10^{-123}$ as a mnemonic scale, but not as the working threshold of the model.

3 Cosmological Computational Limit

The calculation is carried out in the flat ΛCDM model: FRW metric with $\Omega_k = 0$, $\Omega_m + \Omega_\Lambda + \Omega_r = 1$; the radiation contribution in the present epoch is negligible ($\Omega_r < 0{,}01\%$).

Take the Hubble horizon $R = c/H$ and Friedmann's second equation for energy density ρ and pressure p

$$\dot{H} = -\frac{4\pi G}{c^2}(\rho + p). \tag{5}$$

Horizon growth rate:

$$\frac{dR}{dt} = -c\,\frac{\dot{H}}{H^2} = \frac{4\pi G}{c^3}(\rho + p)\,R^2. \tag{6}$$

Then $\rho = \rho_m + \rho_\Lambda$, $p_m = 0$, $p_\Lambda = -\rho_\Lambda$, whence $\rho + p = \rho_m$. Therefore in the adopted approximation entropy change and available operations are determined by non-relativistic matter rest energy; the dark-energy contribution vanishes because $\rho_\Lambda + p_\Lambda = 0$. For a flat universe $\rho_m = \Omega_m \cdot 3c^4/(8\pi G R^2)$, and

$$\frac{dR}{dt} = \frac{3}{2}\,c\,\Omega_m. \tag{7}$$

Horizon entropy $S = \pi k R^2/\ell_P^2$:

$$\frac{dS}{dt} = \frac{2\pi k R}{\ell_P^2}\frac{dR}{dt} = \frac{3\pi k c^4}{\hbar G}\,\Omega_m R. \tag{8}$$

Matter energy inside the horizon $E_m = c^4\Omega_m R/(2G)$. Lloyd's limit [4, 5]

$$\dot{N}_m = \frac{2E_m}{\pi\hbar} = \frac{c^4\Omega_m R}{\pi\hbar G}. \tag{9}$$

Dark energy produces no entropy ($\rho_\Lambda + p_\Lambda = 0$). The ratio

$$\frac{\dot{S}}{k\,\dot{N}_m} = 3\pi^2 \tag{10}$$

— $3\pi^2$ nats (about 42,7 bits) per orthogonal matter operation. Ω_m and R cancel: the relation is holographic and epoch-independent while matter and Λ dominate.

de Sitter entropy is finite; its value is given above (§ 2). The identity $S_{\rm dS} - S = S_{\rm dS}\Omega_m$ is exact: $R_H^2 = R_{\rm dS}^2\Omega_\Lambda$ for a flat universe without radiation. Remaining orthogonal operations

$$I_{\rm ost}(t) = \int_t^\infty \dot{N}_m\,dt' = \frac{S_{\rm dS} - S}{3\pi^2} = \frac{S_{\rm dS}\Omega_m}{3\pi^2} \approx 3{,}5 \times 10^{120} \tag{11}$$

The value for the present epoch is given here. The integral is finite because the horizon asymptotically tends to $R_{\rm dS}$. In homogeneous Friedmann–Robertson–Walker (FRW) cosmology the quantity $I_{\rm ost}(t)$ is the same for comoving observers at one cosmological time.

As Ω_m decreases, $I_{\rm ost}(t)$ decreases: at recombination $I_{\rm ost} \approx 1{,}1\times10^{121}$, now $I_{\rm ost} \approx 3{,}5\times10^{120}$, and in the asymptotic future $I_{\rm ost}(t) \to 0$. At laboratory timescales the change in Ω_m can be neglected.

4 Computational Estimate of the Threshold

In quantum Darwinism, objectivity arises when information about the system state is multiply reproduced in independent fragments of the environment [18]; this picture develops Wheeler's "It from Bit" [19]. Binary distinction here is understood in Wheeler's sense—a yes/no answer to a posed question, i.e. an act rather than a stored record.

The quantities in § 2 and § 3 measure different things. $N_{\rm cell,max}$ is region memory capacity: how many binary facts fit in it. $I_{\rm ost}$ is the number of orthogonal operations: how many binary distinctions are executable in it. The ratio $12\pi^2/\Omega_m \approx 376$ is the difference between two counts, not a correction to one. The same gap between horizon entropy and the processable budget was noted by Krauss and Starkman [7]: their limit on energy harvestable in de Sitter gives the order 10^{120}, but not an identity for $I_{\rm ost}$. Hereafter the count $I_{\rm ost}$ is adopted: physically a branch exists insofar as it can be distinguished from absence, not insofar as it can be stored.

Matching the operation remainder to the number of binary cells on the limiting sphere, we obtain

$$\frac{N_{\rm cell,max}}{I_{\rm ost}} = \frac{12\pi^2}{\Omega_m} \sim 4 \times 10^2.$$

Therefore the mnemonic scale $1/N_{\rm cell,max} \sim 10^{-123}$ overestimates resolvability: memory is still sufficient, but operations to distinguish all addresses are lacking. On a logarithmic scale the same factor lowers the recorded limit from $\log_2 N_{\rm cell,max} \approx 409$ to $N_* = \log_2 I_{\rm ost} \approx 400{,}4$, i.e. by about 8,6 qubits. Davies already noted the same order 400 [13], comparing 2^n coefficients of an entangled state with $I_{\rm universe} \sim 10^{122}$; that is a count of Hilbert-space dimension and memory capacity, not of executable distinctions, and it supplies no trimming rule. Working resolution is limited by the number of executable distinctions, not memory capacity.

The same limit is a threshold on distinguishable probability. A fair coin in a series of N_* independent tosses assigns a given sequence weight $2^{-N_*} \approx \varepsilon$; as *one* equiprobable alternative among 2^N strings at $N > N_*$ such a run lies below the threshold and is indistinguishable from absence. This is not a ban on a long classical record. A 500-bit file can be stored and copied: the bits are written sequentially, each with weight of order unity, not drawn at once from 2^{500} equal strings. Possessing a prespecified string is not the same as obtaining it by randomly enumerating all bits. As in bitcoin mining: writing a message that begins with fifty zeros is trivial; obtaining the same prefix as a random hash output takes of order 2^{50} trials. The expected particular

500-bit file under random search carries weight $2^{-500} < \varepsilon$ and is indistinguishable, whereas the same string, written bit by bit, remains ordinary memory.

The quantity I_{ost} is not a consumable quota distributed among devices, but the total measurement budget of the causal patch. The operational meaning of ε is that over I_{ost} measurements one can accumulate a distinguishable signal of order unity; per operation

$$\varepsilon = \frac{1}{I_{\text{ost}}} \approx 3 \times 10^{-121}. \tag{12}$$

If an einselection object's contribution to this total signal is weaker than ε, no protocol within the patch distinguishes it. In the Collapse Threshold Rule (§ 5) this budget is realized for one branch or one coherence; trace distance [20] describes the shift of outcome distribution in one measurement, but the threshold is set by accumulated statistics over I_{ost} operations. Therefore ε characterizes the resolving power of the entire region, has the same value for every entangled system, and does not grow with the number of devices. It is a property of the causal patch, not the laboratory apparatus: the local environment determines which branches and coherences are tested, but not the value of ε.

5 Collapse Threshold Rule

We put forward the hypothesis and propose the postulate that a description of physical state cannot contain more independently distinguishable elements than I_{ost} binary distinctions (§ 4). Hence the general rule of removal by indistinguishability: an element of an einselection object to which no distinction is allocated is physically indistinguishable from absence and is removed. For a branch this is projection onto the sector complement. For coherence the same rule yields objectivation: indistinguishable interference is not retained in the description, and the full matrix is left with a single outcome. The rule takes the full density matrix of all entangled particles and acts on it as a whole, without a system/environment split.

We introduce an integer micromodel specifying nonnegative integers $n_i \in \mathbb{Z}_{\geq 0}$, $\sum_i n_i = I_{\text{ost}}$; weights live on the grid n_i/I_{ost}. Between zero and $\varepsilon = 1/I_{\text{ost}}$ there is no admissible value, so excess is removed by deletion rather than coarse-graining all weights. The full dynamical model is discussed in § 12.

The Collapse Threshold Rule implements this postulate for einselection objects within a causal patch with resolving power ε. By "description" we mean not the unitary circuit that prepared the state, but what experiment can afterwards distinguish: Born weights of branches and subsystem correlations readable from measurement statistics (§ 4).

Branch removal. A branch is one outcome in the einselection basis with diagonal weight $p_i = \rho_{ii}$; off-diagonal elements describe interference between branches. Physical interaction with the environment defines orthogonal sectors P_i, not an arbitrary matrix representation. The Collapse Threshold Rule applies to the entire entangled system within which these records are linked.

The Collapse Threshold Rule tests one einselection object: one branch or the coherence of that branch with its complement. Let P_i be the sector projector, $p_i = \operatorname{Tr}(P_i\rho)$ the Born weight of the branch. Elementary branch removal and elementary Lüders dephasing [21] take the form

$$\mathcal{C}_i^{\text{del}}(\rho) = \frac{(I-P_i)\rho(I-P_i)}{\operatorname{Tr}\big((I-P_i)\rho\big)}, \qquad \mathcal{C}_i^{\text{ph}}(\rho) = P_i\rho P_i + (I-P_i)\rho(I-P_i). \tag{13}$$

Removing the corresponding rows and columns leaves the principal submatrix of a positive semidefinite matrix ρ, and Lüders dephasing is completely positive. Therefore Hermiticity, positivity, and normalization are preserved [22, 21].

An object is removed if it cannot be distinguished from absence within the budget of I_{ost} measurements.

Recorded branch. Count how many times branch i occurs in a series of I_{ost} independent repetitions. The expected number of hits equals $I_{\text{ost}}p_i$; a distinguishable signal of order unity requires $I_{\text{ost}}p_i \gtrsim 1$, whence

$$p_i < \varepsilon = \frac{1}{I_{\text{ost}}}. \tag{14}$$

Coherent branch. The same branch may enter superposition with its complement and be tested by interferometry: the signal goes through amplitude $\sqrt{p_i}$, not frequency p_i. Over I_{ost} repetitions of the same interferometric measurement amplitudes add as $\sqrt{I_{\text{ost}}p_i}$; the requirement $\sqrt{I_{\text{ost}}p_i} \gtrsim 1$ is equivalent to $I_{\text{ost}}p_i \gtrsim 1$ and again yields $p_i < \varepsilon$. Interferometry is more sensitive in a single measurement—in this sense trace distance to $\mathcal{C}_i(\rho)$ is of order $\sqrt{p_i}$, not p_i [20]—but no second operation budget is allocated: each repetition consumes the same unit from I_{ost}. Therefore the condition $p_i < \varepsilon^2$ that would follow from a single measurement ($\sqrt{p_i} < \varepsilon$) applies only to a measurement without statistical accumulation; with the full causal-patch budget both cases reduce to one invisibility condition $p_i < \varepsilon$.

Physically distinguishable objects are measurement outcomes: branch probabilities and interference between them, recoverable from experiment statistics, including quantum tomography. The density matrix is only a description of this operationally accessible information, not an independent physical object; the observed state is a branch. The Collapse Threshold Rule is deliberately applied to individual branches, not to distinguishability of the full description: otherwise any joint removal of invisible branches would be forbidden as soon as their total weight becomes observable, and the model would add no new predictions, remaining in principle unfalsifiable. Therefore the Collapse Threshold Rule cuts off einselection objects indistinguishable by outcome statistics and does not require a small change of the entire ρ in coordinate representation: joint removal of many invisible branches and coherences can change the matrix by order unity, although each such branch or coherence individually is experimentally indistinguishable.

Removal of invisible branches generally leaves several live sectors with $p_i \geq \varepsilon$. That is not a single outcome. Residual superposition of the full matrix in the einselection basis is

$$\Delta_P(\rho) = \sum_i P_i \rho P_i, \qquad C(\rho) = \frac{1}{2}\big\|\rho - \Delta_P(\rho)\big\|_1. \tag{15}$$

For two orthogonal branches with weights p and $1-p$ one has $C = 2\sqrt{p(1-p)}$; for a Bell pair $C = 1$.

Irreversibility of branch removal. Since $I_{\text{ost}}(t)$ decreases monotonically, $\dot{\varepsilon} > 0$: the threshold only rises with time. A branch removed at an earlier threshold value remains below it and by the Collapse Threshold Rule is not restored. Unitary evolution may subsequently create a branch with the same quantum numbers, but it will be a new branch, not restoration of the removed one.

Renormalization. After identifying

$$S = \big\{i : \ p_i < \varepsilon\big\} \tag{16}$$

invisible sectors are removed and the remainder is renormalized if its weight is positive. Renormalization covers all surviving branches of the entire entangled system. Independent cutoff of matrix cells by $|\rho_{ij}| < \varepsilon$ is not applied: it may violate positivity [22], and visible interference of one sector pair may sum from many cells within blocks.

Let $Q = I - \sum_{i\in S} P_i$ be the projector onto the surviving part of the state. Then the Collapse Threshold Rule acts on the full density matrix of the entangled system in one step:

$$\mathcal{N}(\rho) = \frac{Q\rho Q}{\operatorname{Tr}(Q\rho Q)}. \tag{17}$$

For K equiprobable alternatives each weight equals $1/K$. Since both frequency counting and interferometry require $p_i \geq \varepsilon$ for visibility, the condition $p_i < \varepsilon$ yields the same ceiling

$$K \lesssim I_{\text{ost}}, \qquad N_* \approx \log_2 I_{\text{ost}} \approx 400{,}4. \tag{18}$$

In further estimates it is convenient to use the round reference value 400. This estimate refers to one equiprobable alternative within one connected system, not to a factorized product or sequential classical recording. How to count branches for a product, GHZ, a Haar state, and RCS is in § 6.

Dephasing. Unitary entanglement with the environment moves inter-branch blocks into correlations and does not by itself reduce C of the full ρ. For coherence of live sectors to fall to ε, dephasing of the full matrix is required; the mechanism may vary. As an example we take the hypothesis of independent proper-time clocks (BDPT) [23]: blocks $P_a \rho P_b$ decay as $e^{-\chi}$, $\chi \propto (Mc^2)^2 t^{2/3}$, where M is the rest mass of branch matter; radiation and Λ energy do not enter the phase. This is a separate hypothesis, not a consequence of I_{ost}.

Objectivation. Objectivation is the occurrence of a single outcome in the full ρ: a pure state of one branch. It is not an improper mixture, in which a subsystem looks mixed because of the environmental trace while the global superposition still lives, nor a proper classical mixture of several live branches without interference. The quantity $C(\rho)$ is the residual coherence of the full matrix in the einselection basis: a single trace distance to the mixture $\Delta_P(\rho)$, not the coherence of one branch; it includes all blocks $P_i \rho P_j$ with $i \neq j$. If more than one sector is live and $C(\rho) \leq \varepsilon$,

$$\mathcal{C}^{\text{sel}}(\rho) = \frac{P_k \rho P_k}{p_k}, \qquad \Pr(k) = p_k = \operatorname{Tr}(P_k \rho). \tag{19}$$

Trimming is not required for this: when all branches are visible, $\mathcal{N}$ is the identity, and selection is produced by dephasing. If trimming already left one branch, objectivation is trivial. While $C > \varepsilon$ and several branches remain visible, the description carries them all: the classical outcome is not yet singled out.

Degenerate case. If *all* branches are invisible, joint removal would zero ρ. Objectivation is then the same jump $\mathcal{C}^{\text{sel}}$, with conditional probability

$$\Pr(i \mid S) = \frac{p_i}{\sum_{k \in S} p_k}. \tag{20}$$

The trigger is absence of a visible remainder, not $C \leq \varepsilon$. This case arises for a flat distribution and determines the fate of Shor's algorithm (§ 8).

6 Examples of the Rule

Spatial tail of the $1s$ state. For a Coulomb $1s$ state the integral weight outside radius r equals

$$w(r) = e^{-2x}(1 + 2x + 2x^2), \qquad x = r/a_0. \tag{21}$$

The quantity w is the Born probability to find the electron outside a sphere of radius r in a position measurement. However before such a measurement the state has the form

$$|\psi\rangle = \sqrt{1 - w}\,|\text{in}\rangle + \sqrt{w}\,|\text{out}\rangle. \tag{22}$$

Removing the outer tail changes the positional outcome probability by w. In a coherent state, interferometry over I_{ost} repetitions distinguishes amplitude $\sqrt{w}$ when $I_{\text{ost}} w \gtrsim 1$, i.e. at the

same $w \gtrsim \varepsilon$. Therefore for both coherent and already recorded tails trimming requires $w < \varepsilon$; this gives $r \gtrsim 7{,}6$ nm for a hydrogen atom in the ground state. This is removal of a small weight, including before environment recording. A single outcome "inside or outside" occurs only if both sectors survive and $C \leq \varepsilon$, or if the outer tail is already cut and one branch remains.

Factorized product. For factorized $|+\rangle^{\otimes 400}$ each qubit gives two outcomes of weight 1/2, altogether $400 \times 2 = 800$—far below I_{ost}. Branch removal does not arise because each factor carries weight $1/2 \gg \varepsilon$. The weight of a particular computational string of all 400 qubits, e.g. $|0\ldots0\rangle$, equals 2^{-400}, but that is a product of independent weights 1/2, not the weight of one branch.

GHZ. For GHZ all 400 qubits are entangled, but the global state is a superposition of only two strings, $|0\ldots0\rangle$ and $|1\ldots1\rangle$, each with weight 1/2; one cannot sum over qubits. Branch removal does not arise because both live branches carry weight $1/2 \gg \varepsilon$.

Haar state. In a Haar state in dimension $D = 2^N$ there is no factorization: a particular computational string, e.g. $|0\ldots0\rangle$, is already a branch of typical weight $p_i \sim 2^{-N}$, and simultaneously of order 2^{N-1} such branches are distinguishable. At $N \gtrsim 400$ this exceeds I_{ost}, and the Collapse Threshold Rule removes almost all strings with $p_i < \varepsilon$; it is here that the ceiling $N_* \approx 400$ applies, not in factorized $|+\rangle^{\otimes N}$ nor in GHZ.

RCS. Random circuit sampling (RCS) [24, 25] prepares a state close to Haar, so the same ceiling applies; the removal front is in § 8.

Effective number of branches. We define the effective number of branches of an entangled system via the inverse participation ratio (IPR) [26, 27]:

$$K_{\text{eff}} \;=\; \frac{1}{\sum_i p_i^2}. \tag{23}$$

Normalization is $\sum_i p_i = 1$; the sum of squares equals unity only if a single branch is live. For K equal weights $p_i = 1/K$ one has $\sum_i p_i^2 = 1/K$ and $K_{\text{eff}} = K$, so the ceiling $K \lesssim I_{\text{ost}}$ may saturate. For a Haar state in dimension $D = 2^N$ the situation is different:

$$\mathbb{E}\left[\sum_i p_i^2\right] = \frac{2}{D+1}, \qquad K_{\text{eff}} \approx \frac{D+1}{2} \approx 2^{N-1}. \tag{24}$$

The last equality uses IPR concentration at large D; in general the mean of the inverse is not the inverse of the mean. After removing branches with $p_i < \varepsilon$ surviving weights satisfy $p_i \geq \varepsilon$ and $K_{\text{eff}} \leq I_{\text{ost}}$. If $\rho = \bigotimes_a \rho_a$, the number of independently distinguishable elements equals $\sum_a K_{\text{eff}}(\rho_a)$; for non-factorizable ρ—$K_{\text{eff}}(\rho)$. Condition: $\sum_a K_{\text{eff}}(\rho_a) \leq I_{\text{ost}}$. Hence

$$\log_2 K_{\text{eff}} \leq N_* \approx 400{,}4. \tag{25}$$

For the product above the sum over factors is 800, for GHZ $K_{\text{eff}} = 2$, for a Haar state and RCS $K_{\text{eff}} \approx 2^{N-1}$. As coupling between initially independent systems strengthens, the spectrum of reduced ρ and K_{eff} change continuously; at zero coupling the Collapse Threshold Rule applies to each system separately.

7 Non-conservation of Energy at Collapse

We denote by t_{coll} the duration of the transition between two physical descriptions at trimming $\mathcal{N}$ (removal of branches with $p_i < \varepsilon$) or selection $\mathcal{C}^{\mathrm{sel}}$ (a single outcome; § 5).

The jump cannot be localized in time more sharply than its duration, hence the uncertainty of its moment is $\Delta t \sim t_{\mathrm{coll}}$. The uncertainty relation for the act is $\Delta E_{\mathrm{unc}}\, \Delta t \gtrsim \hbar$. In a thermal environment interactions decay on a timescale of order $\hbar/k_B T$, including entanglement on which coherence depends; hence

$$t_{\mathrm{coll}} \lesssim \frac{\hbar}{k_B T}. \tag{26}$$

At 300 K this is ~ 25 fs.

The jump need not preserve unitary norm: before renormalization the total weight of removed branches may differ from unity. If energy non-conservation does not exceed ΔE_{unc}, then

$$|\Delta E| \lesssim \frac{\hbar}{t_{\mathrm{coll}}}. \tag{27}$$

This is an upper estimate via the same act uncertainty, not a direct derivation of $\Delta E\, \Delta t \lesssim \hbar$; without asserting that the corresponding energy is necessarily emitted or absorbed by the environment.

Branch-trimming energy. Mean state energy is $\langle H \rangle = \mathrm{Tr}(H\rho)$. After removing invisible sectors $S = \{i : p_i < \varepsilon\}$ we have $\mathcal{N}(\rho) = Q\rho Q/\,\mathrm{Tr}(Q\rho Q)$, and the energy shift is defined as

$$\Delta E = \mathrm{Tr}(H\mathcal{N}(\rho)) - \mathrm{Tr}(H\rho). \tag{28}$$

This is not a new assumption but a consequence of the renormalization rule (§ 5). Let sectors be approximately energy eigensectors, $[H, P_i] \approx 0$, and $E_i = \mathrm{Tr}(HP_i\rho P_i)/p_i$ the mean energy of branch i. Then $\mathrm{Tr}(H\rho) = \sum_i p_i E_i \equiv \langle E \rangle$, $\mathrm{Tr}(H\mathcal{N}(\rho)) = \langle E \rangle_{\bar{S}}$ and $\mathrm{Tr}(HQ\rho Q) = \sum_{i \notin S} p_i E_i$, where $\bar{S}$ is the complement of S. Denoting $W = \sum_{i \in S} p_i$, $\langle E \rangle_S = (\sum_{i \in S} p_i E_i)/W$ and $\langle E \rangle_{\bar{S}} = (\sum_{i \notin S} p_i E_i)/(1 - W)$,

$$\Delta E = W\big(\langle E \rangle_{\bar{S}} - \langle E \rangle_S\big) = \langle E \rangle_{\bar{S}} - \langle E \rangle. \tag{29}$$

The shift is finite for any $W < 1$ and does not exceed the spread of energies over branches: $|\Delta E| \le W|\langle E \rangle_{\bar{S}} - \langle E \rangle_S|$. Under partial trimming it can be large if removed and surviving branches systematically differ in E_i. Selection $\mathcal{C}^{\mathrm{sel}}$ gives $E_k - \langle E \rangle$ in a single shot, zero on average over Born. In the degenerate case mean energy after the act equals $\sum_i p_i E_i = \langle E \rangle$, i.e. $\langle \Delta E \rangle = 0$ identically—there is no systematic sink or source by construction; in a single act only fluctuations of order ordinary for the system remain. Where energy goes at $W < 1$ and whether collapse is compatible with balance remains an open question (§ 12).

8 Threshold Collapse in Quantum Computing

For a nearly equiprobable distribution substantial removal is expected as K_{eff} approaches the ceiling I_{ost}. For a Porter–Thomas distribution [28, 24] rare tails cross the threshold earlier: at $N = N_*$ already $K_{\mathrm{eff}} \approx 2^{N-1} = I_{\mathrm{ost}}/2$, and removed weight is substantial. Therefore the exact RCS front below is determined directly by the weight distribution, not by equality $K_{\mathrm{eff}} = I_{\mathrm{ost}}$.

RCS limit. For random circuit sampling (RCS) [24, 25] the number of qubits N and distribution width K_{eff} are critical. Let N be the number of qubits in the circuit and $W = \sum_{i:\, p_i < \varepsilon} p_i$ the total Born weight of removed branches. The processor computational basis is the pointer-state basis set by qubit definition in interaction with the environment [2, 18]; final readout only

strengthens and registers the outcome in the same basis, it does not create it. Laboratory isolation suppresses decoherence *between* pointer states and preserves high fidelity, but does not cancel the pointer-state basis. After entangling gates joint strings $|x\rangle$ are einselection branches of the entire register with weights $p_x = |\langle x|\psi\rangle|^2$, no longer reducible to a product of independent factors (§ 6). Trimming is imposed on these weights as they grow during circuit operation, before readout. For a coherent superposition of K strings C is of order unity, not ε: BDPT on qubit mass over microseconds does not set the wall. Readout fixes the already altered distribution; if many strings are still live at readout, a single outcome comes from selection at $C \le \varepsilon$ or the degenerate edge.

The reverse question is whether residual noise pushes part of the qubits into sequential classical recording mode. An error event records which-path information about roughly one qubit and may entangle it with environment matter, in which case that dust enters $E_{\rm br}$; then $\langle k\rangle = Nde_{2q}/2$. At the working point $N \approx 400$, $d = 40$, $e_{2q} = 10^{-4}$ this is $\langle k\rangle \approx 0{,}8$—less than one qubit out of four hundred, wall shift within window $N \approx 400 \pm 3$. The window in which the test works by fidelity and the window in which leakage is negligible coincide. (Upper estimate: the coherent component of which-path recording error does not create it.)

For circuits on N qubits whose output distribution approaches Haar, weights obey the Porter–Thomas distribution: $P(p) = 2^N e^{-2^N p}$. Typical minimum weight is of order $p_{\min} \sim 2^{-2N}$. The condition $p < \varepsilon$ touches the first rare tails around $N \approx 200$, but their total weight is negligible; a typical branch becomes invisible around $N \approx 400$. The result also depends on circuit depth d—the number of sequential two-qubit gate layers: Porter–Thomas distribution for two-dimensional connectivity is established at saturation depth $d_{\rm sat} \sim \sqrt{N}$, over which the light cone covers the entire chip; at $d \gtrsim d_{\rm sat}$ typical weights are already Haar, while in smaller circuits branch removal starts later.

Integrating the Porter–Thomas distribution up to threshold $p_{\rm thr}$ gives

$$W(u) \;=\; 1 - (1+u)e^{-u} \;\simeq\; \frac{u^2}{2}, \qquad u = 2^N p_{\rm thr}. \tag{30}$$

Threshold $p_{\rm thr} = \varepsilon$ and $u = 2^{N-N_*}$. For a Haar state $u \approx 2K_{\rm eff}/I_{\rm ost}$, therefore condition $u = 1$ corresponds to $K_{\rm eff} \approx I_{\rm ost}/2$, not saturation of the formal ceiling. At sensitivity $\delta = 10^{-3}$ the effect becomes visible around $N = 396$ qubits. At $N \approx 400{,}4$ one removes $W = 1 - 2/e \approx 26{,}4\%$; at $N = 403$ already $W \approx 98\%$.

It is important to distinguish removed *mass* W and the number of surviving *strings*. The fraction of branches with $p_i \ge \varepsilon$ equals e^{-u}, whence

$$n = De^{-u} = 2^N \exp\!\left(-2^{N-N_*}\right). \tag{31}$$

The limit $W \to 1$ means only that almost all Born mass went into tails below ε, not that one string remains: at $N = 405$ still $n \sim 10^{111}$. After renormalization

$$K_{\rm eff} = \frac{D(1+u)^2 e^{-u}}{u^2 + 2u + 2}. \tag{32}$$

For the Porter–Thomas distribution the model gives two parameter-free markers: soft trimming front at $N = N_* \approx 400{,}4$ ($u = 1$) and *classicality* $N_{\rm cl}$ at $n = 1$, i.e. at

$$N_{\rm cl} \ln 2 = 2^{N_{\rm cl} - N_*}, \qquad N_{\rm cl} \approx 408{,}5 \tag{33}$$

($K_{\rm eff} \to 1$; at $N = 401, 403, 405, 408$ we have $K_{\rm eff} \approx 2^{398{,}6}, 2^{394}, 2^{370}, 2^{128}$). $N_{\rm cl}$ is classicality of the *pre-measurement state*: one pointer string remains. An ordinary apparatus flash is the other path: few fat branches and $C \le \varepsilon$ after BDPT on Mc^2 of the record. The Collapse Threshold Rule removes all invisible branches even if joint coarse-graining changes the state by order unity.

Shor algorithm limit. Threshold influence is determined not by qubit number per se but by branching width in the pointer-state basis. Partition $\{P_i\}$ is set by the environment and therefore is the same for the entire register: it is the same computational Z-basis as in RCS above. A periodic comb is a coherent superposition of pointer strings, not a pointer state, so it cannot be taken as a separate branch: that would be a representation convenient for the algorithm, not a partition set by interaction with the environment (§ 5). Coarse-graining to combs would require the environment to record the second register and not record the first; meanwhile orthogonality of the second register by itself is not irreversible classical recording.

Consider stages of the standard Shor algorithm [29]. Let m be the width of the counting register, $Q = 2^m$ and n the module to factor. After Hadamard transforms the first register remains a product of single-qubit states ("coherent product" mode, § 6), therefore 2^{-m} is a product of weights $1/2$, not the weight of one branch; branch removal does not arise. Modular exponentiation entangles the registers:

$$|\psi\rangle = \frac{1}{\sqrt{Q}} \sum_{k=0}^{Q-1} |k\rangle\, |a^k \bmod n\rangle, \tag{34}$$

All Q pointer-basis strings carry equal weight $p_i = 1/Q$. Factorizability of this state in the Z-basis is determined by order r: the quantity $a^k \bmod n$ depends only on $k \bmod r$, therefore exact decomposition into independent factors is possible if and only if $r \mid 2^m$, i.e. when $r = 2^j$. In this case, setting $k = k_{\rm hi} \cdot 2^j + k_{\rm lo}$, we obtain $|\psi\rangle = |+\rangle^{\otimes(m-j)} \otimes \frac{1}{\sqrt{r}} \sum_{k_{\rm lo}} |k_{\rm lo}\rangle |a^{k_{\rm lo}}\rangle$, and by additivity $\sum_a K_{\rm eff}(\rho_a) = r + 2(m-j)$. At $j < m$ all weights equal $1/2$ or $1/r$ and remain above ε while $r \le I_{\rm ost}$ (i.e. $j \lesssim N_*$); at $j = m$ the sole weight is 2^{-m}, and the wall arrives at $m \gtrsim N_*$ as for a flat distribution. While all weights $\ge \varepsilon$, branch removal does not arise. At $r \ne 2^j$ the state is non-factorizable and $K_{\rm eff} = Q = 2^m$ (§ 5). Hereafter we assume $r \ne 2^j$. For RSA module $n = pq$ the fraction of bases a whose order is a power of two is inverse to the odd part of the exponent of group $\mathbb{Z}_n^*$ and is negligibly small; moreover at $r = 2^j$ factorization does not require full period finding. Therefore the caveat does not weaken the conclusion about cryptographic modules but is necessary for correctness of the criterion.

The Collapse Threshold Rule is imposed during the circuit, not only on this final state. Modular exponentiation is implemented by sequential controlled-U on one control at a time; after t steps $\rho = \rho_{t+\text{target}} \otimes |+\rangle^{\otimes(m-t)}$, and by additivity $\sum_a K_{\rm eff}(\rho_a) = 2^t + 2(m-t)$. At $r \ne 2^j$ the entangled block carries 2^t equal branches of weight 2^{-t}; unprocessed controls in $|+\rangle$ are not yet touched (at $r = 2^j$ the block itself splits into $|+\rangle^{\otimes(t-j)}$ and an r-string part after $t > j$ steps, and the estimate below does not apply)—global weight 2^{-m} is not used here since the state is factorized. At $t > N_* \approx 400{,}4$ all block branches are below ε, the degenerate edge fires (§ 5), and the block collapses to one string $|k_0\rangle|a^{k_0}\rangle$; the algorithm breaks already in the middle of exponentiation. With further entanglement $K_{\rm eff}$ grows again, but the final limit remains $m \lesssim N_*$. Order r enters the criterion only through non-factorizability $r \ne 2^j$; given that condition the value of r itself does not enter the limit.

At the final step $t = m$ the distribution is flat, not Porter–Thomas, therefore the front is razor-sharp: at $m \le 400$ one has $p_i = 2^{-m} \ge \varepsilon$ and all strings are visible, while at $m \ge 401$ condition $p_i < \varepsilon$ holds immediately for all. The second case is exactly the degenerate edge of the Collapse Threshold Rules (§ 5): joint removal would zero ρ, therefore one string $|k\rangle|a^k\rangle$ is preserved with conditional Born probability. One string of periodic structure carries nothing, and subsequent QFT gives an almost uniform spectrum instead of peaks at multiples of Q/r. The value of r does not enter this condition beyond the already made assumption $r \ne 2^j$.

For Shor and QPE what matters is non-factorizability of the state in the pointer-state basis (for period finding—condition $r \ne 2^j$) and counting-register width; neither order r itself nor circuit size enter the criterion given the first condition:

$$\log_2 K_{\rm eff} = m \lesssim N_* \approx 400{,}4. \tag{35}$$

Standard Shor for an L-bit module uses $m = 2L$, therefore it is viable only at $L \lesssim 200$ and fails already at $L \approx 201$ ($m = 402$). For RSA-512 ($m = 1024$) it *collapses* in the model.

However m need not equal $2L$. Ekerå–Håstad schemes [30] reduce RSA factorization to short discrete logarithm and use a counting register of width $m \approx L/2 + L/s$ with compromise parameter s. For RSA-512 at $s \geq 4$ this gives $m \leq 384 < N_*$: string weight remains above threshold, the wall does not fire, and the module factorizes. The model therefore does not protect RSA-512—it forbids only standard period finding with $m = 2L$. Since $m \geq L/2$ for any s, all known period-finding schemes for RSA-1024 and RSA-2048 fall in zone $m > N_*$; the period-finding boundary lies around $L \gtrsim 800$.

Period finding is not the only path. The Collapse Threshold Rule removes invisible branches and renormalizes the remainder, so the wall can be used as a filter. By Coppersmith's theorem [31], knowing the high $L/4$ bits of a factor p factors $N = pq$ classically in polynomial time. For RSA-1024 this is 256 bits $< N_*$. The circuit: superposition over those bits; a reversible flag "Coppersmith found a divisor" (polynomial circuit, $K_{\text{eff}} = 2^{256}$); controlled padding of $m \geq 145$ qubits on unsolved branches. After padding the state is non-factorizable: wrong strings carry global weight $2^{-(256+m)} \leq 2^{-401} < \varepsilon$, correct ones (high bits of p or q) carry $2^{-256} \geq \varepsilon$. This is not the degenerate edge: a visible remainder exists, trimming leaves 1–2 strings, and measurement plus classical Coppersmith finishes factorization. The path works at $L/4 \lesssim N_*$, i.e. up to about $L \simeq 1600$. RSA-2048 needs 512 known bits: already the starting superposition gives every string, including the correct one, weight $2^{-512} < \varepsilon$, and the degenerate edge fires. No more economical reduction for balanced RSA is known. The protection boundary therefore shifts from $L \gtrsim 800$ to $L \gtrsim 1600$: the model does not forbid factorization of RSA-1024, but forbids RSA-2048.

General branching wall. From the above follows a general prohibition not tied to period finding. Any *non-factorizable* superposition with equal weights $p_i = 2^{-m}$ over more than 2^m pointer-basis strings at $m \geq 401$ is invisible as a whole: the degenerate edge fires (§ 5), and the register collapses to one string. Under the same condition fall Grover search on $N > N_*$ qubits (uniform superposition over 2^N strings), QPE and amplitude estimation at coherent counting-register width $m > N_*$; only factorization saves (§ 6)—$|+\rangle^{\otimes N}$, intermediate Shor stages, narrow Ekerå–Håstad registers. For a *flat* distribution the degenerate edge arrives immediately at $m \geq 401$. In RCS the Porter–Thomas distribution is different: at $N = 403$ already $W \approx 98\%$, at $N = 405$—$W \to 1$, but still $n \sim 10^{111}$ strings survive; one string remains only around $N_{\text{cl}} \approx 408{,}5$. The soft front of removed mass is steep in window $N \approx 400$–405, classicality $N_{\text{cl}} \approx 408{,}5$.

This is the main falsifiable prediction of the model—stronger and closer to experiment than the consequence for RSA-2048: mirror circuit, local marginal test, and RCS scan on one 404-qubit processor (§ 9) directly test the geometry of this wall, whereas factorization of modules above $L \gtrsim 1600$ is not yet reachable by witness width.

Experiment does not contradict the model even where analog simulators involve hundreds and thousands of entangled particles: particle number is not branch number. For a GHZ state of 1000 spins $K_{\text{eff}} = 2$, not 2^{1000}; spin chains and cold atoms evolve by Hamiltonian, not maintaining equal superposition over 2^N strings of computational Z-basis. The Collapse Threshold Rule counts branches in the environment pointer basis (§ 5), not arbitrary many-body entanglement; direct conflict would arise only upon demonstration of stable uniform superposition wider than N_* in this basis—no such experiment exists.

In a fault-tolerant scheme physical qubits may be far more than N_*, but the limit concerns not their number but logical algorithm structure—width m of the logical counting register. Syndrome measurements record only errors in the environment and do not reveal logical state (stabilizers commute with logical operators), therefore period-finding strings by themselves are not converted to classical recorded mode. For the same reason estimate $\langle k \rangle$ does not apply to the logical register: correction does not let the environment accumulate which-path recording of

logical state specifically.

For modules above $L \gtrsim 1600$ one cannot assert that a more economical reduction cannot exist, packing a verifiable witness into N_* qubits and using trimming as a filter; no such reduction is known for balanced RSA. Nor is an algorithm known that factors RSA-2048 while never exceeding $K_{\text{eff}} \leq I_{\text{ost}}$: the Coppersmith filter does not bypass the wall, but uses it as a resource.

9 Experimental Test

The absolute value $\varepsilon \sim 10^{-121}$ is unreachable in the laboratory as a direct measurement. Therefore we propose to capture boundary geometry, the break between two chains, and the nonlinear signal of the local marginal test—a shift of one qubit's probabilities without measuring the entire register. The scan runs over qubit number N, circuit depth d, and chain structure. These are not three independent threshold parameters: the Collapse Threshold Rule compares branch weights with ε; N, d, and structure only set what distribution $\{p_i\}$ the circuit prepares. In RCS after saturation $d \gtrsim d_{\text{sat}}$ the wall sits around $N \approx 400$; a smaller circuit shifts it in d, and narrow branching at the same N and d gives no wall.

Boundary geometry. Control under independent errors: hyperbola $N \cdot d \approx \text{const}$ (error budget) is an accumulation-model approximation, not a consequence of standard quantum mechanics and not the Collapse Threshold Rule threshold geometry; the working control curve is built from measured noise channels. Threshold wall model (§ 8): at $d < d_{\text{sat}}$ the boundary depends on depth, and after Porter–Thomas distribution is established at $d \gtrsim d_{\text{sat}}$ it bends toward vertical. The bend lies in region $N \approx 400 \pm 3$. Front $W = 1 - (1 + u)e^{-u}$ with $u = 2^{N-N_*}$; per qubit u doubles. At $u \ll 1$ we have $W \simeq u^2/2$, therefore W grows roughly fourfold per qubit; in the transition window ($N \sim 400$) growth of W slows. The threshold is not tuned, therefore bend position should not follow reduction of ordinary error rate.

Mirror circuit. The idea is random-circuit echo: a register from $|0\rangle^{\otimes N}$ is run through scrambler U and reverse mirror $U^\dagger$; under standard unitary evolution the state returns to zero, while the Collapse Threshold Rule inside U removes branches and breaks compensation. Metric is return probability $F = \left|\langle 0|U^\dagger U_{\text{phys}}|0\rangle\right|^2$ [1]. Predicting F classically is not required: it suffices to run $UU^\dagger$ repeatedly on the chip and measure the fraction of returns to $|0\rangle^{\otimes N}$. Poisson: $M \sim 100/F$ copies. Under independent stochastic two-qubit gate noise a rough estimate is $F \approx e^{-Nde_{2q}}$ (in each of two passes of order $Nd/2$ gates). At $e_{2q} = 3 \times 10^{-3}$, $d = 40$, $N \sim 400$ this is $F \sim 10^{-21}$, which practically closes the test; at $e_{2q} = 10^{-4}$—of order 0,20. Testability condition $e_{2q} \lesssim 5/(Nd)$ (so that $F \gtrsim 10^{-2}$). Randomized compiling [32] separates stochastic noise from coherent compensation. Sensitivity δ of the mirror circuit sets the lower edge of the window: at $\delta = 10^{-3}$ removed weight becomes visible around $N = 396$ qubits.

Discrimination from hardware noise. At fixed N, d, and circuit structure the ordinary noise boundary should shift when measured e_{2q} changes, whereas threshold front position in N is set by ε and does not shift; lowering e_{2q} only changes the ability to see this front. Therefore repeating the scan on devices or calibrations with different e_{2q} is the main distinguishing test. Unknown correlated noise not following measured e_{2q} remains a competing explanation [33, 34].

Paired circuits. Comparison on one chip at the same N and same depth d; arms differ by chain structure, not depth. Branching width is set by how many pointer-basis strings (computational Z) carry weight *at each layer*, not by whether all qubits are entangled. Arm A, wide: random single-qubit rotations and two-dimensional layers of two-qubit gates (RCS). Superposition fills almost all 2^N strings, and for Haar distribution $K_{\text{eff}} \approx 2^{N-1}$ after $d \gtrsim d_{\text{sat}}$. Arm B,

narrow: K_{eff} is small *throughout* the circuit. A classical reversible network (CNOT, SWAP on computational states) holds one string, $K_{\text{eff}} = 1$. A CNOT chain after one H gives GHZ: all qubits are entangled, but only two live branches, $K_{\text{eff}} = 2$. Echo $U^\dagger U$ on the narrow path is *not* such: in the middle of the first U the distribution is already Haar, and the threshold fires before unwind. "Narrow path" means small branching width, not smaller d and not disabling part of the qubits. In the corresponding transition region arm B works normally; A/B difference captures wall geometry. Signature is residual deviation of A not disappearing when ordinary errors are reduced. Isolation does not close the gap: chain structure helps, not only increasing relaxation and dephasing times T_1/T_2 (ordinary hardware qubit coherence scales). Scan along hyperbola $N \cdot d \approx \text{const}$ is a separate control of the ordinary error-accumulation model (noise budget almost the same at different N), not a difference of depths A and B; at each such (N, d) arm systematics differ.

Local marginal test. Circuit: 404 qubits on one chip—control C, 402 ancilla, qubit A entangled with C by a Bell pair. Qubit C is not measured. The bit is encoded by operation: Z-control—controlled-U; X-control—H on C, controlled-U, H back. Conditional circuit on ancillas: at $C = 0$—identity, at $C = 1$—scrambler U; mirror $U^\dagger$ is not needed—marginal is fixed at trimming moment. Metric is marginal of qubit A: $\rho_A = \text{Tr}_{C,\text{anc}}(\rho)$, i.e. $P(A{=}0)$ and $P(A{=}1)$.

In standard quantum mechanics both operations are local relative to A, and its marginal is strictly 1/2. In the model Z-control cuts fraction $W(2u)$ of the scrambled arm, and after renormalization

$$P_Z(A{=}1) = \frac{1 - W(2u)}{2 - W(2u)}, \qquad P_X(A{=}1) = \tfrac{1}{2}, \tag{36}$$

where $u = 2^{N-N_*}$. Under X-control scrambled micro-branches are symmetrically linked to $A = 0$ and $A = 1$; removal shifts both outcomes equally. Scan over N: Signature is monotonic fall of

N	$P_Z(A{=}1)$	$P_X(A{=}1)$
400	35,6%	50%
401	16,3%	50%
402	**1,6%**	**50%**

Table 1: Marginal probabilities of qubit A under Z- and X-control; N is the number of ancillas.

$P_Z(A{=}1)$ with fixed $P_X(A{=}1) = 1/2$. Table values are ensemble frequencies after trimming; in a single shot after selection the outcome of A is 0 or 1. Hardware fault reproduces suppression of both arms, not one. The test does not depend on choice of preferred layer (§ 11): at centimeter baseline shift $vl/c^2 \sim 4 \times 10^{-14}$ s against readout time ~ 100 ns, i.e. the layer is unobservable and the result does not depend on it.

Failure of standard Shor. Signature is QPE/Shor on a counting register of growing width m: collapse at $m \to 400$ for any order r that is not a power of two (§ 8). Base a with $r = 2^j$ gives an extra zero control: the counting register factorizes, and at $j \lesssim N_*$ all weights remain above ε—the wall should not fire until the same m as for factorized $|+\rangle^{\otimes N}$. A pair of runs at the same m and d with $r = 2^j$ and $r \neq 2^j$ is the same differential test as paired A/B circuits above, but inside one algorithm. Standard Shor with $m = 2L$ in the model does not factorize modules from $L \approx 201$ bits, whereas RSA-512 remains vulnerable to Ekerå–Håstad schemes with $m \leq 384$. The RCS transition is not shifted by hardware improvement: the front remains around 400–404 qubits. A 404-qubit processor enables the local marginal test and superluminal channel. Competing explanation is correlated noise [33]: its boundary follows error rate, whereas the threshold boundary is set by ε. Statistical test of RCS and its dependence on noise model are discussed in [34]. Comparison with GRW/CSL [35, 36, 37]: they have plane (λ, r_c), and mass

enters the spontaneous collapse rate; here the actualization threshold $\varepsilon = 1/I_{\text{ost}}$ is not tuned, and mass enters only through BDPT. Structural choices—einselection basis, renormalization (§ 5), and width measure—remain.

10 Superluminal Telephone

After confirmation of the local marginal test (§ 9) the same circuit is moved to a separated baseline of distance l: qubit A, entangled with C by a Bell pair, is delivered to Bob in advance by telecom fiber, and the effect becomes a communication channel. Working point is 404 qubits (402 ancillas; C and A are not included in N). Alice does not measure C; the bit is encoded by Z- or X-control over the entire ensemble of M acts. At $C = 0$—identity, at $C = 1$—scrambler U. Bob measures A in the Z basis.

In ordinary quantum mechanics Alice's local unitary operations on C and ancillas do not change ρ_A at Bob: when measuring A in the Z basis he always sees $P(A{=}1) = 50\%$, regardless of Alice's choice between Z- and X-control. In the model trimming $\mathcal{N}$ changes global branches of system $C+\text{ancilla}+A$ and renormalizes the remainder (§ 5). That is the superluminal shift of the marginal. Z- and X-control yield *different* full matrices ρ, not two ensembles of the same ρ. On Alice's isolated chip C does not fall to ε in microseconds: qubit mass is too small for BDPT, and there are no copy-records. Selection at Bob actualizes the already shifted bit. At $N = 402$:

$$P_Z(A{=}1) = \frac{1 - W(2u)}{2 - W(2u)} \approx 1{,}62\%, \qquad P_X(A{=}1) = \tfrac{1}{2}, \tag{37}$$

difference $\approx 48{,}4$ percentage points. In ordinary quantum mechanics both probabilities equal 50%.

The signal arises precisely from the *absolute* threshold. If the threshold depended only on relative weights, renormalization would not change Bob's observable probabilities. In GRW-type collapse models averaged evolution is linear, therefore different preparations of one ρ are indistinguishable and the superluminal channel is closed [35, 36, 37]. For Weinberg's nonlinear quantum mechanics Gisin and Polchinski showed that local operations on one side of an EPR pair change statistics on the other [38, 39, 40]. Here U_Z and U_X are both local relative to A, therefore in linear theory ρ_A does not change under either; nonlinearity of $\mathcal{N}(\rho)$ violates precisely this marginal preservation (§ 5).

Bob's unit frequency is the mean of M coin tosses with probability q; width equals $\sqrt{q(1-q)/M}$. For $\pm 2\sigma$ intervals around $P_Z(A{=}1)$ and $1/2$ not to overlap,

$$\left|P_Z(A{=}1) - \tfrac{1}{2}\right| \geq 2\Big(\sqrt{P_Z(A{=}1)\big(1 - P_Z(A{=}1)\big)/M} + \sqrt{1/(4M)}\Big). \tag{38}$$

Hence at $N = 402$ we obtain $M \approx 7$ acts per bit. Each act is direct controlled-U on ancillas at $C = 1$ (at $C = 0$—identity). On a two-dimensional lattice saturation is $d_{\text{sat}} \sim \sqrt{N} \approx 20$ layers (§ 8) at layer duration of tens of nanoseconds giving $t_{\text{run}} \approx 0{,}5\ \mu$s per act; $M \cdot t_{\text{run}} \approx 3{,}5\ \mu$s, bitrate ≈ 280 kbit/s. Each act consumes one Bell pair C–A (after readout of A new entanglement is needed), therefore a stable channel requires pair distribution rate $R_{\text{Bell}} = 1/t_{\text{run}} \approx 2 \times 10^6$ pairs/s; at lower R_{Bell} real bitrate is limited by R_{Bell}/M, not only by t_{run} on the chip. Readout of qubit A at Bob takes nanoseconds and is not included in t_{run}.

The superluminal signal itself is a trimming jump of duration t_{coll} (§ 7), simultaneous on the FRW layer (§ 11), not continuous $\chi(t)$. The channel beats light if $M \cdot (t_{\text{run}} + t_{\text{coll}}) < l/c$; at room-temperature estimate $t_{\text{coll}} \sim 25$ fs minimum baseline $\approx 1{,}05$ km. The local marginal test remains on one 404-qubit chip (baseline in centimeters); the race with light is set by separating A and Alice's station by at least that distance after effect confirmation. Synchronization of control and readout on a unified comoving time scale is described in § 11.

Two-way communication. Two-way communication is possible with symmetric 404-qubit setups at each station: each has 402 ancillas and its own half of a Bell pair C–A. In Alice→Bob tick C and ancillas are at Alice, A at Bob; in Bob→Alice tick roles swap. C and A are control and marginal readout, not "Alice's and Bob's qubits." One act consumes a pair: after readout of A new entanglement is needed, the same pair is unavailable for the next bit. In the active tick the pair-source station must provide the same $R_{\mathrm{Bell}} \approx 2 \times 10^6$ pairs/s. With one pair one cannot simultaneously transmit and receive—either we speak or listen. Opposing ticks are separated on the comoving layer (§ 11), not launched simultaneously. Other engineering assumptions are in § 12.

11 CMB Frame

The model *assumes* that the quantity I_{ost} and simultaneity of trimming and selection jumps are set not by laboratory clocks but on a layer of constant cosmological time of homogeneous isotropic Friedmann–Robertson–Walker (FRW) metric—standard geometry of the expanding Universe. In this idealization layer $t_{\mathrm{FRW}} = \mathrm{const}$ is one for the entire metric: all comoving observers (zero peculiar velocity) agree on simultaneity. Practically the same preferred frame is the rest frame of the cosmic microwave background (CMB); it is to this frame that the model attaches the jump, not to local simultaneity of each laboratory and not to continuous $\chi(t)$ of the BDPT hypothesis. An Earth setup moves relative to the CMB at speed of order 370 km/s, therefore its Einstein or GPS synchronization is shifted from this layer by vl/c^2 (below). Entangled pairs are separated subluminally and remain inside one causal patch, therefore the operation budget is defined over the entire experiment. The jump is simultaneous on this layer in the idealization of homogeneous FRW metric; in inhomogeneous or rotating geometries a global comoving layer is not naturally defined. Rest relative to the CMB defines a preferred frame; compatibility with constraints on Lorentz-invariance violation [41, 42] is not discussed here. Open questions of extending the CMB layer to local systems, preferred frame, and causal consistency are in § 12.

Preferred-layer test. The laboratory moves relative to the CMB at speed $v \approx 370$ km/s. Shift vl/c^2 exceeds one readout time τ at $l \gtrsim (c^2/v)\tau$; for $\tau \sim 100$ ns this is $l \gtrsim 24$ km. Events are launched *deliberately simultaneously* in the laboratory. Then orientation of the baseline relative to the CMB dipole decides whose recording first crosses the FRW layer, and Bob's statistics should oscillate with *sidereal* period ($\approx$ 23 h 56 min), not solar days. Solar motion (temperature, grid) is filtered out by this period.

The Alice→Bob channel is not switched off: if Alice launches measurement earlier than Bob in the laboratory with margin greater than vl/c^2 (at 24 km this is ~ 100 ns), she is earlier on the FRW layer too at any time of day. This margin already covers $t_{\mathrm{coll}} \sim 25$ fs.

This is a separate protocol, not a race with light. The goal is to verify that the preferred layer coincides with rest relative to the CMB. Laboratory simultaneity is set by Einstein synchronization: light pulse round trip, midpoint of flight is the common tick; in practice GPS/GNSS or two-way fiber transmission do the same. Such synchronization is tied to Earth time, not CMB: layer difference is precisely shift vl/c^2.

Synchronization by comoving clocks. The model postulates collapse simultaneity on the CMB layer. Since laboratory clocks do not coincide with CMB-layer simultaneity, to synchronize by absolute collapse time local clock readings must be converted to cosmological time t_{FRW}. Alice and Bob first synchronize atomic clocks in the usual way, then convert proper times along known worldlines of the stations and metric model. For nearby Earth stations the main simultaneity correction to first order equals $\Delta t_{\mathrm{CMB}} \simeq \mathbf{v}\cdot\Delta\mathbf{x}/c^2$, accounting for velocity and baseline orientation relative to the CMB. If, for example, Alice is on Earth and Bob in open space, this correction

alone is insufficient: in a weak field local clock rate is linked to comoving time by

$$\frac{d\tau}{dt_{\mathrm{FRW}}} \simeq 1 + \frac{\Phi}{c^2} - \frac{v^2}{2c^2}, \tag{39}$$

therefore along each worldline one integrates gravitational potential Φ and kinematic time dilation separately. Additionally one accounts for Earth rotation and orbital motion, space-station trajectory, propagation delay of calibration signals, and local metric perturbations.

On this common scale one can schedule control and readout moments of the superluminal channel in advance. Such synchronization is especially convenient at large, including space, distances: both stations use one cosmological time coordinate rather than determining order of remote events by exchanging light signals during the session. For baseline ~ 1 km maximum kinematic correction is about 4 ns, small compared with act $\sim 0{,}5\ \mu$s but available for calibration; at space baseline combined kinematic and gravitational correction becomes substantial. This is not a separate physical signal of global clocks but conversion of local clock readings to the assumed preferred frame; its accuracy is limited by ephemeris precision, gravitational field model, and extension of the CMB layer to inhomogeneous geometries (§ 12).

12 Discussion and Open Questions

Integer micromodel. The postulate of limited description and sketch of integer grid n_i/I_{ost} are given in § 5. Full microtheory from this sketch is not yet derived. Open obstacles: under tensor product of subsystems weights of form $n_i m_j/I_{\mathrm{ost}}^2$ do not lie on the same grid—a composition theory with preserving rounding is needed; as $I_{\mathrm{ost}}(t)$ decreases (§ 3) the entire discrete distinguishability scale is rebuilt; the moment of applying map $\mathcal{N}$ is not specified—after a gate, after einselection recording by the environment, or continuously during evolution—which determines whether a new branch can grow to $p_i \geq \varepsilon$. Until these are resolved the Collapse Threshold Rule specifies an effective one-step rule on the full density matrix of an entangled system implementing the postulate for einselection objects, not a closed discrete quantum mechanics with unambiguous evolution $\rho(t)$.

Local Bekenstein bound. In this work the threshold is determined by the global causal-patch budget and is the same for all entangled systems. An alternative microtheory is possible in which available precision is additionally limited by the Bekenstein bound $S \leq 2\pi ER/(\hbar c)$ for a region containing physical carriers of entanglement [3]. For a compact system this local memory limit may be stricter than the cosmological operation budget; then ε and critical qubit number N_* will depend on setup size and energy. Whether quantum state should be treated as locally stored information or information of the entire causal patch remains an open question.

Palmer invariant set. A related idea is Palmer's invariant set [43]: reality is limited to a fractal invariant set. Here the limitation is different: operational distinguishability of states within a causal patch, not geometry of an attractor.

Branches. Absolute threshold requires unambiguously defined branches. Difficulties of their definition and counting are discussed by Saunders [44]. Here probabilities are set by absolute Born weights $p_i = \mathrm{Tr}(P_i\rho)$ in the einselection basis, not by branch number; the rule is a function of a single ρ. Contrast between Z- and X-control in the local marginal test and in its separated realization does not arise from two decompositions of one ρ into ensembles but from different unitary operations changing ρ itself; dependence on coarse-graining level remains an open question. Practically the weight threshold acts on pointer strings of the full matrix, while C and BDPT act on already trimmed sectors; whether these partitions always coincide is not shown.

BDPT and the budget I_{ost}. Proper-time dephasing and selection itself at $C \leq \varepsilon$ are not derived from I_{ost}: the threshold sets visibility only. Which dust grains already enter Mc^2 of a branch is a separate input. If all matter with which the system is even slightly entangled is written into E_{br}, the cluster spreads to the horizon. Weak coupling of two experiments: infinitesimal entanglement formally makes them one ρ; whether to count them as two clusters is open. The narrow P1 of [23] via $\mathcal{D}_s$ is stronger than "all matter of the branch"; for an apparatus with which-path recording both rules give Mc^2 of the dust.

CMB frame and causality. In § 11 simultaneity of trimming and selection jumps is postulated on a constant FRW time layer; on cosmological background it coincides with the CMB rest frame. Continuous $\chi(t)$ is not tied to this layer. For inhomogeneous gravitational field no unambiguous extension of this layer from cosmological background to local Earth and spacecraft systems is defined. Conversion of station proper times to t_{FRW} depends on worldlines, ephemerides, and metric model; required precision and experimental way to test preferred frame remain open. The preferred layer sets a single order of remote events and thereby may exclude causal loops arising under Lorentz-invariant superluminal transmission only if all nonlinear dynamics strictly obey this foliation; causal consistency of the full microtheory is not yet shown.

Non-conservation of energy. Estimate (27) links non-conservation scale with t_{coll} via act uncertainty ($\Delta E_{\text{unc}} \gtrsim \hbar/t_{\text{coll}}$, non-conservation $\lesssim \Delta E_{\text{unc}}$); collapse (29) gives exact renormalization shift and does not contain t_{coll}. Under partial trimming it may be noticeable, but in degenerate Born choice mean shift vanishes (§ 7), unlike GRW/CSL models [35, 36, 37] where parameter λ is limited by spontaneous heating. Whether energy is emitted or absorbed, whether a perpetual motion machine is possible—beyond this work. Energy balance at collapse is a topic for a separate article.

Objectivation only. The postulate also admits a narrower rule: invisible branches are neither removed nor renormalized, and at $C \leq \varepsilon$ only objectivation $\mathcal{C}^{\text{sel}}$ acts. The superluminal channel (§ 10) is then closed: the marginal shift is produced by map $\mathcal{N}$, not by selecting one live branch. The working model takes both maps; which of them enters the dynamics remains open.

Superluminal telephone engineering. The scheme in § 10 sets preliminary architecture and working estimates for direct controlled-U. The engineering bottleneck is not t_{run} on the chip nor t_{coll}, but the Bell-pair pipeline: at $M \approx 7$ and $t_{\text{run}} \approx 0{,}5$ μs one needs rate $R_{\text{Bell}} \approx 2 \times 10^6$ pairs/s per direction, orders of magnitude above typical quantum communication lines. Possible optimization is to prepare $U|0\rangle$ offline on a buffer register parallel to the previous act, and in the tick at $C = 1$ substitute it by controlled-SWAP (at $C = 0$ ancillas remain in $|0\rangle$). In standard quantum mechanics this is equivalent to controlled-U at the level of the final global ρ; for nonlinear Collapse Threshold Rules equivalence requires a separate assumption of no threshold event during preparation and storage of the prefabricated state. If such optimization is admissible, act duration shrinks to $t_{\text{run}} \approx 50$ ns (loading prefabricated state and Z/X-control on C, without rerunning d_{sat} layers), bitrate rises to $\approx 2{,}8$ Mbit/s, minimum baseline to ≈ 100 m, but R_{Bell} rises to $\approx 2 \times 10^7$ pairs/s. In the nonlinear model final state by itself may not determine outcome: threshold may fire already during offline preparation, therefore equivalence of prefabricated state and controlled-U must be specified by microdynamics and tested experimentally. Also open are resources and duration of multi-qubit controlled-SWAP, buffer reset, pair generation and delivery at rate R_{Bell}, Bell-pair preservation, and readout synchronization.

References

[1] B. Swingle and N. Yunger Halpern. Resilience of scrambling measurements. *Physical Review A*, 97:062113, 2018.

[2] W. H. Zurek. Decoherence, einselection, and the quantum origins of the classical. *Reviews of Modern Physics*, 75:715–775, 2003.

[3] R. Bousso. The holographic principle. *Reviews of Modern Physics*, 74:825–874, 2002.

[4] S. Lloyd. Ultimate physical limits to computation. *Nature*, 406:1047–1054, 2000.

[5] S. Lloyd. Computational capacity of the universe. *Physical Review Letters*, 88:237901, 2002.

[6] T. Banks and W. Fischler. An holographic cosmology. arXiv:hep-th/0111142, 2001.

[7] L. M. Krauss and G. D. Starkman. Universal limits on computation. arXiv:astro-ph/0404510, 2004.

[8] R. V. Buniy, S. D. H. Hsu, and A. Zee. Is Hilbert space discrete? *Physics Letters B*, 630:68–72, 2005.

[9] R. V. Buniy, S. D. H. Hsu, and A. Zee. Discreteness and the origin of probability in quantum mechanics. *Physics Letters B*, 640:219–223, 2006.

[10] Y. I. Ozhigov. Algorithmic approach to quantum physics. arXiv:quant-ph/0412196, 2004.

[11] R. Srikanth. A computational model for quantum measurement. arXiv:quant-ph/0302160, 2003.

[12] R. Srikanth. The quantum measurement problem and physical reality: A computation theoretic perspective. arXiv:quant-ph/0602114, 2006.

[13] P. C. W. Davies. The implications of a cosmological information bound for complexity, quantum information and the nature of physical law. *Fluctuation and Noise Letters*, 7:C37–C50, 2007. arXiv:quant-ph/0703041.

[14] F. Del Santo and N. Gisin. Physics without determinism: Alternative interpretations of classical physics. *Physical Review A*, 100:062107, 2019.

[15] R. Gambini, L. P. García-Pintos, and J. Pullin. Undecidability as solution to the problem of measurement: Fundamental criterion for the production of events. *International Journal of Modern Physics D*, 20:909–918, 2011. arXiv:1009.3817.

[16] D. Harlow and P. Hayden. Quantum computation vs. firewalls. *Journal of High Energy Physics*, 2013(6):085, 2013.

[17] G. W. Gibbons and S. W. Hawking. Cosmological event horizons, thermodynamics, and particle creation. *Physical Review D*, 15:2738–2751, 1977.

[18] W. H. Zurek. Quantum Darwinism. *Nature Physics*, 5:181–188, 2009.

[19] J. A. Wheeler. Information, physics, quantum: The search for links. In W. H. Zurek, editor, *Complexity, Entropy, and the Physics of Information*, pages 3–28. Addison-Wesley, Redwood City, 1990.

[20] J. Watrous. *The Theory of Quantum Information*. Cambridge University Press, Cambridge, 2018.

[21] G. Lüders. Über die zustandsänderung durch den meßprozeß. *Annalen der Physik*, 8:322–328, 1951.

[22] R. A. Horn and C. R. Johnson. *Matrix Analysis*. Cambridge University Press, Cambridge, 2 edition, 2013.

[23] O. Ponfilenok. Branch-dependent proper time. viXra:2607.0087, 2026.

[24] S. Boixo, S. V. Isakov, V. N. Smelyanskiy, R. Babbush, N. Ding, Z. Jiang, M. J. Bremner, J. M. Martinis, and H. Neven. Characterizing quantum supremacy in near-term devices. *Nature Physics*, 14:595–600, 2018.

[25] F. Arute, K. Arya, R. Babbush, D. Bacon, J. C. Bardin, R. Barends, R. Biswas, S. Boixo, F. G. S. L. Brandao, D. A. Buell, et al. Quantum supremacy using a programmable superconducting processor. *Nature*, 574:505–510, 2019.

[26] D. J. Thouless. Electrons in disordered systems and the theory of localization. *Physics Reports*, 13:93–142, 1974.

[27] R. Grobe, K. Rzążewski, and J. H. Eberly. Measure of electron-electron correlation in atomic physics. *Journal of Physics B*, 27:L503–L508, 1994.

[28] C. E. Porter and R. G. Thomas. Fluctuations of nuclear reaction widths. *Physical Review*, 104:483–491, 1956.

[29] P. W. Shor. Polynomial-time algorithms for prime factorization and discrete logarithms on a quantum computer. *SIAM Journal on Computing*, 26:1484–1509, 1997.

[30] M. Ekerå and J. Håstad. Quantum algorithms for computing short discrete logarithms and factoring RSA integers. In *Post-Quantum Cryptography (PQCrypto 2017)*, volume 10346 of *Lecture Notes in Computer Science*, pages 347–363. Springer, 2017.

[31] D. Coppersmith. Small solutions to polynomial equations, and low exponent RSA vulnerabilities. *Journal of Cryptology*, 10:233–260, 1997.

[32] J. J. Wallman and J. Emerson. Noise tailoring for scalable quantum computation via randomized compiling. *Physical Review A*, 94:052325, 2016.

[33] G. Kalai. The quantum computer puzzle. *Notices of the American Mathematical Society*, 63:508–516, 2016.

[34] Y. Rinott, T. Shoham, and G. Kalai. Statistical aspects of the quantum supremacy demonstration. *Statistical Science*, 37(3):322–347, 2022.

[35] G. C. Ghirardi, A. Rimini, and T. Weber. Unified dynamics for microscopic and macroscopic systems. *Physical Review D*, 34:470–491, 1986.

[36] G. C. Ghirardi, P. Pearle, and A. Rimini. Markov processes in Hilbert space and continuous spontaneous localization of systems of identical particles. *Physical Review A*, 42:78–89, 1990.

[37] A. Bassi, K. Lochan, S. Satin, T. P. Singh, and H. Ulbricht. Models of wave-function collapse, underlying theories, and experimental tests. *Reviews of Modern Physics*, 85:471–527, 2013.

[38] S. Weinberg. Testing quantum mechanics. *Annals of Physics*, 194:336–386, 1989.

[39] N. Gisin. Weinberg's non-linear quantum mechanics and supraluminal communications. *Physics Letters A*, 143:1–2, 1990.

[40] J. Polchinski. Weinberg's nonlinear quantum mechanics and the Einstein-Podolsky-Rosen paradox. *Physical Review Letters*, 66:397–400, 1991.

[41] D. Mattingly. Modern tests of lorentz invariance. *Living Reviews in Relativity*, 8:5, 2005.

[42] S. Liberati. Tests of lorentz invariance: an update. *Classical and Quantum Gravity*, 30:133001, 2013.

[43] T. N. Palmer. The Invariant Set Postulate: a new geometric framework for the foundations of quantum theory and the role played by gravity. *Proceedings of the Royal Society A*, 465:3165–3185, 2009.

[44] S. Saunders. Branch-counting in the Everett interpretation of quantum mechanics. *Proceedings of the Royal Society A*, 477:20210600, 2021.