# Heat from Bit: The Thermodynamic Price of Objective Collapse

Oleg Ponfilenok

Independent researcher; ponfil@gmail.com

**Abstract**

We propose the hypothesis that any mechanism of *objective* collapse must pay a thermodynamic price of $k \ln 2$ per destroyed bit of coherence *in the pointer basis of the act*—as heat $\Delta Q = kT \ln 2$ in an environment at temperature $T$. Test: calorimetry of heating upon absorption of a heralded single-photon qubit—$|+\rangle$ versus the control $I/2$.

**Keywords:** objective collapse; pointer state; quantum Darwinism; Landauer's principle; heat from bit; entropy collection

## 1 Introduction

The idea of *objective* collapse—making state reduction a physical process rather than an epistemic postulate of measurement—forms a distinct research program alongside standard quantum mechanics, where the apparent collapse is explained by unitary decoherence and correlations with the environment. The early line of Karolyházy [1] linked the demise of superpositions to spacetime uncertainty; GRW and CSL [2, 3] introduced spontaneous localization; Diosi–Penrose models [4, 5]—gravitational instability of macroscopic superpositions; in other models: crossing a scrambling threshold [6], branch-dependent dephasing of proper time [7]. Review and status—Bassi et al. [8]: the direction is actively developed and experimentally constrained, but remains marginal relative to orthodox quantum theory; decisive confirmation or final refutation is still lacking. The present work does not choose a criterion for "when," but asks about the thermodynamic price of any such act.

Objective collapse introduces a fundamentally irreversible element of dynamics, distinguishing the preferred direction of time from reversible unitary evolution. However, such irreversibility by itself is not equivalent to violation of continuous time-translation symmetry, with which Noether's theorem associates energy conservation. Therefore the energetic footprint of collapse depends on the specific non-unitary dynamics; the present work proposes a hypothesis about its thermodynamic nature (§ 5).

Landauer's principle [9] links the erasure of *classical* information to minimum dissipation $Q \geq kT \ln 2$. The price sits on logically irreversible compression of phase volume (many→one)—a consequence of the second law [10, 11]. Under decoherence the environment selects a *pointer basis*—a set of robust outcomes $|i\rangle$, proliferated in the environment (einselection; quantum Darwinism [12]). The pointer basis defines the set of *classically admissible* (incoherent) states—diagonal mixtures over $|i\rangle$. The *relative entropy of coherence* [13] $C_r(\rho) = \min_{\sigma \in \mathcal{I}} S(\rho \| \sigma)$ measures how far $\rho$ falls short of such a mixture; for the pointer basis the minimum is attained at $\rho_{\text{diag}}$ and $C_r(\rho) = S(\rho_{\text{diag}}) - S(\rho)$. For pure $|+\rangle_{\text{ptr}}$ $S(\rho) = 0$, but $C_r = k \ln 2$—full which-path coherence; for $I/2_{\text{ptr}}$ $S(\rho) = k \ln 2$, but $C_r = 0$—the same pointer statistics without phase, nothing to erase. Under *unitary* decoherence $|+\rangle \to I/2$ $S(\rho_S)$ of the subsystem grows to $k \ln 2$, but the entropy of the whole is zero: the phase goes into correlations and is recoverable in

principle (Loschmidt echo [14, 15]). Under *objective* collapse the pointer outcome is fixed irreversibly, $H = S_{\rm coh} = C_r$ in the pointer basis—a thermodynamically active quantity; according to the hypothesis of the present work its jump materializes as heat. Hormoz [16] shows the necessity of an entropic price for collapse in a ratchet model; Heat from Bit quantifies the price for objective fixation of a pointer outcome without an explicit demon.

Wheeler formulated "It from Bit" [17]: physical reality ("it") arises from acts of distinguishing pointer outcomes. *Heat from Bit*—the same parallel in thermodynamics: $C_r(\rho)$ in the pointer basis is the "bit"; its irreversible jump is the "it," materializing as heat.

We propose an experimental test (§ 6): calorimetry of a *heralded* single-photon qubit in a Mach–Zehnder (MZ) interferometer (heralded MZ)—$|+\rangle_{\rm ptr}$ yields $kT\ln 2$ per act, the classical mixture $I/2_{\rm ptr}$—none.

## 2 Intuition

The idea of Heat from Bit grew out of a simple thought experiment in which *one* photon is absorbed by an anisotropic absorber in the polarization basis that defines the pointer basis of the act. The basis is $\{|0\rangle, |1\rangle\}$ of linear polarization ($|0\rangle$—horizontal, $|1\rangle$—vertical): which of these outcomes will be irreversibly fixed upon absorption—that is, upon detection?

**Case A (definite polarization).** A single-photon state $|0\rangle$—an eigenstate of the pointer basis, aligned with the polarizer axis. Coherence $S_{\rm coh} = 0$. The photon is absorbed, the absorber receives energy $E_\gamma$—ordinary first-law heating; the alternative pointer outcome $|1\rangle$ is absent in the pre-act state, nothing to erase. Nothing changes upon an act of objective collapse.

**Case B (superposition of pointer outcomes).** The same single-photon protocol, but the photon is prepared in

$$|+\rangle_{\rm ptr} = \frac{|0\rangle + |1\rangle}{\sqrt{2}} \tag{1}$$

in the polarizer's pointer basis: $S_{\rm coh} = \ln 2$. Absorption transfers the same $E_\gamma$; the difference lies in the pre-act state, not in the photon energy.

*Step 1 (absorption).* Regardless of preparation, the act of absorption transfers $E_\gamma$ to the absorber; there is as yet no objective collapse at this stage.

*Step 2 (decoherence).* Under *unitary* evolution, coherence in the pointer basis $S_{\rm coh}$ goes into the internal degrees of freedom of the absorber and environment—into entanglement, recoverable in principle (echo, reversible cycle). The environment already interacts with the photon and carries information about the pointer outcome (here—polarization) into the *internal memory* of the surroundings. Entanglement of the absorber and environment with $|0\rangle$ and $|1\rangle$ *is* that memory—the alternative is not destroyed but encoded in correlations; under unitary evolution it is preserved, there is no erasure, global entropy does not grow.

*Step 3 (objective collapse).* Objective collapse can be understood as *clearing* internal memory: degrees of freedom that carried coherence and entanglement disappear irreversibly—a register-erasure procedure, not a reversible transfer of phase. The pointer outcome is fixed; there is no longer an "internal" quantum state storing the alternative $|0\rangle$ vs $|1\rangle$, $S_{\rm coh}$ vanishes. This is an act of erasing one bit of pointer outcome: from the superposition $|+\rangle$ irreversibly one outcome $|0\rangle$ or $|1\rangle$ remains. We propose the hypothesis that the fundamental destruction of this information obeys a thermodynamic constraint analogous to Landauer's principle [9]: the environment of the act—the reservoir where pointer modes and their correlations with the absorber lived—must receive additional heat

$$\Delta Q = kT\ln 2 \tag{2}$$

*for each* objectively fixed single-photon act ($|+\rangle_{\rm ptr} \to |0\rangle$ or $|1\rangle$), above ordinary heating by $E_\gamma$. The excess of case B over A at equal $E_\gamma$—the prediction of Heat from Bit.

**Case C (classical mixture).** The same protocol, but the photon is prepared as $I/2$ in the pointer basis:

$$I/2 = \frac{|0\rangle\langle 0| + |1\rangle\langle 1|}{2} \tag{3}$$

—a diagonal mixture of $|0\rangle$ and $|1\rangle$ without phase correlations: $S_{\text{coh}} = 0$. This is not a superposition of pointer outcomes but a statistical record: the outcome is already "measured" at the level of the pre-act state, only classical uncertainty remains as to *which* of them was realized. In the act of absorption the environment does not *erase* an alternative bit but merely *copies* an already existing classical record; copying classical information is energetically free. There is neither decoherence nor objective collapse in the act.

## 3 Pointer basis and coherence entropy

Upon interaction with the environment the system decoheres into a *pointer basis* $\{|i\rangle\}$—robust outcomes proliferated in the environment (einselection; quantum Darwinism [12]). A classical record is a specific pointer outcome $|i\rangle$. $|+\rangle$ and $I/2$ coincide in $\rho_{\text{ptr}}$ and $\langle\hat{H}\rangle$; a bolometer sees only the diagonal in the pointer basis. The tariff is by coherence in the pointer basis *of the act*, not by $S(\rho)$ abstractly.

**Coherence in the pointer basis.** In the resource theory of coherence [13, 18]:

$$S_{\text{coh}}(\rho) = S(\rho_{\text{ptr}}) - S(\rho), \tag{4}$$

where $\rho_{\text{ptr}}$ is the diagonal of $\rho$ in the pointer basis of the act. For $|+\rangle$ $S_{\text{coh}} = \ln 2$, for $I/2$—zero; this is the *relative entropy of coherence* $C_r(\rho)$ [13, 18] relative to the same basis. Kammerlander and Anders [19] show that the thermodynamics of *projection* into a basis (dephasing, nonselective measurement) is nontrivial only when there is coherence in that basis; for diagonal mixtures $\Delta S = 0$. In an optimal process for $|+\rangle$ from such projection one can extract work $\langle W\rangle = kT\ln 2$. Heat from Bit transfers this distinguishability to the pointer basis of the act and postulates dissipation linked to $S_{\text{coh}}$ upon objective fixation of the outcome.

**Tariff entropy $H$.** Let $H$ denote the *tariff entropy* of the pre-act state in the pointer basis *of the act*; $S(\rho)$, $S_{\text{coh}}$, and $H$ are in *nats*:

$$H(\rho) = S_{\text{coh}}(\rho). \tag{5}$$

$H$ is neither the full $S(\rho)$ nor the classical $-\sum_i p_i \ln p_i$, but precisely the coherence in the pointer basis destroyed upon irreversible fixation of the outcome; it sets the thermodynamic price. Diagonal $\rho_{\text{ptr}}$ $(I/2)$: $H = 0$; pure superposition $|+\rangle_{\text{ptr}}$: $H = \ln 2$.

**Number of bits**:

$$N_{\text{bit}}(\rho) = H(\rho)/\ln 2. \tag{6}$$

**Coherent states of the field** $(\hat{a}|\alpha\rangle = \alpha|\alpha\rangle)$ are eigenstates of the operator $\hat{a}$ [20]; they define the classical amplitude of a mode and a CW laser beam. The term "coherent" here does not mean $S_{\text{coh}}$ in the pointer basis: in $|\alpha\rangle$ there is no superposition of pointer outcomes, only Poisson statistics, $H = 0$; a "photon" of a coherent beam has no separate pre-act $\rho$. "Angle 45°" of a CW laser does not reproduce case B from § 2:

$$|\alpha\rangle_{45^\circ} = |\alpha/\sqrt{2}\rangle_0\,|\mathrm{i}\alpha/\sqrt{2}\rangle_1, \tag{7}$$

not a flux of $|+\rangle_{\text{ptr}}$. For a pure pre-act $\rho = |\psi\rangle\langle\psi|$, $|\psi\rangle = \sum_i c_i|i\rangle$, we have $p_i = |c_i|^2$ and $H = -\sum_i p_i \ln p_i$ (then $S(\rho) = 0$). In the general case $H = S(\rho_{\text{ptr}}) - S(\rho)$; for mixed $I/2$ this is $H = 0$, although $-\sum_i p_i \ln p_i = \ln 2$.

## 4 Thermodynamics of objective collapse

The tariff $H$ (§ 3) is a definition ($H = S_{\rm coh}$ in the pointer basis of the act), not an assumption of the model.

Under *unitary* dephasing $S(\rho_{\rm ptr})$ does not change, the growth of $S(\rho)$ of the subsystem is the decrease of $S_{\rm coh}$ (§ 3); an isolated subsystem under unitary $\pi/2$ does not change $S(\rho)$ (pure remains pure), and a reversible cycle $\pi/2 \to -\pi/2$ restores the state; tariff $H$ is not activated. Heat does *not* follow from the growth of $S(\rho)$ of the subsystem and does not accompany unitary "encryption" of the pointer-basis phase into correlations with the environment.

**Postulate (Heat from Bit).** Testable hypothesis of the work (§ 6): under *objective* collapse with $H > 0$ the jump of the tariff is identified with thermodynamic entropy of the reservoir $S_{\rm th} = kH$ and heat $\Delta Q = T\Delta S_{\rm th} = kT\Delta H$ in the environment at temperature $T$; the pointer outcome is fixed irreversibly. Average over acts

$$\langle Q\rangle = kT\,H = kT\,C_r(\rho). \tag{8}$$

For a given $H$ the distribution of $Q_i$ does *not* follow from this (for example, $Q_i \equiv kTH$ gives the same $\langle Q\rangle$). For a *pure* pre-act $|\psi\rangle = \sum_i c_i|i\rangle$ ($S(\rho) = 0$, $p_i = |c_i|^2$) the model *additionally* adopts a per-trajectory tariff

$$\boxed{Q_i = kT\ln(1/p_i)} \tag{9}$$

—informational "surprise" of outcome $i$ (consistent with the average via $H = -\sum p_i \ln p_i$): a rare outcome "erases" more alternative probability mass. Then $\Delta S_{{\rm th},i} = k\ln(1/p_i)$,

$$\langle Q\rangle = \sum_i p_i\,Q_i = kTH, \qquad {\rm Var}(Q) = (kT)^2\,{\rm Var}(\ln p_i). \tag{10}$$

For a mixed pre-act state ($H = 0$, e.g. $I/2$) the per-act tariff is not activated: $\langle Q\rangle = 0$. At equal $p_i = 1/n$ all outcomes are equivalent: $Q_i = kT\ln n$; for binary $|+\rangle$ ($p_i = \frac{1}{2}$) this is $\Delta Q = kT\ln 2$, $\Delta H = \ln 2$ and $\Delta S_{\rm th} = k\ln 2$ ($\Delta N_{\rm bit} = 1$). Example $p_0 = 0{,}9$, $p_1 = 0{,}1$: an act with outcome 0 releases $kT\ln(10/9) \approx 0{,}11\,kT$, with outcome 1—$kT\ln 10 \approx 2{,}3\,kT$; the average $\approx 0{,}33\,kT = kT\,h_2(0{,}9)$.

The per-outcome tariff is consistent with Born's rule and stochastic thermodynamics [11]. At equal amplitudes ($|+\rangle$) $Q_i$ are equal, no fluctuations; the difference between versions arises only for asymmetric superposition and per-act calorimetry (currently unattainable; § 6).

Proliferation of an already existing record (CNOT, redundancy in the sense of quantum Darwinism [12]) does not create a new bit; unitary entanglement (GHZ, cat states) does not multiply the tariff. Unitary preparation of $|+\rangle_{\rm ptr}$ ($\pi/2$, which-path encoding $|x_0\rangle + |x_1\rangle$, etc.)—reversible operations; no price.

On a series of objective acts (average over outcomes):

$$dH = \ln 2\,dN_{\rm bit}, \qquad \langle\delta Q\rangle = kT\,dH = \mu_{\rm bit}\,dN_{\rm bit}, \qquad \mu_{\rm bit} = kT\ln 2. \tag{11}$$

For a pure pre-act state one act with outcome $i$: $\Delta Q_i = kT\ln(1/p_i)$; $\langle\Delta Q\rangle = kT\,\Delta H = kT\ln 2\,\Delta N_{\rm bit}$. The payment is not $\langle\hat H\rangle$ of the Hamiltonian but the thermodynamic price of fixing a pointer outcome; $\delta Q$ is dissipated in the reservoir, on average—at tariff $\mu_{\rm bit}$.

## 5 Origin of heat energy

In concrete objective-collapse models the non-unitary dynamics itself may specify energy released upon localization: in CSL and GRW [2, 3, 8] the collapse operator raises the mean energy of the system. Then tariff heat $\delta Q = kT\Delta H$ (§ 4) could in principle come from that reservoir. In other models such a reservoir may not be explicit or may be negligible.

We consider the duration of the collapse act $\tau_{\rm col}$—non-unitary compression of the superposition to a single pointer outcome. Under decoherence the system entangles with environmental degrees of freedom of the act—"receptors" of pointer information; the reservoir at the absorber temperature $T$ is populated with modes on the scale $kT$, and the characteristic time of thermal oscillations is $\tau_{\rm th} \equiv \hbar/kT$. In many objective-collapse theories the moment of localization depends on the number of environmental particles decohered with the system or on the effective mass in the superposition [2, 8, 12, 6, 7]. This mass grows as new "receptors" join over a time of order $\tau_{\rm th}$. The value $\tau_{\rm col}$ is set by the particular collapse model; however, heuristically, when the criterion grows with the number of receptors, non-unitary compression must complete within $\tau_{\rm col} \lesssim \tau_{\rm th}$—before one more environmental degree of freedom joins the superposition and the criterion shifts.

In an act of duration $\tau_{\rm col}$ energy is defined no better than $\Delta E \sim \hbar/(2\tau_{\rm col})$ (the Mandelstam–Tamm relation [21] bounds the *rate* of change of an observable); for the full payment $kT\ln 2$ to fit in this window, we link $\tau_{\rm col}$ to the tariff payment $\delta Q = kT\ln 2$:

$$\tau_{\rm col} \lesssim \frac{\hbar}{2kT\ln 2} \approx \frac{\tau_{\rm th}}{2\ln 2}, \tag{12}$$

A single act is not microscopically auditable; over a series of $N$ acts one observes the total excess $\Delta Q_{\rm tot} = N\,kT\Delta H$ (§ 6). Under *unitary* evolution the mean Hamiltonian is conserved, but objective collapse is non-unitary, therefore the mechanism forbidding local energy imbalance is switched off at the moment of pointer-outcome fixation. Thus, in the absence of an explicit reservoir in the collapse model, the energy for tariff heat $\delta Q$ may be drawn from the energy uncertainty of the act duration and dissipated in the thermal reservoir.

## 6 Experimental proposal

The proposed test is differential: the postulated term *adds* heat to ordinary heating ($Q_{\rm out} = Q_{\rm std} + \mu_{\rm bit}\,\Delta N_{\rm bit}$ when $\Delta N_{\rm bit} > 0$), therefore a systematic excess over a control with $H = 0$ is measured. Here $Q_{\rm std}$ is heat without tariff (absorption $W_{\rm in}$, ordinary $Q_{\rm dissip}$); $\Delta N_{\rm bit}$ is the number of pointer bits objectively erased in acts (§ 3): one act from $|+\rangle_{\rm ptr}$ gives $\Delta N_{\rm bit} = 1$, from $I/2_{\rm ptr}$—zero. Heralded MZ—$\sim 10^{10}$ at $R = 10^8$ s$^{-1}$ and $\tau \sim 10^2$ s; all test $\langle Q\rangle = kTH$ (§ 4), not distinguishing fluctuations of $Q_i$ by outcome. A test correlating $Q$ with outcome requires asymmetric superposition and per-act calorimetry and is *currently* unattainable: $\Delta Q_i \sim 10^{-2}$ eV per herald at $T = 300$ K is comparable to single-event bolometer noise, while heralded MZ targets $\langle Q\rangle$ via lock-in and accumulation of $\sim 10^{10}$ acts.

An ordinary CW laser after 50/50 is a product of coherent modes $|\alpha/\sqrt{2}\rangle_L|{\rm i}\alpha/\sqrt{2}\rangle_R$, not a flux of $|+\rangle_{\rm ptr}$ qubits; the same for polarization $|\alpha\rangle_{45^\circ}$ (§ 3). Here—a pure flux of qubits: SPDC + SNSPD herald $\to |1\rangle_{\rm Fock} \to$ Mach–Zehnder $\to$ two pointer zones of absorption.

**Pointer basis and states.** The pointer basis of the act is *localization of absorption*: $|x_0\rangle$, $|x_1\rangle$—two separated zones on one micro-island (left and right arm of the M–Z). Single-photon path state after the first M–Z splitter:

$$|\psi\rangle = \frac{|1,0\rangle + e^{{\rm i}\phi}|0,1\rangle}{\sqrt{2}} \quad \Leftrightarrow \quad |+\rangle_{\rm ptr}:\ H = S_{\rm coh} = C_r = \ln 2. \tag{13}$$

Preparation is unitary ($S(\rho) = 0$); the tariff is activated only upon objective collapse (§ 4), not upon splitting or phase rotation. Control—*classical* path preparation without coherence (not a phase scrambler):

$$\rho_{\rm mix} = \tfrac{1}{2}|1,0\rangle\langle 1,0| + \tfrac{1}{2}|0,1\rangle\langle 0,1| \quad \Leftrightarrow \quad I/2_{\rm ptr}:\ H = 0; \tag{14}$$

diagonals of $\rho_{\text{ptr}}$ coincide with $|+\rangle_{\text{ptr}}$, but $C_r(\rho_{\text{mix}}) = 0$. The control is implemented *before* the interferometer: an optical switch by a classical bit (physically $|1,0\rangle$ or $|0,1\rangle$ per herald); the ensemble is $I/2_{\text{ptr}}$. Fast randomization of $\phi$ does *not* work: in each run the state remains pure $|+\rangle$ with unknown phase ($H = \ln 2$ on the trajectory); $I/2$ is obtained only after tracing over the scrambler (§ 3).

**Why which-path, not polarization.** § 3 shows the same tariff on a single-photon $|+\rangle_{\text{ptr}}$ in the polarization basis; heralded MZ implements an equivalent $|+\rangle_{\text{ptr}}$ in the localization basis $|x_i\rangle$—more convenient for event-triggered calorimetry and control of $I/2_{\text{ptr}}$.

**Setup.**

- **Source**: SPDC in a monolithic or external resonator (cavity SPDC) or several parallel SPDC pairs on a common herald; $\lambda = 1550$ nm, SNSPD on idler; signal $|1\rangle_{\text{Fock}} \to$ MZ (50/50, $\phi$ fixed). Design herald rate $R = 10^8$ s$^{-1}$ (by $\sim$ 1–2 orders of magnitude above typical waveguide SPDC via resonator and/or multiplexing).
- **Absorber**: suspended micro-island (SiN/graphene/Ge–InGaAs), two pointer zones on one island; $T = 300$ K. Thermal link to substrate $G \sim 1$ $\mu$W/K (not mW/K: otherwise the phonon limit NEP $= \sqrt{4k_BT^2G}$ makes lock-in infeasible). Quantum efficiency of the path "herald $\to$ absorption in pointer zone" $\eta = 0{,}85$ (SNSPD, AR coatings, focus on zone).
- **Synchronization**: herald pulse $\to$ bolometer integration window $\ll T_2^*$ of path; lock-in upon switching $|+\rangle_{\text{ptr}} \leftrightarrow I/2_{\text{ptr}}$ at $f_{\text{mod}} \sim 1$ kHz.

**Signal.** For *each* objectively fixed heralded act in channel $|+\rangle_{\text{ptr}}$: $\Delta Q_{\text{bit}} = kT\ln 2 \approx 1{,}8 \times 10^{-2}$ eV ($\sim 2{,}2\%$ of $E_\gamma$ at $\lambda = 1550$ nm); in $I/2_{\text{ptr}}$—zero. At design $R = 10^8$ s$^{-1}$ and $\eta = 0{,}85$ the average excess power

$$\Delta P = \eta\, R\, kT \ln 2 \approx 2{,}4 \times 10^{-13}\ \text{W} \quad (T = 300\ \text{K}). \tag{15}$$

The fraction $\Delta Q_{\text{bit}}/E_\gamma$ is constant; verification is by slope $\Delta P/R$ and event-triggered calorimetry. At $R = 10^8$ s$^{-1}$ contamination from two-pair SPDC events ($g^{(2)}(0) > 0$) dilutes the differential $|+\rangle_{\text{ptr}} - I/2_{\text{ptr}}$; $g^{(2)}(0) \ll 1$ on herald or low duty cycle of the source is required.

**Noise budget.** Phonon limit of the island: NEP $= \sqrt{4k_BT^2G}$. At design $G \sim 1$ $\mu$W/K NEP $\approx 2{,}2\times 10^{-12}$ W/$\sqrt{\text{Hz}}$. Lock-in integration time to SNR $\sim 1$ in power: $\tau \approx (\text{NEP}/\Delta P)^2 \sim 10^2$ s at the $\Delta P$ above.

**Channels (equal $R$, equal $\eta$). Channel $|+\rangle_{\text{ptr}}$:** MZ with fixed $\phi$; $\langle Q\rangle = kT\ln 2$ per heralded act.

**Channel $I/2_{\text{ptr}}$:** same $R$; before MZ classical which-path (L/R switch); $\langle Q\rangle = 0$.

The differential $|+\rangle_{\text{ptr}} - I/2_{\text{ptr}}$ isolates the price for one pointer bit. At $R = 10^8$ s$^{-1}$ over $\tau \sim 10^2$ s $\sim 10^{10}$ acts accumulate and total excess $\Delta Q_{\text{tot}} \sim 10^{-11}$ J.

**What we record.**

1. Event-triggered energy excess $\Delta Q_{\text{bit}} \approx kT\ln 2$ per herald in channel $|+\rangle_{\text{ptr}}$; zero in $I/2_{\text{ptr}}$.
2. Linearity of total $\Delta Q$ in $R$ and $T$; slope $\Delta P/R \propto \eta\, kT\ln 2$.
3. Lock-in peak at $f_{\text{mod}}$; absence of peak at constant $I/2_{\text{ptr}}$.

**Outcomes.** Confirmation: stable differential $|+\rangle_{\text{ptr}} - I/2_{\text{ptr}}$ at $R = 10^8$ s$^{-1}$. Null at SNR $\gg 1$ $\Rightarrow$ bound on tariff $kT\ln 2$.

# 7 Discussion

The Heat from Bit model postulates a single thermodynamic price—for objective collapse ($Q_i = kT\ln(1/p_i)$ for pure coherent states, $\langle Q\rangle = kTH$; § 4); unitarity is free. The existence of such acts is taken as a premise (introduction); the specific criterion for "when" the article does not fix. Tariff entropy $H = S_{\text{coh}}$ in the pointer basis of the act (§ 3); $H = 0$ for equilibrium and P-positive light, $I/2_{\text{ptr}}$, and coherent $|\alpha\rangle$ (§ 3). Experimental test—heralded MZ (§ 6).

**Thermal sector and detailed balance.** Thermal light, solar and CMB radiation in the model are *P-positive* states: a mixture of coherent $|\alpha\rangle$ with Gaussian weight [20]; each component is an eigenstate of $\hat{a}$ ($\hat{a}|\alpha\rangle = \alpha|\alpha\rangle$), in the pointer basis of the act $H = 0$ (§ 3): fixing the outcome does not change $\rho$. This is not an empirical coincidence but a *compatibility condition* with thermodynamics: take a cavity with matter and radiation at one $T$. Spontaneous emission carries away energy $E_\gamma$; if absorption of an equilibrium photon gave $E_\gamma + kT\ln 2$, but emission only $E_\gamma$, detailed balance breaks: the cavity would spontaneously heat from its own blackbody radiation. The model must guarantee $H = 0$ for equilibrium modes *structurally*. Consequence—no conflict with the CMB spectrum (FIRAS) [22] and global energy balance of Earth [23]: $\sim 10^{17}$ W of solar thermal light falls on the surface; even a fraction $\sim 2\%$ tariff per photon would give $\sim 10^{15}$ W excess heating—by factors to tens above the measured energy imbalance ($\sim 0{,}9$ W/m$^2$, $\sim 5\times 10^{14}$ W).

**Spontaneous emission and tariff boundary.** Microscopically a thermal photon and a fluorescence photon are spontaneous emission; the tariff depends on the pre-act $\rho$ of the subsystem of the act (§ 3–§ 4). An equilibrium mode is a P-positive mixture ($H = 0$); controlled test—$|+\rangle_{\text{ptr}}$ (§ 6).

**Relation to existing data.** There is no published *direct* experimental bound on the tariff by $T$ and the postulate $Q_i = kT\ln(1/p_i)$; below—why adjacent measurements neither exclude nor duplicate the proposed test (§ 6).

**Optics.** Single-photon calorimetry on TES (Transition Edge Sensor) [24] resolves photon energy at 1550 nm to $\sim 0{,}1$–1 eV, but at $T_{\text{bath}} \sim 0{,}1$–$0{,}3$ K. At tariff "temperature of the act environment" the excess $kT\ln 2$ at such $T$ ($\sim 10^{-5}$ eV per bit) is three to four orders below TES resolution. Cold TES data do *not* bound the tariff by $T$; the heralded-MZ test (§ 6) with event-triggered calorimetry does not repeat existing single-photon measurements.

**Laser radiometry.** The primary scale of optical power compares warm thermal detectors with cryogenic electrical-substitution radiometers ($T_{\text{bath}} \sim 4$–5 K, tariff $kT\ln 2$ there $\sim 0{,}03\%$ of $E_\gamma$); agreement of services and detector types is 0.02–1.2%. Naive reading "coherent laser $\Rightarrow$ extra $\sim 2\%$ heat at 300 K" does *not* follow from the model: the tariff pays not field coherence in the amplitude basis but superposition *in the pointer basis of the act* (§ 3)—$|+\rangle_{\text{ptr}}$ vs $I/2_{\text{ptr}}$. An ordinary power meter: one beam, one spot, no 50/50 splitter and no controlled inter-zone phase; this is not the $|+\rangle_{\text{ptr}}$ protocol from § 6 and not a bound on the tariff by $T$ for such a state. The proposed heralded-MZ test (§ 6) is the differential $|+\rangle_{\text{ptr}} - I/2_{\text{ptr}}$ on single-photon qubits; radiometry did not reproduce it.

**Bounds on heating of collapse models.** Experiments on GRW/CSL/DP [8, 25] (cold atoms, cantilevers, underground setups) bound *any* excess heating from spontaneous localization—"background" heating of the collapse model itself. Heat from Bit asks something else: how much heat does *irreversible fixation of a pointer outcome* give if before the act there was coherence in the pointer basis ($H > 0$). Existing bounds would refute Heat from Bit only if in those setups many pointer bits were destroyed per event. Usually this does not happen: the system is either already a classical mixture ($H \approx 0$), or a "large" superposition is not in the pointer basis where the tariff is counted—then $H = 0$ and additional heating by Heat from Bit is also

$\approx 0$. Measured zero heating in GRW/CSL/DP tests does not forbid the model at tariff by $T$. In § 6 $H$ and the number of bits per act are made large intentionally.

**Falsification.** Direct test of the model—§ 6.

**Comparison with neighboring approaches.** From measurement engines [26, 27, 28, 29, 30, 31] the Heat from Bit model differs by autonomy of the act without apparatus; from quantum thermodynamics of projections [19]—by the postulate of *objective* collapse and tariff in the pointer basis of the act, not a unitary measuring device; from information reservoirs [32, 33, 34, 35]—by duality (bit as energy carrier at $\Delta S_{\text{tot}} = 0$); from GRW/CSL/DP [8]—by organizing heating by bit flux ($kT \ln 2$ per $\Delta N_{\text{bit}}$): in standard formulations their heating is diffuse and does not correlate with informational acts (diffuse background by itself does not cancel the Heat from Bit model).

**Local first law.** The first law is violated only locally, in a sub-$\tau_{\text{th}}$ act of non-unitary fixation (§ 5); globally the second law is preserved—$\delta Q$ is dissipated in the reservoir with entropy $k\Delta H$. Per-event energy imbalance is below calorimeter resolution; heralded MZ measures systematic excess over the ensemble (§ 6).

**Table 1.** Comparison of standard QM, GRW/CSL/DP models, and the Heat from Bit model.

| | **Standard QM** | **GRW/CSL/DP** | **Heat from Bit** |
|---|---|---|---|
| Objective collapse | no | yes | yes |
| First law | yes | no | locally in act |
| Second law | yes | yes | yes |
| Heat in objective act | 0 | diffuse heating | $\langle Q \rangle = kTH$ |
| Distinctive test | — | Gran Sasso; LISA Pathfinder; ultracold atoms | heralded MZ |

## Declarations

**Funding.** The author received no specific grant from any funding agency.
**Data availability.** Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.
**Competing interests.** The author declares no competing interests.
**Author contributions.** OP conceived the model, performed the calculations, and wrote the manuscript.

## References

[1] F. Károlyházy. Gravitation and quantum mechanics of macroscopic objects. *Il Nuovo Cimento A*, 42:390–402, 1966.

[2] G. C. Ghirardi, A. Rimini, and T. Weber. Unified dynamics for microscopic and macroscopic systems. *Physical Review D*, 34:470–491, 1986.

[3] P. Pearle. Combining stochastic dynamical state-vector reduction with spontaneous localization. *Physical Review A*, 39:2277–2289, 1989.

[4] L. Diósi. Models for universal reduction of macroscopic quantum fluctuations. *Physics Letters A*, 140:451–454, 1989.

[5] R. Penrose. On gravity's role in quantum state reduction. *General Relativity and Gravitation*, 28(5):581–600, 1996. doi:10.1007/BF02105068.

[6] O. Ponfilenok. Objective collapse above the scrambling threshold. Companion manuscript, 2026.

[7] O. Ponfilenok. Branch-dependent proper time. Companion manuscript, 2026. doi:10.5281/zenodo.21497431.

[8] A. Bassi, K. Lochan, S. Satin, T. P. Singh, and H. Ulbricht. Models of wave-function collapse, underlying theories, and experimental tests. *Reviews of Modern Physics*, 85:471–527, 2013.

[9] R. Landauer. Irreversibility and heat generation in the computing process. *IBM Journal of Research and Development*, 5:183–191, 1961.

[10] C. H. Bennett. The thermodynamics of computation—a review. *International Journal of Theoretical Physics*, 21:905–940, 1982.

[11] T. Sagawa and M. Ueda. Second law of thermodynamics with discrete quantum feedback control. *Physical Review Letters*, 100:080403, 2008.

[12] W. H. Zurek. Quantum Darwinism. *Nature Physics*, 5:181–188, 2009.

[13] T. Baumgratz, M. Cramer, and M. B. Plenio. Quantifying coherence. *Physical Review Letters*, 113:140401, 2014.

[14] T. Gorin, T. Prosen, T. H. Seligman, and M. Żnidariç. Dynamics of Loschmidt echoes and fidelity decay. *Physical Review Letters*, 96:150603, 2006.

[15] A. Goussev, R. A. Jalabert, H. M. Pastawski, and D. A. Wisniacki. Loschmidt echo. *Scholarpedia*, 7(8):11687, 2012. doi:10.4249/scholarpedia.11687.

[16] Sahand Hormoz. Quantum collapse and the second law of thermodynamics. *Physical Review E*, 87:022129, 2013. doi:10.1103/PhysRevE.87.022129.

[17] J. A. Wheeler. Information, physics, quantum: The search for links. In *Complexity, Entropy, and the Physics of Information*. Addison-Wesley, 1989.

[18] A. Streltsov, G. Adesso, and M. B. Plenio. Colloquium: Quantum coherence as a resource. *Reviews of Modern Physics*, 89:041003, 2017.

[19] Philipp Kammerlander and Janet Anders. Coherence and measurement in quantum thermodynamics. *Scientific Reports*, 6:22174, 2016. arXiv:1502.02673, doi:10.1038/srep22174.

[20] R. J. Glauber. Coherent and incoherent states of the radiation field. *Physical Review*, 131:2766–2788, 1963.

[21] L. Mandelstam and I. Tamm. The uncertainty relation between energy and time in non-relativistic quantum mechanics. *J. Phys. (USSR)*, 9:249–254, 1945.

[22] D. J. Fixsen, E. S. Cheng, J. M. Gales, J. C. Mather, R. A. Shafer, and E. L. Wright. The cosmic microwave background spectrum from the full COBE FIRAS data set. *The Astrophysical Journal*, 473:576–587, 1996.

[23] N. G. Loeb, B. A. Wielicki, D. R. Doelling, et al. Toward optimal closure of the Earth's top-of-atmosphere radiation budget. *Journal of Climate*, 25:470–491, 2012.

[24] B. Karimi, J. P. Pekola, D. S. Golubev, Yu. A. Pashkin, F. Giazotto, et al. Reaching the ultimate energy resolution of a nanocalorimeter. *Nature Communications*, 11:367, 2020. doi:10.1038/s41467-019-14247-2.

[25] S. Donadi, A. Bassi, C. Curceanu, et al. Underground test of gravity-related wave function collapse models using ultracold atomic beams. *Nature Physics*, 17:528–532, 2021.

[26] C. Elouard, D. Herrera-Martí, B. Huard, and A. Auffèves. Extracting work from quantum measurement in Maxwell's demon engines. *Physical Review Letters*, 118:260603, 2017.

[27] C. Elouard and A. N. Jordan. Efficient quantum measurement engines. *Physical Review Letters*, 120:260601, 2018.

[28] J. Yi, P. Talkner, and Y. W. Kim. Work extraction from a single thermal bath via quantum measurement. *Physical Review E*, 96:022108, 2017.

[29] L. Bresque, G. Súarez, J. Ankerhold, et al. Quantum heat engine boosted by Maxwell's demon feedback. *Physical Review Letters*, 126:120605, 2021.

[30] R. Dassonneville, C. Elouard, R. Cazali, R. Assouly, A. Bienfait, A. Auffèves, and B. Huard. Amplifying microwave pulses with a single qubit engine fueled by quantum measurements. *Physical Review Research*, 8:013228, 2026. arXiv:2501.17069, doi:10.1103/PhysRevResearch.8.013228.

[31] K. Koshihara and K. Yuasa. No-go theorem for quantum heat engines powered purely by quantum measurements in the steady regime, 2026. arXiv:2604.22376.

[32] D. Mandal and C. Jarzynski. Work and information processing in a solvable implementation of Maxwell's demon. *Proceedings of the National Academy of Sciences*, 109:11641–11645, 2012.

[33] D. Mandal, H. T. Quan, and C. Jarzynski. Maxwell's refrigerator: An exactly solvable model. *Physical Review Letters*, 111:030602, 2013.

[34] S. Deffner and C. Jarzynski. Information processing and the second law of thermodynamics: An inclusive, Hamiltonian approach. *Physical Review X*, 3:041003, 2013.

[35] K. Ptaszyński and M. Esposito. Thermodynamics of quantum information flows. *Physical Review Letters*, 122:150603, 2019.