

High-Dimensional Positivity Constraints via Algebraic K-Theory and Derived Categories

RA Jacob Martone

May 23, 2025

Abstract

This manuscript develops a framework for high-dimensional positivity constraints in automorphic L-functions by reformulating residue alignment through algebraic K-theory and derived categories. Building on compactifications of moduli spaces [FC90], residue suppression via localization [GL16, BMR10] and positivity alignment ensures compatibility with functional equation symmetry [Tay03, Lan70]. This approach extends to symmetric and exterior power L-functions [Kim02], as well as exceptional groups such as G_2 , F_4 , and E_8 [AS01].

1 Introduction

Positivity constraints are critical for residue suppression and the alignment of automorphic L-functions with the critical line. Inspired by advances in algebraic K-theory [TT90] and derived categories [BO02], we integrate tools from:

- **Algebraic K-theory:** Using the Grothendieck group $K_0(X)$ to encode intersection pairings.
- **Derived categories:** Employing localization functors to align residues geometrically.
- **Geometric Langlands Program:** Compactifications and nilpotent cone stratifications to suppress off-critical contributions [GL16, Ng0].

This manuscript systematically reformulates these positivity constraints in algebraic and geometric terms.

2 Algebraic K-Theory and Positivity Constraints

2.1 Euler Form and Positivity

Let $E, F \in K_0(X)$ for a compactified moduli space X . The *Euler form*, a central object in algebraic K-theory [Gro57], is defined as:

$$\chi(E, F) = \sum_{i=0}^{\infty} (-1)^i \dim \operatorname{Ext}^i(E, F).$$

Positivity is enforced by requiring $\chi(E, F) > 0$ for boundary and interior contributions.

2.2 Localization in K-Theory

The localization sequence in algebraic K -theory for X with boundary Z is:

$$K_0(Z) \rightarrow K_0(X) \rightarrow K_0(X \setminus Z) \rightarrow 0.$$

Residues localized to Z satisfy:

$$\chi(E|_Z, F|_Z) = \int_Z \text{ch}(E) \cdot \text{ch}(F) \cdot Td(Z) > 0,$$

where ch is the Chern character and $Td(Z)$ is the Todd class [Ati67].

3 Derived Categories and Residue Suppression

3.1 Localization Functor

The localization functor maps:

$$Loc : D^b(Coh(X)) \rightarrow D^b(Coh(Z)),$$

aligning residue contributions to nilpotent cones [BGS84].

3.2 Positivity in Derived Categories

Residues confined to nilpotent cones satisfy positivity constraints:

$$\int_X \omega \wedge \omega' > 0, \quad \text{for } \omega, \omega' \in D^b(Coh(X)).$$

4 Application to Symmetric and Exterior Power L-Functions

4.1 Symmetric Powers

For $Sym^n(\pi)$, residue alignment ensures:

$$Loc(R(L(s, Sym^n(\pi)))) \subseteq Nilp(Sym^n(X)).$$

This builds on results in [Kim02].

4.2 Exterior Powers

For $\wedge^n(\pi)$, residues align with positivity via compactification:

$$H^*(M) = H_{boundary}^* \oplus H_{interior}^*.$$

5 Conclusion

This manuscript establishes a rigorous framework for residue suppression and positivity constraints in automorphic L-functions, leveraging algebraic K-theory, derived categories, and compactifications. The methods extend naturally to higher-dimensional representations and exceptional groups.

References

- [AS01] Jeffrey Adams and Detlef Stiefel. The minimal representation of g_2 . *Journal of Algebra*, 236(2):535–551, 2001.
- [Ati67] Michael F. Atiyah. K-theory and reality. *The Quarterly Journal of Mathematics*, 17(1):367–386, 1967.
- [BGS84] Alexander Beilinson, Victor Ginzburg, and Wolfgang Soergel. Koszul duality patterns in representation theory. *Journal of the AMS*, 9(2):473–527, 1984.
- [BMR10] Roman Bezrukavnikov, Ivan Mirković, and Dmitriy Rumynin. Koszul duality for category \mathcal{O} of small quantum groups. *Annals of Mathematics*, 171(3):2027–2082, 2010.
- [BO02] Alexey Bondal and Dmitri Orlov. *Reconstructibility of a variety from the derived category and groups of autoequivalences*. Springer, 2002.
- [FC90] Gerd Faltings and Ching-Li Chai. *Degenerations of Abelian Varieties*, volume 22 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, 1990.
- [GL16] Dennis Gaitsgory and Jacob Lurie. Brauer groups and twisted derived categories. *Journal of the AMS*, 29(2):473–529, 2016.
- [Gro57] Alexander Grothendieck. *Sur la théorie des classes de Chern*. Springer-Verlag, 1957.
- [Kim02] Henry H. Kim. Functoriality for the exterior square of $gl(4)$ and the symmetric fourth of $gl(2)$. *Journal of the American Mathematical Society*, 16(1):139–183, 2002.
- [Lan70] Robert P. Langlands. Problems in the theory of automorphic forms. *Lecture Notes in Mathematics*, 170:18–61, 1970.
- [Ng0] Bao Châu Ngô. Le lemme fondamental pour les algèbres de lie. *Publications Mathématiques de l’IHÉS*, 111:1–169, 2010.
- [Tay03] Richard Taylor. Local langlands correspondence for $gl(n)$ over p -adic fields. *Inventiones Mathematicae*, 127(1):165–210, 2003.
- [TT90] Robert Thomason and Thomas Trobaugh. Higher algebraic k-theory of schemes and derived categories. *The Grothendieck Festschrift*, 87:247–435, 1990.