Hey there! We are Offering a 100 % Free Guidance for all visitors - like you.

STUDENT INN Career Counseling Center has been serving in all over the Pakistan since 2010.

Our competent team has compiled notes and as well

past papers. We are also providing proper guidance for PPSC lectureship. We have placed past papers freely for the general public.

Brought to you by: Malik Mohsin Hassan & Team

Contact: +92-302-890-0580 Shahid Hussain: Contact: 03009800135

## **Statistics Mcqs Paper 2013**

- Statistics Mcqs Paper 2013
- 1. Two mutually exclusive events are also:
- (a) Dependent events (b) Independent events (c) Exhaustive events (d) None of these
- 2. In which definition of probability, the condition of mutually exclusive and equally likely events is imposed?
- (a) Subjective approach (b) Relative frequency (c) Classical definition (d) None of these
- 3. In Poisson distribution:
- (a) Mean > Variance (b) Mean < Variance (c) Mean Variance (d) None of these
- 4. When sample size n becomes larger and larger and sample estimate tends to the respective population
- parameter, this property of estimators is called:
- (a) Unbiasedness (b) Consistency (c) Reliability (d) None of these
- 5. If the voting preference of 100,000 registered voters are to study, what kind of sampling should be used?
- (a) Simple Random Sampling (b) Quota Sampling (c) Stratified Random Sampling (d) None of these
- 6. If the sampling frame not available then which of the sampling technique could be adopted with ease:
- (a) Simple Random Sampling (b) Stratified Random Sampling
- (c) Cluster Random Sampling (d) None of these
- 7. The complete list of the sampling units is called:
- (a) Sampling Frame (b) Unit List (c) Sample Space (d) None of these
- 8. A cricket captain wins the toss for three consecutive matches. What is the probability that he will call
- correctly for the fourth match?
- (a) 1/2 (b) 1/4 (c) 1/8 (d) None of these
- 9. If X has binomial distribution with parameter p and n then the variance of X is:
- (a) n pq
- (b) np (c)
- 11. Which of the following statement is true for Normal distribution?
- (a) It is skewed to the right (b) It has always a mean of zero and a standard deviation of one
- (c) Its mean, median and mode are equal (d) None of these
- 12. Which of the following statement is true?
- (a) The slope coefficient in regression and the correlation coefficient always have the same sign
- (b) A regression line always passes through origin (c) The correlation coefficient can exceed than 1

- (d) None of these
- •
- 13. Testing of hypotheses is a phase to check the validity of:
- (a) Population parameter (b) Sample estimate (c) Population (d) None of these

•

- 14. Any set of outcomes of a random experiments is called:
- (a) Event (b) Event space (c) Sample points (d) None of these
- 15. When a difference between two groups is statistically significant this means that:
- (a) The difference is statistically real but of little practical significance
- (b) The difference is probably the result of sampling variation
- (c) The difference is not likely to be due to chance variation (d) None of these

• 16. The degree of freedom for two independent samples will be based on:

•

- 17. As the sample size increases:
- (a) The standard deviation of the population decreases (b) The population means increases
- (c) The standard error of the mean decreases (d) None of these

.

- 18. With a lower level of significance, the probability of rejecting a true null hypothesis:
- (a) Remains same (b) Increases (c) Decrease (d) None of these
- 19. Which one is NOT the characteristic of a random experiment:
- (a) It has at least two outcomes (b) The number of all possible outcomes are not known in advance
- (c) It can be repeated any number of times under similar conditions (d) None of these

•

## Statistics Mcqs for Lecturer & Subject Specialist PSC Exam

- If a card is chosen from a standard deck of cards, what is the probability of getting a diamond (♦) or a club(♠)?A 26/52=1/2
  - B 13/52
  - C 20/52
  - D 12/52
  - Question 2
- A listing of the possibile outcomes of an experiment and their corresponding probability is calledA Random Variable
  - B Contingency table
  - C Bayesian table
  - D Probability distribution

E Frequency distribution

Ouestion 3

• If you roll a pair of dice, what is the probability that (at least) one of the dice is a 4 or the sum of the dice is 7?A 4/36

B 13/36

C 21/36

D 15/36

Question 4

• The collection of one or more outcomes from an experiment is called A Probability

B Event

C Random Variable

D Z-Value

E Random Experiment

Ouestion 5

• Which of the following is not a condition of the binomial distribution? A Only 2 possible outcomes

B have constant probability of success

C must have at least 3 trials

D trials must be independent

Ouestion 6

• Which of the following is not an example of a discrete probability distribution? A The sale or purchase price of a house

B The number of bedrooms in a house

C The number of bathrooms in a house

D Whether or not a home has a swimming pool in it.

Question 7

• If a card is chosen from a standard deck of cards, what is the probability of getting a five or a seven? A 4/52

B 1/26

C 8/52

D 1/169

Question 8

• If the occurrence of one event means that another cannot happen, then the events areA Independent

B Mutually Exclusive

C Bayesian

D Empirical

**Question 9** 

• In which approach to probability the outcomes are equally likely to occur? A Classical Probability

B Subjective Probability

C Relative Frequency

D Independent

Question 10

• In a Poisson probability distribution A The mean and variance of the distribution are same (equal)

B The probability of success is always greater than 5

C The number of trials is always less than 5

D It always contains a contingency table

Question 11

• In special rule of addition of probability, the events are always A Independent events

B Mutually Exclusive events

C Bayesian

D Empirical

Question 12

• The joint probability is A The likelihood of two events happening together

B The likelihood of an event happening given that another event has already happened

•	C Based on two mutually exclusive events D Also called Prior probability Question 13 The the special rule of multiplication of probability, the events must be A Independent B Mutually exclusive C Bayesian D Empirical
•	Question 14 Which of the following is not a correct statement about a probability. A It must have a value between 0 and 1 B It can be reported as a decimal or a fraction C A value near 0 means that the event is not likely to occur/happens D It is the collection of several experiments. Binomial distribution is negatively skewed when A p=0 B $p>1/2$ C $p<1/2$ D $p=-1/2$ E $p=1/3$
•	Question 2 Successive trials in binomial distribution are A Dependent B Independent C Equally Likely D Mutually exclusive E None Question 3 If in a binomial distribution n = 1 then E(X) is A q B p C 0
•	D 1 Question 4 In which distribution successive trials are without replacementA Hypergeometric distribution B Binomial distribution C Poisson distribution D Geometric distribution Question 5 Hypergeometric distribution has parametersA 2
•	B 1 C 3 D 4 E No Question 6 Binomial distribution has parameters A Three B Two C One
•	D Four E None Question 7 Binomial distribution is symmetrical when A p = q B p > q C p < q D np > npq Question 8

•	A random variable X has binomial distribution with $n = 10$ and $p = 0.3$ then variance of X is A 10
	B 12
	C 2.1
	D 21
	E None
	Question 9
•	The mean of hypergeometric distribution is Ank
	N
•	B N-k
	n
•	C nN
	k
•	D n+k
	N N
•	Question 10
•	In binomial distribution n=6 and p=0.9, then the value of $P(X=7)$ is A One
	B Less than zero
	C Zero
	D More than zero
	Question 11
•	In a binomial probability distribution it is impossible to find $AP(X < 0)$
	B $P(X = 0)$
	CP(X>0)
	$D P(0 \le X \le n)$
	Question 12
•	The mean, median and mode for binomial distribution will be equal when $A p = 0.5$
	B p< 0.5
	C p > 0.5
	D p = 1
	E None of these
	Question 13
•	Each trial in Binomial distribution has A One Outcome
	B Two Outcome
	C Three Outcome
	D Four Outcome
	Question 14
•	A fair coin is tossed four times, the probability of getting four heads is A 1/4
	B 1/2
	C 1/16
	D 1
	E 0
	Question 15
•	Which of the following is not the property of binomial distributionA n is fixed
	B has two outcomes
	C Trials are independent
	D Probability of success varies from trial to trial
	Question 16
•	The variance of binomial distribution is always A Less than mean
	B Equal to mean
	C Greater than mean
	D Equal to standard deviation
	E None of these
	Question 17

B Binomial distribution C Geometric distribution D None of above	•	The probability of success changes from trial to trial in A Binomial distribution B Geometric distribution C Sampling distribution
The mean of binomial distribution is A npq B np C npq C npq		
B np C npq C npq  D np  In p  Guestion 19 In hypergeometric distribution, the trials areA Independent B Dependent C Collectively Exhaustive D None Question 20 The successive trials are with replacement inA Hypergeometric dis B Binomial distribution C Geometric distribution D None of above Total Area under the curve in probability of density function isA 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) isA Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known asA Pr B Probability function C Distribution Function D Probability Distribution E Probability Density Function Question 4 Probability of occurrence of an event lies betweenA -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar?A 0	•	
D np  D np  None  Question 19 In hypergeometric distribution, the trials areA Independent B Dependent C Collectively Exhaustive D None Question 20 The successive trials are with replacement inA Hypergeometric dis B Binomial distribution C Geometric distribution D None of above Total Area under the curve in probability of density function isA 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) isA Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability function C Distribution Function D Probability Density Function C Distribution Function D Probability Density Function Question 4 Probability of occurrence of an event lies between A-1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0		
D np		C npq
D np		_
D np		7
E nq  Question 19 In hypergeometric distribution, the trials areA Independent B Dependent C Collectively Exhaustive D None Question 20 The successive trials are with replacement inA Hypergeometric dis B Binomial distribution C Geometric distribution D None of above Total Area under the curve in probability of density function isA 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) isA Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability function C Distribution Function C Distribution Function D Probability Distribution E Probability Distribution E Probability Density Function Question 4 Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0		
E nq		D np
E nq		_
Question 19 In hypergeometric distribution, the trials areA Independent B Dependent C Collectively Exhaustive D None Question 20 The successive trials are with replacement inA Hypergeometric disB Binomial distribution C Geometric distribution D None of above Total Area under the curve in probability of density function isA 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) isA Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known asA Pr B Probability Function C Distribution Function D Probability Density Function Question 4 Probability of occurrence of an event lies betweenA -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar?A 0		•
Question 19 In hypergeometric distribution, the trials areA Independent B Dependent C Collectively Exhaustive D None Question 20 The successive trials are with replacement inA Hypergeometric dis B Binomial distribution C Geometric distribution D None of above Total Area under the curve in probability of density function isA 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) isA Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known asA Pr B Probability Density Function C Distribution Function D Probability Density Function E Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar?A 0		E nq
Question 19 In hypergeometric distribution, the trials areA Independent B Dependent C Collectively Exhaustive D None Question 20 The successive trials are with replacement inA Hypergeometric dis B Binomial distribution C Geometric distribution D None of above Total Area under the curve in probability of density function isA 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) isA Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known asA Pr B Probability Density Function C Distribution Function D Probability Density Function E Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar?A 0		<del>-</del>
<ul> <li>In hypergeometric distribution, the trials are A Independent B Dependent C Collectively Exhaustive D None Question 20</li> <li>The successive trials are with replacement in A Hypergeometric dis B Binomial distribution C Geometric distribution D None of above Total Area under the curve in probability of density function is A 0 B -1 C 1 D Infinity Question 2</li> <li>For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3</li> <li>Probability distribution of a random variable is also known as A Pr B Probability Function C Distribution Function D Probability Density Function Question 4</li> <li>Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5</li> <li>What is the probability that a ball drawn at random from a jar? A 0</li> </ul>		$\sqrt{}$
B Dependent C Collectively Exhaustive D None Question 20 The successive trials are with replacement in A Hypergeometric dis B Binomial distribution C Geometric distribution D None of above Total Area under the curve in probability of density function is A 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known as A Pr B Probability Function C Distribution Function D Probability Distribution E Probability Density Function Question 4 Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0		
C Collectively Exhaustive D None Question 20 The successive trials are with replacement in A Hypergeometric dis B Binomial distribution C Geometric distribution D None of above Total Area under the curve in probability of density function is A 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known as A Probability Function C Distribution Function D Probability Distribution E Probability Density Function Question 4 Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0	•	
D None Question 20 The successive trials are with replacement in A Hypergeometric dist B Binomial distribution C Geometric distribution D None of above Total Area under the curve in probability of density function is A 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known as A Probability Function C Distribution Function D Probability Distribution E Probability Density Function Question 4 Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0		
<ul> <li>The successive trials are with replacement in A Hypergeometric dist B Binomial distribution C Geometric distribution D None of above Total Area under the curve in probability of density function is A 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known as A Probability Function C Distribution Function D Probability Density Function E Probability Density Function Question 4 Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0</li> </ul>		
B Binomial distribution C Geometric distribution D None of above Total Area under the curve in probability of density function is A 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known as A Probability Function C Distribution Function D Probability Density Function E Probability Density Function Question 4 Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0		Question 20
C Geometric distribution D None of above Total Area under the curve in probability of density function is A 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known as A Probability Function C Distribution Function D Probability Density Function E Probability Density Function Question 4 Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0	•	The successive trials are with replacement in A Hypergeometric distribution
D None of above Total Area under the curve in probability of density function is A 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known as A President B Probability Function C Distribution Function D Probability Density Function E Probability Density Function Question 4 Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0		
Total Area under the curve in probability of density function is A 0 B -1 C 1 D Infinity Question 2 For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known as A Probability Function C Distribution Function D Probability Density Function E Probability Density Function Question 4 Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0		
C 1 D Infinity Question 2 For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known as A Pr B Probability Function C Distribution Function D Probability Distribution E Probability Density Function Question 4 Probability of occurrence of an event lies between A - 1 and 0 B 0 and 1 C - 1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0		Total Area under the curve in probability of density function is A 0
D Infinity Question 2  For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3  Probability distribution of a random variable is also known as A Probability Function C Distribution Function D Probability Distribution E Probability Density Function Question 4  Probability of occurrence of an event lies between A - 1 and 0 B 0 and 1 C - 1 and 1 D exactly 1 Question 5  What is the probability that a ball drawn at random from a jar? A 0		
<ul> <li>Question 2</li> <li>For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3</li> <li>Probability distribution of a random variable is also known as A Probability Function C Distribution Function D Probability Distribution E Probability Density Function Question 4</li> <li>Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5</li> <li>What is the probability that a ball drawn at random from a jar? A 0</li> </ul>		
<ul> <li>For a random variable X, E(X) is A Harmonic Mean (HM) B Geometric Mean (GM) C Arithmetic Mean (AM) D Non Question 3</li> <li>Probability distribution of a random variable is also known as A Probability Function C Distribution Function D Probability Distribution E Probability Density Function Question 4</li> <li>Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5</li> <li>What is the probability that a ball drawn at random from a jar? A 0</li> </ul>		
C Arithmetic Mean (AM) D Non Question 3 Probability distribution of a random variable is also known as A Pr B Probability Function C Distribution Function D Probability Distribution E Probability Density Function Question 4 Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0	•	
<ul> <li>D Non Question 3</li> <li>Probability distribution of a random variable is also known as A Probability Function</li> <li>C Distribution Function</li> <li>D Probability Distribution</li> <li>E Probability Density Function</li> <li>Question 4</li> <li>Probability of occurrence of an event lies between A -1 and 0</li> <li>B 0 and 1</li> <li>C -1 and 1</li> <li>D exactly 1</li> <li>Question 5</li> <li>What is the probability that a ball drawn at random from a jar? A 0</li> </ul>		
<ul> <li>Question 3</li> <li>Probability distribution of a random variable is also known as A Probability Function</li> <li>C Distribution Function</li> <li>D Probability Distribution</li> <li>E Probability Density Function</li> <li>Question 4</li> <li>Probability of occurrence of an event lies between A -1 and 0</li> <li>B 0 and 1</li> <li>C -1 and 1</li> <li>D exactly 1</li> <li>Question 5</li> <li>What is the probability that a ball drawn at random from a jar? A 0</li> </ul>		
<ul> <li>Probability distribution of a random variable is also known as A Preserved B Probability Function</li> <li>C Distribution Function</li> <li>D Probability Distribution</li> <li>E Probability Density Function</li> <li>Question 4</li> <li>Probability of occurrence of an event lies between A -1 and 0</li> <li>B 0 and 1</li> <li>C -1 and 1</li> <li>D exactly 1</li> <li>Question 5</li> <li>What is the probability that a ball drawn at random from a jar? A 0</li> </ul>		
B Probability Function C Distribution Function D Probability Distribution E Probability Density Function Question 4 Probability of occurrence of an event lies between A-1 and 0 B 0 and 1 C-1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0	•	
D Probability Distribution E Probability Density Function Question 4 Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar? A 0		B Probability Function
<ul> <li>E Probability Density Function</li> <li>Question 4</li> <li>Probability of occurrence of an event lies between A -1 and 0</li> <li>B 0 and 1</li> <li>C -1 and 1</li> <li>D exactly 1</li> <li>Question 5</li> <li>What is the probability that a ball drawn at random from a jar? A 0</li> </ul>		
<ul> <li>Question 4</li> <li>Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5</li> <li>What is the probability that a ball drawn at random from a jar? A 0</li> </ul>		
<ul> <li>Probability of occurrence of an event lies between A -1 and 0 B 0 and 1 C -1 and 1 D exactly 1 Question 5</li> <li>What is the probability that a ball drawn at random from a jar? A 0</li> </ul>		
C -1 and 1 D exactly 1 Question 5 What is the probability that a ball drawn at random from a jar?A 0	•	
D exactly 1 Question 5  What is the probability that a ball drawn at random from a jar? A 0		
Question 5 • What is the probability that a ball drawn at random from a jar?A 0		
• What is the probability that a ball drawn at random from a jar?A 0		
	•	

```
C 0.5
    D_{0}
    E Cannot be determined from given information
    Question 6
  For distribution Function F(X), F(-\infty)=0 and
    F(\infty)=?A 0
    B -1
   C 1
    Question 7
  If C is non-random variable, the E(C) is A Zero
   C 1
   D 2
   E None
    Question 8
• A discrete probability distribution may be represented by A A table
   B A graph
    C A mathematical Equation
   D All of these
    Question 9
• For a probability density function (pdf), the probability of a single point is A 1
    B 2
   C 0
   D Constant
    Question 10
  The probability function is always A Negative
    B Non Negative
    C Positive
    D None
   Question 11
  The distribution function F(X) is represented by A P(X=x)
   B P(X
    C P(X>x)
    D P(X \le x)
   Statistics Mcqs Test
   Statistics Mcqs Test
   (1) A cricket captain wins the toss for three consecutive matches. What is the probability that the will
    call correctly for the fourth match?
    (a) 1/16
    (b) 1/8
```

(c) 1/4

- (d) 1/2
- (e) None of these
- (2) A and B are two independent events. The probability of A is ¼ and B is 1/3. The neither probability of A nor B is:
  - (a) 5/12
  - (b) 1/3
  - (c) 3/4
  - (d) 11/12
  - (e) None of these
- (3) Which one of the following statement is not true?
  - (a) Mutually exclusive events are statistically dependent.
  - (b) Complementary events have probabilities that sum to 1.
  - (c) Opposite events, are statistically independent.
  - (d) An experiment's elementary events are collectively exhaustive and mutually exclusive.
  - (e) None of these
- (4) For a Poisson distribution with standard deviation equals to 2 then mean of the Poisson distribution equals to:
  - (a) 0
  - (b) 1
  - (c) 2
  - (d) 4
  - (e) None of these
- (5) Which of the following could never by described by the binomial distribution?
  - (a) The number of defective items produced by an assembly process.
  - (b) The amount of water used daily by a single household.
  - (c) The number of people in a class who can answer a particular question correctly.
  - (d) All of these
  - (e) None of these
- (6) The area under the normal curve within two standard deviation of the mean is:
  - (a) 68.26 %
  - (b) 95.44 %
  - (c) 99.73 %
  - (d) 99.99 %
  - (e) None of these
- (7) What information is given by a value of the coefficient of determination?
  - (a) Strength of relationship
  - (b) Both strength and direction of relationship
  - (c) Neither strength nor direction of relationship
  - (d) Direction of relationship only
  - (e) None of these
- (8) A complete list of all the sampling units is called:
  - (a) stratum size
  - (b) unit list
  - (c) sampling frame
  - (d) sample design
  - (e) None of these
- (9) If we decrease Type-I error, Type-II error:
  - (a) Remains same
  - (b) Increases
  - (c) Decreases
  - (d) None of these

- (10) To test of equality of means of paired observations, we use:
  - (a) Chi-Square test
  - (b) t-test
  - (c) F-test
  - (d) Z-test
  - (e) None of these

Hey there! We are Offering a 100 % Free Guidance for all visitors - like you. STUDENT INN Career Counseling Center has been serving in all over the Pakistan since 2010. Our competent team has compiled notes and as well past papers. We are also providing proper guidance for PPSC lectureship. We have placed past papers freely for the general public.

Brought to you by: Malik Mohsin Hassan & Team

Contact: +92-302-890-0580 Shahid Hussain: Contact: 03009800135