

# **MULTIPLE MYELOMA** presenting as acute kidney injury secondary to **igA light chain kappa deposits**

H.MEFTAH<sup>1</sup>, O.KALLACH<sup>1</sup>, S.BOUKABOUS<sup>1</sup>, D.BENALI<sup>1</sup>, M.OUYAHIA<sup>1</sup>, Y.BENTATA<sup>1,2</sup>, I.HADDIYA<sup>1,2</sup>

<sup>1</sup> Department of nephrology-dialysis and kidney transplantation , mohammed VI university hospital , faculty of medicine of OUJDA-MOROCCO.

<sup>2</sup> Epidemiology laboratory , faculty of medicine and pharmacy of OUJDA-MOROCCO.


### **BACKGROUND:**


Multiple myeloma (MM) is a commonly encountered hematological malignancy with significant renal involvement. the most common cause of severe acute kidney injury (AKI) in these patients is a tubulopathy which results from very high concentrations of circulating monoclonal immunoglobulin free light chains.

We present a case of IgA MM revealed by severe AKI.

# **CASE PRESENTATION:**

A 69-year-old patient with no medical history, who presented with severe AKI without hypercalcemia requiring hemodialysis, during his hospitalization in





treatment was based on multiple sessions of hemodialysis using a high cut-off dialysis membrane, the evolution was marked by a partial recovery of renal function after 5 cycles of (cyclophosphamidethalidomide and dexamethasone) chemotherapy.

## **DISCUSSION & CONCLUSION:**

Renal monoclonal immunoglobulin deposition disease (MIDD) is a rare disease defined by deposition of monoclonal light chains and/or heavy chains on basement membranes and vascular walls of the kidney. This case highlights the importance of an early diagnosis and assessment to prevent end stage kidney disease.

#### REFERENCES









Figure 1: myeloma involving bone marrow