

# CHIP-FIRING GAMES, POTENTIAL THEORY ON GRAPHS, AND SPANNING TREES

MATTHEW BAKER AND FARBOD SHOKRIEH

**ABSTRACT.** We study the interplay between chip-firing games and potential theory on graphs, characterizing reduced divisors ( $G$ -parking functions) on graphs as the solution to an energy (or potential) minimization problem and providing an algorithm to efficiently compute reduced divisors. Applications include an “efficient bijective” proof of Kirchhoff’s matrix-tree theorem and a new algorithm for finding random spanning trees. The running times of our algorithms are analyzed using potential theory, and we show that the bounds thus obtained generalize and improve upon several previous results in the literature. We also extend some of these considerations to metric graphs.

## 1. INTRODUCTION

Chip-firing games on graphs arise in several different fields of research: in theoretical physics they relate to the “abelian sandpile” or “abelian avalanche” models in the context of self-organized critical phenomena ([5, 28, 29]); in arithmetic geometry, they appear implicitly in the study of component groups of Néron models of Jacobians of algebraic curves ([49, 41, 7]); and in algebraic graph theory they relate to the study of flows and cuts in graphs ([4, 13, 14]). We recommend the recent survey article [39] for a short but more detailed overview of the subject.

There is a close connection between chip-firing games and potential theory on graphs. In this paper, we explore some new aspects of this interplay. Conceptually, this connection should not come as a surprise; in both settings the Laplacian operator plays a crucial rule. However, in chip-firing games an extra “integrality condition” is imposed; in the language of optimization theory, chip-firing games lead to integer programming problems whose associated linear programming relaxations can be solved using potential theory on graphs. Our potential theory methods allow us to prove some new results about chip-firing games and to give new proofs and/or generalizations of some known results in the subject. We also show that certain “ad-hoc” techniques in the literature are naturally explained or unified by our approach.

---

*Date:* September 27, 2016.

*2010 Mathematics Subject Classification.* 05C05, 05C25, 05C50, 05C57, 05C85.

We would like to thank Omid Amini, Jan Draisma, Xander Faber, Greg Lawler, Ye Luo, Sam Payne, and the anonymous referees for their comments on an earlier version of this manuscript. The authors’ work was supported in part by NSF grants DMS-0901487 and CCF-0902717.

Our main potential-theoretic tool is the *energy pairing* (see §3.3), which is a canonical positive definite bilinear form defined on the set of divisors of degree zero<sup>1</sup> on  $G$ . This pairing can be computed using any *generalized inverse* of the Laplacian matrix of  $G$ . The energy pairing can be used to define two functions  $\mathcal{E}_q$  and  $b_q$  (for a fixed vertex  $q$ ) on  $\text{Div}(G)$  which interact in a useful way with *chip-firing moves*; after firing a set of vertices  $A$  (not containing the exceptional vertex  $q$ ), the value of  $\mathcal{E}_q$  goes down by at least the size of the associated cut (see Proposition 4.1), and the value of  $b_q$  goes down by *exactly* the size of  $A$  (see Proposition 4.5).

Chip-firing moves induce a natural equivalence relation on  $\text{Div}(G)$  called *linear equivalence* of divisors. If we once again fix a vertex  $q$ , then a particularly nice set of representatives for linear equivalence classes is given by the  *$q$ -reduced divisors* (see §4.3). We show that the  $q$ -reduced divisor equivalent to a given divisor  $D$  can be characterized as the unique element of  $|D|_q$  (the set of divisors  $D' = \sum_v a'_v(v)$  linearly equivalent to  $D$  for which  $a'_v \geq 0$  whenever  $v \neq q$ ) minimizing the functional  $\mathcal{E}_q$ ; see Theorem 4.12.. A similar result holds with  $\mathcal{E}_q$  replaced by  $b_q$ ; see Theorem 4.14. Using this result we are able to give a new proof of the important fact that there is a unique  $q$ -reduced divisor in each linear equivalence class.

In order to check whether or not a given divisor is  $q$ -reduced, according to the definition (given in §4.3), one needs to check a certain inequality for *all subsets* of  $V(G) \setminus \{q\}$ . But there is in fact a much more efficient procedure called *Dhar's burning algorithm* (after Dhar [28]); see §5.1. Using a modification of Dhar's burning algorithm, it is possible to obtain an "activity preserving" bijection between  $q$ -reduced divisors and spanning trees of  $G$ ; this was originally discovered (using different terminology) by Cori and Le Borgne [26]. In §5.2 we formulate the Cori-Le Borgne algorithm in the language of reduced divisors.

We then turn to the problem of *computing* the  $q$ -reduced divisor equivalent to a given divisor. Dhar's algorithm shows that one can efficiently *check* whether a given divisor is  $q$ -reduced; we show in §5.3 (specifically Algorithm 4) that one can efficiently *find* the  $q$ -reduced divisor equivalent to a given divisor as well. Algorithm 4 can be viewed as the *search* version of Dhar's *decision* algorithm. The main challenge here is the running-time analysis; we use potential theory (specifically, the function  $b_q$ ) to give a bound on the running time of Algorithm 4. As we have already mentioned, the key point is that after firing a set  $A \subseteq V(G) \setminus \{q\}$ , the value of  $b_q$  goes down by exactly the size of  $A$ ; this makes the function  $b_q$  a powerful tool for running-time analysis in chip-firing processes. The seemingly different techniques of Tardos [53], Björner–Lovász–Shor [17], Chung–Ellis [22], van den Heuvel [32], and Holroyd–Levine–Mészáros–Peres–Propp–Wilson [34] all give bounds which are specializations of the running-time bound which we derive using  $b_q$ ; see Remark 5.9.

We next turn to applications of Algorithm 4. The first application (see §6.1) is an "efficient bijective" proof of Kirchhoff's celebrated matrix-tree theorem (stated in a

---

<sup>1</sup>A *divisor* on a finite graph  $G$  is an element  $\sum_{v \in V(G)} a_v(v)$  of the free abelian group  $\text{Div}(G)$  on the set  $V(G)$  of vertices of  $G$ . The *degree* of a divisor is the sum of the  $a_v$  over all  $v$ .

more canonical way than usual in Theorem 6.2). The efficient bijective matrix-tree theorem provides a new approach to the *random spanning tree* problem, which we state in §6.2. This problem has been extensively studied in the literature and there are essentially two known types of algorithms for it: determinant based algorithms (e.g. [31, 23, 38]) and random walk based algorithms (e.g. [19, 1, 55, 36] and [43, Chapter 4]). See Remark 6.3 for possible advantages of our new approach.

Finally, we study some analogous questions in the context of *metric graphs* (or “abstract tropical curves”). While potential theory on finite graphs can be developed purely in the context of linear algebra, potential theory on metric graphs is more conveniently formulated in terms of measure theory. We assume in Appendix A that the reader is familiar with some basic facts concerning potential theory on metric graphs as explained in [6, 8]. Our main new result is Theorem A.7 (the metric graph analogue of Theorem 4.12); using it, we give a new proof of the existence and uniqueness of reduced divisors in the metric graph setting. We conclude with a discussion of Dhar’s algorithm for metric graphs and Luo’s search version of this theorem [42].

The paper is structured as follows. In §2 we fix our notation. In §3 we recall some basic facts from potential theory on graphs and define the energy pairing. The functionals  $\mathcal{E}_q$  and  $b_q$  are introduced in §4 and the interplay between chip-firing dynamics and potential theory on graphs is studied. The algorithmic applications of this interplay, most notably Algorithm 4, are discussed in §5. Some applications of Algorithm 4, including an “efficient bijective” proof of Kirchhoff’s matrix-tree theorem and a new algorithm for the random spanning tree problem, are discussed in §6. Finally, in Appendix A we extend some of our results to metric graphs.

## 2. NOTATION AND TERMINOLOGY

Throughout this paper, a *graph* will mean a finite, connected, unweighted multigraph with no loop edges. The set of vertices of a graph  $G$  is denoted by  $V(G)$  and the set of edges by  $E(G)$ . We let  $n = |V(G)|$  and  $m = |E(G)|$ . For  $A \subseteq V(G)$  and  $v \in A$ , we denote by  $\text{outdeg}_A(v)$  the number of edges between  $v$  and  $V(G) \setminus A$ .

Let  $\text{Div}(G)$  be the free abelian group generated by  $V(G)$ . An element  $\sum_{v \in V(G)} a_v(v) \in \text{Div}(G)$  is called a *divisor* on  $G$ . The coefficient  $a_v$  of  $(v)$  in  $D$  is denoted by  $D(v)$ . For  $D \in \text{Div}(G)$ , let  $\deg(D) = \sum_{v \in V(G)} D(v)$  and let  $\text{Div}^0(G)$  be the subgroup of  $\text{Div}(G)$  consisting of divisors of degree zero. We denote by  $\mathcal{M}(G) = \text{Hom}(V(G), \mathbb{Z})$  the group of integer-valued functions on the vertices. For  $A \subseteq V(G)$ ,  $\chi_A \in \mathcal{M}(G)$  denotes the  $\{0, 1\}$ -valued characteristic function of  $A$ ; note that  $\{\chi_{\{v\}}\}_{v \in V(G)}$  generates  $\mathcal{M}(G)$ .

The *Laplacian operator*  $\Delta : \mathcal{M}(G) \rightarrow \text{Div}(G)$  is defined by  $\Delta(f) = \sum_{v \in V(G)} \Delta_v(f)(v)$ , where

$$\Delta_v(f) = \sum_{\{v, w\} \in E(G)} (f(v) - f(w)).$$

This definition naturally extends to all rational or real-valued functions on vertices.

Let  $\text{Prin}(G)$  (the group of *principal divisors*) be the image of the Laplacian operator  $\Delta : \mathcal{M}(G) \rightarrow \text{Div}(G)$ . It is easy to see that  $\text{Prin}(G) \subseteq \text{Div}^0(G)$  and that both  $\text{Prin}(G)$  and  $\text{Div}^0(G)$  are free abelian groups of rank  $n - 1$ . As a consequence, the quotient group

$$\text{Jac}(G) = \text{Div}^0(G) / \text{Prin}(G)$$

is finite. Following [4],  $\text{Jac}(G)$  is called the *Jacobian* of  $G$ .

Let  $\{v_1, \dots, v_n\}$  be a labeling of  $V(G)$ . With respect to this labeling, the *Laplacian matrix*  $Q$  associated to  $G$  is the  $n \times n$  matrix  $Q = (q_{ij})$ , where  $q_{ii}$  is the degree of vertex  $v_i$  and  $-q_{ij}$  ( $i \neq j$ ) is the number of edges connecting  $v_i$  and  $v_j$ . It is well-known (and easy to verify) that  $Q$  is symmetric, has rank  $n - 1$ , and that the kernel of  $Q$  is spanned by  $\mathbf{1}$ , the all-ones vector (see, e.g., [12, 18]).

The labeling  $\{v_1, \dots, v_n\}$  of  $V(G)$  induces isomorphisms between the abelian groups  $\text{Div}(G)$ ,  $\mathcal{M}(G)$ , and the group of  $n \times 1$  column vectors with integer coordinates. We use  $[D]$  to denote the column vector corresponding to  $D \in \text{Div}(G)$  and  $[f]$  to denote the column vector corresponding to  $f \in \mathcal{M}(G)$ . Under these isomorphisms, the Laplacian operator  $\Delta : \mathcal{M}(G) \rightarrow \text{Div}(G)$  corresponds to the matrix  $Q$  thought of as a homomorphism  $Q : \mathbb{Z}^n \rightarrow \mathbb{Z}^n$ , i.e., for  $f \in \mathcal{M}(G)$  we have  $[\Delta(f)] = Q[f]$ .

### 3. POTENTIAL THEORY

**3.1. Generalized inverses.** A matrix has an inverse only if it is square and has full rank. But one can define a “partial inverse” for any matrix.

**Definition.** Let  $A$  be a matrix. A matrix  $L$  satisfying  $ALA = A$  is called a *generalized inverse* of  $A$ .

Every matrix  $A$  has at least one generalized inverse. In fact more is true: every matrix has a unique *Moore-Penrose pseudoinverse*<sup>2</sup>.

Let  $Q$  be the Laplacian matrix of a (connected) graph  $G$ . Since  $Q$  has rank  $n - 1$ , it does not have an inverse in the usual sense. But there are several natural ways to obtain generalized inverses for  $Q$ . Here are some examples.

**Construction 3.1.** Fix an integer  $1 \leq i \leq n$  and let  $Q_i$  be the invertible  $(n-1) \times (n-1)$  matrix obtained from  $Q$  by deleting  $i^{\text{th}}$  row and  $i^{\text{th}}$  column from  $Q$  ( $Q_i$  is sometimes called the *reduced Laplacian* of  $G$  with respect to  $i$ ). Let  $L_{(i)}$  be the  $n \times n$  matrix obtained from  $Q_i^{-1}$  by inserting a row of all zeros after the  $(i-1)^{\text{st}}$  row and inserting a column of all zeros after the  $(i-1)^{\text{st}}$  column. Then  $L_{(i)}$  is a generalized inverse of  $Q$ . Indeed, one checks that  $QL_{(i)} = I + R_{(i)}$ , where  $I$  is the  $n \times n$  identity matrix and  $R_{(i)}$  has all  $-1$  entries in the  $i^{\text{th}}$  row and is zero elsewhere; as  $R_{(i)}Q = 0$ , we obtain  $QL_{(i)}Q = Q$ .

---

<sup>2</sup>The Moore-Penrose pseudoinverse of  $A$  is a generalized inverse of  $A$  having the following additional properties: (i)  $LAL = L$  and (ii)  $AL$  and  $LA$  are both symmetric. See [10] for additional details.

**Construction 3.2.** Let  $J$  be the  $n \times n$  all 1's matrix. Then  $Q + \frac{1}{n}J$  is nonsingular and  $Q^+ = (Q + \frac{1}{n}J)^{-1} - \frac{1}{n}J$  is a generalized inverse of  $Q$ . In fact,  $Q^+$  is the Moore-Penrose pseudoinverse of  $Q$ , since one easily verifies that  $QQ^+ = Q^+Q = I - \frac{1}{n}J$  and  $Q^+QQ^+ = Q^+$ .

One can use the matrices  $L_{(i)}$  from Construction 3.1 to obtain other generalized inverses for  $Q$ :

**Construction 3.3.** Let  $\mu = (\mu_1, \mu_2, \dots, \mu_n)^T \in \mathbb{R}^n$  satisfy  $\sum_{i=1}^n \mu_i = 1$ . Then  $L_\mu = \sum_{i=1}^n \mu_i L_{(i)}$  is a generalized inverse for  $Q$ . The matrix  $L_\mu$  has the additional property that  $L_\mu \mu = c_\mu \mathbf{1}$  for some  $c_\mu \in \mathbb{R}$ ; this follows from the calculation

$$QL_\mu \mu = (I + \sum_{i=1}^n \mu_i R_{(i)})\mu = \mu - \mu = 0.$$

If  $J$  is the all-1's matrix as in Construction 3.2, then  $G_\mu = L_\mu - c_\mu J$  is also a generalized inverse and has the additional property that  $G_\mu \mu = 0$ . The special case where  $\mu_i = 1/n$  for all  $i$  gives the Moore-Penrose pseudoinverse  $Q^+$  from the previous construction.

**3.2. The  $j$ -function.** We can think of a graph  $G$  as an electrical network in which each edge is a resistor having unit resistance.

**Definition.** For  $p, q, v \in V(G)$ , let  $j_q(p, v)$  denote the electric potential at  $v$  if one unit of current enters a network at  $p$  and exits at  $q$ , with  $q$  grounded (i.e., zero potential).

From a more mathematical point of view,  $j_q(p, \cdot)$  is the unique (rational-valued) solution to the Laplace equation  $\Delta f = (p) - (q)$  satisfying  $f(q) = 0$ ; alternatively, one can define  $j_q(p, v)$  to be the  $(p, v)$ -entry of the matrix  $L_{(q)}$  in Construction 3.1 (see, e.g., [21, 6]; note also that  $(pv||q)$  in [13] is the same as our  $j_q(p, v)$  up to scaling).

The following properties of the  $j$ -function are proved, for example, in [6]:

- $j_q(p, q) = 0$ .
- $j_q(p, v) = j_q(v, p)$ .
- $0 \leq j_q(p, v) \leq j_q(p, p)$ .
- $r(p, q) = j_q(p, p) = j_p(q, q)$ , where  $r(p, q)$  denotes the effective resistance between  $p$  and  $q$ .

**3.3. The energy pairing.** Let  $L$  be any generalized inverse of the Laplacian matrix  $Q$ . Then the bilinear form  $\langle \cdot, \cdot \rangle : \text{Div}^0(G) \times \text{Div}^0(G) \rightarrow \mathbb{Q}$  defined by

$$(3.4) \quad \langle D_1, D_2 \rangle = [D_1]^T L [D_2]$$

is independent of the choice of  $L$ . (Indeed, since there are functions  $f_i \in \text{Hom}(V(G), \mathbb{Q})$  such that  $[D_i] = Q[f_i]$  for  $i = 1, 2$ , we have  $\langle D_1, D_2 \rangle = [f_1]^T Q [f_2]$  and it is easy to check that the right-hand side does not depend on the choice of  $f_1, f_2$ .) We call the canonical bilinear form  $\langle \cdot, \cdot \rangle$  the *energy pairing* on  $\text{Div}^0(G)$ .

**Lemma 3.5.** *The energy pairing is positive definite.*

*Proof.* Let  $B$  be the incidence matrix of the graph. Then  $Q = BB^T$ , so if  $[D] = Q[f]$  we have  $\langle D, D \rangle = [f]^T BB^T [f] = \|B^T[f]\|_2^2$ .  $\square$

**Definition.** The *energy* of a divisor  $D \in \text{Div}^0(G)$  is

$$\mathcal{E}(D) = \langle D, D \rangle = [D]^T L[D].$$

Since the energy pairing is positive definite,  $\mathcal{E}(D) \geq 0$  with equality iff  $D = 0$ .

*Remark 3.6.* The name “energy pairing” comes from the fact that if  $D \in \text{Div}^0(G)$  represents an external current in the network, where  $D(v)$  units of current enter the network at  $v$  if  $D(v) > 0$  and  $-D(v)$  units of current exit the network at  $v$  if  $D(v) < 0$ , then  $\mathcal{E}(D)$  is precisely the total energy dissipated (per unit time) in the network.

We emphasize that the energy pairing is independent of the choice of  $L$  only because the divisors are assumed to have degree zero. One can extend the energy pairing to arbitrary divisors by fixing a vertex  $q$  and defining the *q-energy pairing* by

$$\langle D, E \rangle_q = \langle D - \deg(D)(q), E - \deg(E)(q) \rangle$$

for  $D, E \in \text{Div}(G)$ .

### 3.4. The maximum principle.

**Lemma 3.7.** *Let  $f \in \mathcal{M}(G)$ . Let  $A_{\max}$  (resp.  $A_{\min}$ ) be the set of vertices where  $f$  achieve its maximum (resp. minimum) value. Then*

- (a) *For  $v \in A_{\max}$ ,  $\Delta_v(f) \geq \text{outdeg}_{A_{\max}}(v)$ .*
- (b) *For  $v \in A_{\min}$ ,  $\Delta_v(f) \leq -\text{outdeg}_{A_{\min}}(v)$ .*

*Proof.* For part (a) let  $v \in A_{\max}$ . For an edge  $e = vw$ , if  $w \in A_{\max}$  then  $f(v) = f(w)$ , and if  $w \notin A_{\max}$  then  $f(v) - f(w) \geq 1$ . Since  $\Delta_v(f) = \sum_{\{v,w\} \in E(G)} (f(v) - f(w))$ , the result follows. Part (b) follows from part (a) by replacing  $f$  with  $-f$ .  $\square$

One obtains the following well-known corollary:

**Corollary 3.8** (Maximum principle). *Suppose  $f \in \mathcal{M}(G)$  is nonconstant. Then  $f$  achieves its maximum (resp. minimum) value at a vertex  $v$  for which  $\Delta_v(f) > 0$  (resp.  $\Delta_v(f) < 0$ ).*

## 4. CHIP-FIRING DYNAMICS AND POTENTIAL THEORY

**4.1. Chip-firing dynamics on graphs.** Following [7], we define an equivalence relation  $\sim$  (called *linear equivalence*) on the group  $\text{Div}(G)$  as follows:

**Definition.** For  $D_1, D_2 \in \text{Div}(G)$ ,  $D_1 \sim D_2$  if and only if  $D_1 - D_2$  is in the image of  $\Delta : \mathcal{M}(G) \rightarrow \text{Div}(G)$ .

This equivalence relation is closely related to notion of *chip-firing games* or *dollar games* (see, e.g., [17, 13, 14, 7, 28, 5]). Given a divisor  $D \in \text{Div}(G)$ , one can view the integer  $D(v)$  as the number of *dollars* assigned to the vertex  $v$ . If  $D(v) < 0$  then  $v$  is said to be *in debt*. A *chip-firing move* consists of choosing a vertex and having it either

borrow one dollar from each of its neighbors or give (“fire”) one dollar to each of its neighbors. For  $D_1, D_2 \in \text{Div}(G)$ ,  $D_1 \sim D_2$  if and only if starting from the configuration  $D_1$  one can reach the configuration  $D_2$  through a sequence of chip-firing moves.

**4.2. Chip-firing moves and the energy pairing.** For  $D \in \text{Div}(G)$  we define  $\mathcal{E}_q(D) = \langle D, D \rangle_q$ . The following two propositions relate the energy pairing and chip-firing moves, and will be used in the next section.

**Proposition 4.1.** (a) *If  $E = D + \Delta(f) \in \text{Div}(G)$  for some  $f \in \mathcal{M}(G)$ , then*

$$\mathcal{E}_q(E) = \mathcal{E}_q(D) + \sum_{v \in V(G)} (D + E)(v) \cdot f(v) - 2 \deg(E) \cdot f(q) .$$

(b) *If  $E = D - \Delta(\chi_A) \in \text{Div}(G)$  for some  $A \subseteq V(G) \setminus \{q\}$ , then*

$$\mathcal{E}_q(E) = \mathcal{E}_q(D) - \sum_{v \in A} (D + E)(v) .$$

*If, moreover,  $E$  is effective on  $A$  (i.e.  $E(v) \geq 0$  for  $v \in A$ ), then*

$$(4.2) \quad \mathcal{E}_q(E) \leq \mathcal{E}_q(D) - \lambda(A)$$

*where  $\lambda(A)$  denotes the size of the  $(A, G \setminus A)$ -cut (i.e., the number of edges having one end in  $A$  and the other end in  $V(G) \setminus A$ ).*

*Proof.* (a) Let  $\deg(E) = d$ . Then  $\deg(D) = d$  as well. We can find a  $\mathbb{Q}$ -valued function  $g$  so that  $[E + D - 2d(q)] = Q[g]$ . Then

$$\begin{aligned} \mathcal{E}_q(E) &= \langle D + \Delta(f) - d(q), D + \Delta(f) - d(q) \rangle \\ &= \mathcal{E}_q(D) + \langle D + E - 2d(q), \Delta(f) \rangle = \mathcal{E}_q(D) + (Q[g])^T LQ[f] \\ &= \mathcal{E}_q(D) + [g]^T QLQ[f] = \mathcal{E}_q(D) + [g]^T Q[f] \\ &= \mathcal{E}_q(D) + (Q[g])^T [f] = \mathcal{E}_q(D) + [D + E - 2d(q)]^T [f] \\ &= \mathcal{E}_q(D) + \sum_{v \in V(G)} (D + E)(v) \cdot f(v) - 2d \cdot f(q) . \end{aligned}$$

(b) For the first statement let  $f = -\chi_A$  in part (a).  $E(v) \geq 0$  for  $v \in A$  means that  $D(v) \geq \Delta_v(\chi_A) = \text{outdeg}_A(v)$  for  $v \in A$ . So  $\sum_{v \in A} (D + E)(v) \geq \sum_{v \in A} \text{outdeg}_A(v) = \lambda(A)$ .  $\square$

*Remark 4.3.* Proposition 4.1 (a) can be used to give a new solution to the well-known *Pentagon Problem*<sup>3</sup>: “To each vertex of a regular pentagon an integer is assigned in such a way that the sum of all five numbers is positive. If three consecutive vertices are assigned the numbers  $x, y, z$  respectively, and  $y < 0$ , then the following operation is allowed : the numbers  $x, y, z$  are replaced by  $x + y, y, z + y$ , respectively. Such an operation is performed repeatedly as long as at least one of the five numbers is negative. Determine whether this procedure necessarily comes to an end after a finite number of steps”.

<sup>3</sup>Problem 3, 27th IMO 1986. We refer the reader to [56, 54, 52, 2, 47] for some discussions, solutions, and generalizations of this problem.

To see that this process stops for any  $n$ -cycle, let  $D$  be the starting configuration and assume that  $s = \deg(D) \geq 1$ . It follows from Proposition 4.1 (a) that the quantity

$$\mathbb{E}(D) = \sum_{q \in V(G)} \mathcal{E}_q(D) = \sum_{q, p, v \in V(G)} D(p) j_q(p, v) D(v)$$

goes down by exactly  $-2s \cdot D(v) > 0$  after each basic move with  $y = D(v) < 0$ . Since the energy pairing is positive definite, we always have  $\mathbb{E} \geq 0$ , and thus the procedure will necessarily come to an end after a finite number of steps. Note that this method gives a way to compute the number of steps as well.

**Definition.** Let  $\mathbf{1}$  denote the all-1's divisor. For  $D \in \text{Div}(G)$  and  $q \in V(G)$ , we define  $b_q(D) = \langle \mathbf{1}, D \rangle_q$ .

*Remark 4.4.*  $b_q(D)$  is the “total potential” induced by the external current source corresponding to the divisor  $D - \deg(D)(q) \in \text{Div}^0(G)$ .

**Proposition 4.5.** (a) *If  $E = D + \Delta(f) \in \text{Div}(G)$  for some  $f \in \mathcal{M}(G)$ , then*

$$(4.6) \quad b_q(E) = b_q(D) + \sum_{v \in V(G)} (f(v) - f(q)) .$$

(b) *If  $E = D - \Delta(\chi_A) \in \text{Div}(G)$  for some  $A \subseteq V(G) \setminus \{q\}$ , then*

$$(4.7) \quad b_q(E) = b_q(D) - |A| ,$$

where  $|A|$  is the cardinality of the set  $A$ . Thus  $b_q(\cdot)$  is a monovariant<sup>4</sup>.

*Proof.* (a) We can find a  $\mathbb{Q}$ -valued function  $g$  so that  $[\mathbf{1} - n(q)] = Q[g]$ , where  $n = |V(G)|$ . Then

$$\begin{aligned} \langle \mathbf{1}, E \rangle_q &= \langle \mathbf{1}, D \rangle_q + \langle \mathbf{1}, \Delta(f) \rangle_q \\ &= \langle \mathbf{1}, D \rangle_q + (Q[g])^T L Q[f] = \langle \mathbf{1}, D \rangle_q + [g]^T Q L Q[f] \\ &= \langle \mathbf{1}, D \rangle_q + [g]^T Q[f] = \langle \mathbf{1}, D \rangle_q + (Q[g])^T [f] \\ &= \langle \mathbf{1}, D \rangle_q + [\mathbf{1} - n(q)]^T [f] \\ &= \langle \mathbf{1}, D \rangle_q + \sum_{v \in V(G)} (f(v) - f(q)) . \end{aligned}$$

(b) follows from part (a) by setting  $f = -\chi_A$ . □

The  $\mathbb{Q}$ -valued function  $g : V(G) \rightarrow \mathbb{Q}$  in the proof of Proposition 4.5 (a) can be computed explicitly, and this gives a useful formula for  $b_q$ :

**Lemma 4.8.** *Let  $g_q : V(G) \rightarrow \mathbb{Q}$  be the unique function such that  $\Delta(g_q) = \sum_v (v) - n(q)$  and  $g_q(q) = 0$ . Then:*

$$(a) \quad g_q(v) = \sum_{p \in V(G)} j_q(p, v) .$$

---

<sup>4</sup>A quantity which either only goes up or only goes down under some process.



(b) For any divisor  $D \in \text{Div}(G)$ ,

$$(4.9) \quad b_q(D) = \sum_v g_q(v)D(v) = \sum_v \sum_p j_q(p, v)D(v) .$$

In particular, if  $D(v) \geq 0$  for  $v \neq q$ , then  $b_q(D) \geq 0$ .

*Proof.* Part (a) is easy and is left as an exercise. Part (b) follows from Construction 3.1 and the definition of the energy pairing. Alternatively, let  $[\mathbf{1} - n(q)] = Q[g_q]$  and  $[D - d(q)] = Q[f]$  where  $\deg(D) = d$ . Then

$$\begin{aligned} \langle \mathbf{1}, D \rangle_q &= (Q[g_q])^T LQ[f] \\ &= [g_q]^T Q[f] \\ &= [g_q]^T [D - d(q)] = \sum_v g_q(v)D(v) \\ &= \sum_v \sum_p j_q(p, v)D(v) . \end{aligned}$$

The second statement follows because  $j_q(p, v) \geq 0$  and  $j_q(p, q) = 0$ .  $\square$

**4.3. Reduced divisors.** A nice set of representatives for equivalence classes of divisors are given by the “reduced divisors”.

**Definition.** Fix a vertex  $q \in V(G)$ . A divisor  $D \in \text{Div}(G)$  is called *q-reduced*<sup>5</sup> if it satisfies the following two conditions:

- (i)  $D(v) \geq 0$  for all  $v \in V(G) \setminus \{q\}$ .
- (ii) For every non-empty subset  $A \subseteq V(G) \setminus \{q\}$ , there exists a vertex  $v \in A$  such that  $D(v) < \text{outdeg}_A(v)$ .

In other words, every vertex outside  $q$  is nonnegative but simultaneously firing all the vertices in any non-empty subset  $A$  of  $V(G)$  which is disjoint from  $q$  will result in some vertex becoming negative.

The significance of reduced divisors comes primarily from the fact that for every  $D \in \text{Div}(G)$ , there is a *unique*  $q$ -reduced divisor  $D'$  such that  $D' \sim D$ . This basic fact was discovered independently (in different guises) by several different authors (see, e.g., [29, 27, 48, 7]). We give a new proof of this result in Corollary 4.13 below.

We wish to study reduced divisors from a potential-theoretic point of view. Fix a distinguished vertex  $q$  and define

$$|D|_q = \{E \in \text{Div}(G) \mid E \sim D, E(v) \geq 0 \text{ for all } v \neq q\} .$$

**Lemma 4.10.** For every  $D \in \text{Div}(G)$  and any vertex  $q$ , the set  $|D|_q$  is non-empty.

---

<sup>5</sup>Reduced divisors are essentially the same thing as *G-parking functions* [48] or *superstable configurations* [34].

*Proof.* Pick an ordering  $\prec$  on  $V(G)$  with the property that  $q$  is the first vertex in the ordering and every  $v \neq q$  has a neighbor  $w$  with  $w \prec v$ . Starting from the last vertex in the ordering, we can inductively make all vertices other than  $q$  effective by replacing  $D$  with  $D - k\Delta(\chi_w)$  for some neighbor  $w \prec v$  and some sufficiently large integer  $k$ .  $\square$

**Lemma 4.11** (Principle of least action). *Let  $D$  be a  $q$ -reduced divisor. Assume  $E \sim D$  and write  $D = E + \Delta(f)$ .*

- (a) *If  $E \in |D|_q$ , then  $f(v) \leq f(q)$  for all  $v \in V(G)$ .*
- (b) *If  $E + \Delta(g) \in |D|_q$ , then  $f(v) - f(q) \leq g(v) - g(q)$  for all  $v \in V(G)$ .*

*Proof.* Part (a) is a consequence of Lemma 3.7. If  $f$  does not achieve its global maximum at  $q$ , then  $A_{\max} \subseteq V(G) \setminus \{q\}$  and for all  $v \in A_{\max}$  we have  $\Delta_v(f) \geq \text{outdeg}_{A_{\max}}(v)$ . Since  $D$  is  $q$ -reduced, there must be a vertex  $u \in A_{\max}$  such that  $D(u) < \text{outdeg}_{A_{\max}}(u)$ . But then  $E(u) = D(u) - \Delta_u(f) < 0$ , contradicting the assumption that  $E \in |D|_q$ .

For (b), let  $E' = E + \Delta(g)$ . Then  $D = E' + \Delta(f - g)$  and part (a) gives  $f(v) - g(v) \leq f(q) - g(q)$ .  $\square$

**Theorem 4.12.** *Fix  $q \in V(G)$  and let  $D \in \text{Div}(G)$ . Then  $D$  is  $q$ -reduced if and only if  $D \in |D|_q$  and  $\mathcal{E}_q(D) < \mathcal{E}_q(D')$  for all  $D' \neq D$  in  $|D|_q$ .*

*Proof.* If  $D$  is  $q$ -reduced, then  $D \in |D|_q$ . Let  $E \in |D|_q$ . Then  $\mathcal{E}_q(D) \leq \mathcal{E}_q(E)$ ; write  $D = E + \Delta(f)$  with  $f(q) = 0$ . By Lemma 4.11 (a), we have  $f(v) \leq 0$ . By Proposition 4.1 (a), we have

$$\mathcal{E}_q(D) = \mathcal{E}_q(E) + \sum_{v \neq q} (D + E)(v) \cdot f(v) \leq \mathcal{E}_q(E).$$

Now assume  $D \in |D|_q$  and  $\mathcal{E}_q(D) \leq \mathcal{E}_q(E)$  for all  $E \in |D|_q$  but  $D$  is not  $q$ -reduced. Then there exists a non-empty set  $A \subseteq V(G) \setminus \{q\}$  such that  $D_1 = D - \Delta(\chi_A) \in |D|_q$ , so Proposition 4.1 (b) implies

$$\mathcal{E}_q(D_1) = \mathcal{E}_q(D) - \lambda(A) \leq \mathcal{E}_q(D) - 1.$$

It follows that if  $\mathcal{E}_q(D_1) = \mathcal{E}_q(D_2) \leq \mathcal{E}_q(E)$  for all  $E \in |D|_q$ , then both  $D_1$  and  $D_2$  are  $q$ -reduced. By Lemma 4.11 (a), if  $D_2 = D_1 + \Delta(f)$  with  $f(q) = 0$ , then  $f(v) \leq 0$  for all  $v \neq q$ . Similarly  $-f(v) \leq 0$  for all  $v \neq q$ , so  $f = 0$  and  $D_1 = D_2$ .  $\square$

**Corollary 4.13.** *Fix  $q \in V(G)$  and let  $D \in \text{Div}(G)$ . Then there is a unique  $q$ -reduced divisor  $D' \in \text{Div}(G)$  linearly equivalent to  $D$ .*

*Proof.* It follows from Lemma 4.10, Proposition 4.1 (b), and Lemma 3.5 that we may choose  $D' \in \text{Div}(G)$  such that  $\mathcal{E}_q(D') \leq \mathcal{E}_q(D'')$  for all  $D'' \in |D|_q$ . By Theorem 4.12,  $D'$  is the unique  $q$ -reduced divisor linearly equivalent to  $D$ .  $\square$

An analogue of Theorem 4.12 holds with  $\mathcal{E}_q$  replaced by  $b_q$ :

**Theorem 4.14.** *Fix  $q \in V(G)$  and let  $D \in \text{Div}(G)$ . Then  $D$  is  $q$ -reduced if and only if  $D \in |D|_q$  and  $b_q(D) < b_q(D')$  for all  $D' \neq D$  in  $|D|_q$ .*

*Proof.* Note that by Lemma 4.8 (b), the function  $b_q$  does have a minimum in  $|D|_q$ . The rest of the proof mirrors the proof of Theorem 4.12.

If  $D$  is  $q$ -reduced, then  $D \in |D|_q$  by definition. Let  $E \in |D|_q$  be another divisor. Write  $D = E + \Delta(f)$  with  $f(q) = 0$ . By Lemma 4.11 (a) we have  $f(v) \leq 0$ . Now, by Proposition 4.5 (a), we have

$$b_q(D) = b_q(E) + \sum_{v \neq q} f(v) \leq b_q(E) .$$

Now assume  $D \in |D|_q$  and  $b_q(D) \leq b_q(E)$  for all  $E \in |D|_q$  but  $D$  is not  $q$ -reduced. Then there exists a non-empty set  $A \subseteq V(G) \setminus \{q\}$  such that  $D_1 = D - \Delta(\chi_A) \in |D|_q$ . But then Proposition 4.5 (b) gives the contradiction

$$b_q(D_1) = b_q(D) - |A| \leq b_q(D) - 1 .$$

It follows that if  $b_q(D_1) = b_q(D_2) \leq b_q(E)$  for all  $E \in |D|_q$ , then both  $D_1$  and  $D_2$  are  $q$ -reduced. By Lemma 4.11 (a), if  $D_2 = D_1 + \Delta(f)$  with  $f(q) = 0$ , then  $f(v) \leq 0$  for all  $v \neq q$ . Similarly  $-f(v) \leq 0$  for all  $v \neq q$ , so  $f = 0$  and  $D_1 = D_2$ .  $\square$

*Remark 4.15.* Theorem 4.14 remains true if  $b_q(D) = \langle \mathbf{1}, D \rangle_q$  is replaced by  $\langle \mathbf{h}, D \rangle_q$  for any “ $\mathbb{R}$ -divisor”  $\mathbf{h}$  (i.e.  $h(v) \in \mathbb{R}$ ), provided that  $\mathbf{h}(v) > 0$  for  $v \neq q$ .

## 5. ALGORITHMIC ASPECTS OF REDUCED DIVISORS

**5.1. Dhar’s algorithm.** Let  $D$  be a divisor on the graph  $G$ . In order to check whether or not  $D$  is  $q$ -reduced using the definition, one needs to check for *all subsets*  $A \subseteq V(G) \setminus \{q\}$  whether or not there is a vertex  $v \in A$  such that  $D(v) < \text{outdeg}_A(v)$ . But there is in fact a much more efficient procedure called *Dhar’s burning algorithm* (after Dhar [28]).

The idea behind Dhar’s algorithm is as follows. Think of the edges of  $G$  as being made of a flammable material. A fire starts at vertex  $q$  and proceeds along each edge adjacent to  $q$ . At each vertex  $v \neq q$ , there are  $D(v)$  firefighters, each of whom can control fires in a single direction (i.e., edge) leading into  $v$ . Whenever there are fires approaching  $v$  in more than  $D(v)$  directions, the fire burns through  $v$  and proceeds to burn along all the other edges incident to  $v$ . The divisor  $D$  is  $q$ -reduced iff the fire eventually burns through every vertex of  $G$ .

More formally, Dhar’s algorithm is stated in Algorithm 1.

The complexity of Dhar’s algorithm is  $O(n^2)$ : there are at most  $n$  iterations, and at most  $n$  inequalities are tested in each iteration.

**5.2. The Cori-Le Borgne algorithm.** Using a modification of Dhar’s burning algorithm, it is possible to obtain an “activity preserving” bijection between  $q$ -reduced divisors (of a given degree  $d$ ) on  $G$  and spanning trees of  $G$ . This is more or less just a restatement of the work of Cori and Le Borgne in [26] in the language of reduced divisors; however, by using the Cori-Le Borgne algorithm in conjunction with the results of §3 and §4, we are able to obtain new results.

**Input:** A divisor  $D \in \text{Div}(G)$ , and a vertex  $q \in V(G)$ .  
**Output:** TRUE if  $D$  is  $q$ -reduced, and FALSE if  $D$  is not  $q$ -reduced.  
**if**  $D(v) < 0$  for some  $v \in V(G) \setminus \{q\}$  **then** output FALSE and Stop.  
 Let  $A_0 = V(G)$  and  $v_0 = q$ .  
**for**  $1 \leq i \leq n - 1$  **do**  
     Let  $A_i = A_{i-1} \setminus \{v_{i-1}\}$ .  
     **if** for all  $v \in A_i$ ,  $D(v) \geq \text{outdeg}_{A_i}(v)$  **then** output FALSE and Stop.  
     **else** let  $v_i \in A_i$  be any vertex with  $D(v_i) < \text{outdeg}_{A_i}(v_i)$ .  
**end**  
 Output TRUE.

**Algorithm 1:** Dhar’s Burning Algorithm

The idea behind the Cori-Le Borgne algorithm is as follows. In our original formulation of Dhar’s algorithm, we burned through multiple edges at once. We now use an ordering of  $E(G)$  to break ties and implement a “controlled burn” in which only one edge at a time is burnt.

Thus, fix a total order on  $E(G)$  and suppose we are given a  $q$ -reduced divisor  $D$ . We run Dhar’s burning algorithm on  $D$ , starting with a fire at  $q$ . However, any time there are multiple unburnt edges which are eligible to burn, we always choose the *smallest* one. Whenever the fire burns through a vertex  $v$ , we *mark* the edge along which the fire traveled just before burning through  $v$ . Since  $D$  is  $q$ -reduced, the fire eventually burns through every vertex of  $G$ . The set of marked edges is connected, has cardinality  $n - 1$ , and covers all vertices and thus forms a *spanning tree*  $T_D$  of  $G$ . We thus obtain an association  $\{\text{reduced divisors}\} \rightsquigarrow \{\text{spanning trees}\}$  (Algorithm 2).

The remarkable fact discovered by Cori and Le Borgne is that this association is a *bijection*. The inverse map is also completely explicit and can be described as follows (using the same total order on  $E(G)$ ). Suppose we are given a spanning tree  $T$  in  $G$ . A controlled burn starts at the vertex  $q$  and, as before, any time there are multiple unburnt edges eligible to burn we choose the smallest one. The difference is that now the firefighters at  $v$  can control incoming fires in every direction except for those corresponding to edges of  $T$ . Thus the fire burns through a vertex  $v \neq q$  exactly when it travels along an edge  $e \in T$  from some (burnt) vertex  $w$  to  $v$ . At the moment when  $v$  is burned through, we set  $D(v)$  equal to  $|\{\text{burnt edges adjacent to } v\}| - 1$ . Eventually the fire will burn through every vertex and a nonnegative integer  $D(v)$  will have been assigned to each vertex  $v \neq q$ . The value of  $D(q)$  is determined by requiring that  $\deg(D) = d$ . It turns out that the resulting divisor  $D$  is  $q$ -reduced, so we obtain an association  $\{\text{spanning trees}\} \rightsquigarrow \{\text{reduced divisors}\}$  (Algorithm 3) which one checks is *inverse* to Algorithm 2.

**Theorem 5.1.** *The association given by Algorithms 2 and 3 is a bijection. More precisely:*

**Input:**

$G = (V, E)$  is graph with a fixed ordering on  $E$ ,

$q \in V(G)$ ,

$D = \sum_v a_v(v)$ , a  $q$ -reduced divisor of degree  $d$ .

**Output:**

$T_D$  a spanning tree of  $G$ .

**Initialization:**

$X = \{q\}$  (“burnt” vertices),

$R = \emptyset$  (“burnt” edges),

$T = \emptyset$  (“marked” edges).

**while**  $X \neq V(G)$  **do**

$f = \min\{e = \{s, t\} \in E(G) \mid e \notin R, s \in X, t \notin X\}$ ,

    let  $v \in V(G) \setminus X$  be the vertex incident to  $f$ ,

**if**  $a_v = |\{e \text{ incident to } v \mid e \in R\}|$  **then**

$X \leftarrow X \cup \{v\}$ ,

$T \leftarrow T \cup \{f\}$ ,

**end**

$R \leftarrow R \cup \{f\}$

**end**

$\text{ExtAct} = E \setminus R$ ,

$\text{ExtPass} = R \setminus T$ ,

Output  $T_D = T$ .

**Algorithm 2:** Reduced divisor to spanning tree.

- (i) For any  $q$ -reduced divisor  $D$  of degree  $d$ , Algorithm 2 outputs a spanning tree  $T_D$  of  $G$ .
- (ii) For any spanning tree  $T$ , Algorithm 3 outputs a  $q$ -reduced divisor  $D_T$  of degree  $d$  on  $G$ .
- (iii) Algorithms 2 and 3 are inverse to one another:  $T_{D_T} = T$  and  $D_{T_D} = D$ .

Moreover, under the bijection furnished by Algorithms 2 and 3:

- (iv) The set  $R$  is the same at the end of both algorithms.
- (v) The externally active edges<sup>6</sup> for  $T$  are precisely the elements of  $\text{ExtAct} = E \setminus R$ , and the externally passive edges for  $T$  are precisely the elements of  $\text{ExtPass} = R \setminus T$ .
- (vi) The degree of  $\sum_{v \neq q} a_v(v)$  is equal to  $g - \text{ex}(T)$ , where  $g = m - n + 1$ . Equivalently,  $a_q = d - g + \text{ex}(T)$ .

*Remark 5.2.* (1) The complexity of both Algorithms 2 and 3 is the same as that of Dhar’s algorithm (Algorithm 1), namely  $O(n^2)$ .

---

<sup>6</sup>An edge  $e \in E \setminus T$  is called *externally active* for  $T$  if it is the largest element in the unique cycle contained in  $T \cup \{e\}$ , and is called *externally passive* for  $T$  if it is not externally active. The *external activity* of  $T$  is the number of externally active edges for  $T$  and is denoted by  $\text{ex}(T)$ .

**Input:**  
 $G = (V, E)$  is graph with a fixed ordering on  $E$ ,  
 $q \in V(G)$ ,  
 $T$  a spanning tree of  $G$ .  
**Output:**  
 $D_T = \sum_v a_v(v)$ , a  $q$ -reduced divisor of degree  $d$ .  
**Initialization:**  
 $X = \{q\}$  (“burnt” vertices),  
 $R = \emptyset$  (“burnt” edges).  
**while**  $X \neq V(G)$  **do**  
     $f = \min\{e = \{s, t\} \in E(G) \mid e \notin R, s \in X, t \notin X\}$ ,  
    **if**  $f \in T$  **then**  
        let  $v \in V(G) \setminus X$  be the vertex incident to  $f$ ,  
         $a_v := |\{e \text{ incident to } v \mid e \in R\}|$ ,  
         $X \leftarrow X \cup \{v\}$   
    **end**  
     $R \leftarrow R \cup \{f\}$   
**end**  
 $a_q := d - \sum_{v \neq q} a_v$ ,  
 $\text{ExtAct} = E \setminus R$ ,  
 $\text{ExtPass} = R \setminus T$ ,  
Output  $D_T = \sum_v a_v(v)$ .

**Algorithm 3:** Spanning tree to reduced divisor

- (2) A natural choice for the fixed degree is  $d = g$ , in which case it follows from Theorem 5.1 (vi) that  $a_q = ex(T)$ .

*Remark 5.3.* The problem of giving an explicit bijection between reduced divisors (in the guise of  $G$ -parking functions) and spanning trees has been studied in several previous works (see, e.g., [26, 20, 11]). There are also a number of bijections in the literature between  $q$ -critical configurations and spanning trees (see [29, 16, 15, 27, 48, 44]). For a fixed vertex  $q$ ,  $q$ -critical configurations provide another set of representatives for equivalence classes of divisors (see, e.g., [13, 14]). There is a simple relationship between reduced and critical divisors:  $D$  is  $q$ -reduced if and only if  $K^+ - D$  is  $q$ -critical, where  $K^+ = \sum_{v \in V(G)} (\deg(v) - 1)(v)$  [7].

**5.3. Computing the reduced divisor.** Recall that computing the  $q$ -reduced divisor equivalent to some divisor  $D$  can be viewed as the solution to a linear (Theorem 4.14) or quadratic (Theorem 4.12) integer programming problem. Dhar’s algorithm (which runs in time  $O(n^2)$ ) shows that one can efficiently *check* whether a given divisor is the solution to the corresponding integer programming problem. Next we show that in fact one can *find* the solution efficiently as well.

Fix a base vertex  $q \in V(G)$ . Given a divisor  $D \in \text{Div}(G)$ , Algorithm 4 below efficiently<sup>7</sup> computes the  $q$ -reduced divisor  $D' \sim D$ . The idea behind the algorithm is as follows. Starting with a divisor  $D$ , the first step is to replace  $D$  with an equivalent divisor whose coefficients are “small”. This is accomplished by the simple trick of replacing  $[D]$  with  $[D] - Q \lfloor L_{(q)}[D] \rfloor$ , where  $L_q$  is as in Construction 3.1 and  $\lfloor \cdot \rfloor$  denotes the coordinate-wise floor function. The second step is to make the divisor effective outside  $q$ . This is done by having negative vertices borrow from their neighbors in a controlled way. The third step is to iterate Dhar’s algorithm until we reach a  $q$ -reduced divisor. More specifically, if  $D$  is not yet reduced then by running Dhar’s algorithm on  $D$  we obtain a subset  $A$  of  $V(G) \setminus \{q\}$  such that firing all vertices in  $A$  once yields a divisor  $D - \Delta(\chi_A)$  which is still effective outside  $q$ . Replacing  $D$  by  $D - \Delta(\chi_A)$  and iterating this procedure, one obtains (after finitely many iterations) a  $q$ -reduced divisor. Moreover, the number of iterations can be explicitly bounded in terms of the  $j$ -function (§3.2) using formula (4.9).

A formal statement of the resulting algorithm appears below (Algorithm 4).

**Input:**

$Q$  is the Laplacian matrix of the graph  $G$ ,

$D \in \text{Div}(G)$ ,

$q \in V(G)$ .

**Output:**

$D' \sim D$  the unique  $q$ -reduced divisor equivalent to  $D$ .

(Step 1)

Find the generalized inverse  $L_{(q)}$  of  $Q$ , as in Construction 3.1. Compute the divisor  $[D'] = [D] - Q \lfloor L_{(q)}[D] \rfloor$ .

(Step 2)

**while** there exists  $v \neq q$  with  $D'(v) < 0$  **do**  $[D'] \leftarrow [D'] + Q[\chi_{\{v\}}]$ .

(Step 3)

Let  $A_0 = V(G)$ ,  $v_0 = q$ , and  $i = 1$ .

**while**  $i \leq n - 1$  **do**

    Let  $A_i = A_{i-1} \setminus \{v_{i-1}\}$ .

**if** there exists  $v_i \in A_i$  such that  $D'(v_i) < \text{outdeg}_{A_i}(v_i)$  **then**  $i \leftarrow i + 1$ .

**else**  $[D'] \leftarrow [D'] - Q[\chi_{A_i}]$ . Reset  $i = 1$ .

**end**

**Algorithm 4:** Finding the Reduced Divisor

**Correctness of Algorithm 4:**

Assume for the moment that the algorithm actually terminates and produces an output. It is easy to see that the output is linearly equivalent to  $D$ . Also, the output

<sup>7</sup>“Efficient” in this context means that the running time will be polynomial in  $m$  and  $n$  with only  $\log(\deg(D))$ -bit computations involved.

passes Dhar's algorithm and therefore is  $q$ -reduced (in fact, as discussed above, one can view Step 3 as an iterated Dhar's algorithm). Therefore, for the correctness of the algorithm, we only need to show that it terminates (which follows *a posteriori* from the efficiency analysis below).

#### Efficiency of Algorithm 4:

**Proposition 5.4.** *If  $[D'] = [D] - Q[L_{(q)}[D]]$ , then  $|D'(v)| < \deg(v)$  for all  $v \neq q$ .*

*Proof.* Recall from Construction 3.1 that  $QL_{(q)} = I + R_{(q)}$ , where  $I$  is the identity matrix and  $R_{(q)}$  has  $-1$  entries in  $q^{\text{th}}$  row and is zero elsewhere. Therefore  $[D] = QL_{(q)}[D] + \deg(D) \cdot \mathbf{e}_q$ , where  $\mathbf{e}_q$  is the column vector which is 1 in position  $q$  and zero elsewhere. Now

$$\begin{aligned} [D'] &= [D] - Q[L_{(q)}[D]] \\ &= Q(L_{(q)}[D] - [L_{(q)}[D]]) + \deg(D) \cdot \mathbf{e}_q \\ &= Q\mathbf{f} + \deg(D) \cdot \mathbf{e}_q, \end{aligned}$$

where  $\mathbf{f} = L_{(q)}[D] - [L_{(q)}[D]]$  is a vector with entries in  $[0, 1)$ . It is now easy to show that the absolute values of the entries of  $Q\mathbf{f}$  are bounded by the degree of the corresponding vertices.  $\square$

*Remark 5.5.* Computing the generalized inverse  $L_{(q)}$  takes time at most  $O(n^\omega)$ , where  $\omega$  is the exponent for matrix multiplication (currently  $\omega = 2.376$  [25]). Notice that this computation is done only once. The second computation in Step (1) can be done using  $O(n^2)$  operations (multiplication and addition). For bit complexity, one can check that the denominators appearing in the generalized inverse  $L_{(q)}$  are annihilated by the exponent of the Jacobian group. The exponent is bounded above by the number of spanning trees of the graph. If we allow at most  $c$  parallel edges then there are at most  $c^{n-1} \cdot n^{n-2}$  spanning trees. Moreover, one can also show that the absolute value of the entries of  $L_{(q)}$  are bounded above by  $R_{\max}$ , the maximum effective resistance between any two vertices of the graph. Therefore all integers in the algorithm can be represented in  $O(n \cdot \log cn)$  bits.

Now we will use our potential theoretic techniques to bound the number of chip-firing moves in Algorithm 4. As we will see, several different bounds in the literature can be obtained as special cases or corollaries of our general potential theory bound.

**Proposition 5.6.** (a) *Let  $D_1$  be the output of Step 1 of Algorithm 4. Then Step 2 of Algorithm 4 terminates in at most  $b_q(K^+ - D_1)$  borrowing moves, where  $K^+ = \sum_{v \in V(G)} (\deg(v) - 1)(v)$ .*

(b) *Let  $D_2$  be the output of Step 2 of Algorithm 4. Then Step 3 of Algorithm 4 terminates in at most  $b_q(D_2)$  firing moves.*

(c) *Algorithm 4 terminates in fewer than*

$$(5.7) \quad 3 \sum_v \sum_p j_q(p, v) \deg(v)$$

*chip-firing moves.*



*Proof.* (a) By Proposition 5.4,  $|D_1(v)| < \deg(v)$  for all  $v \neq q$ . So for any  $v \neq q$  with  $D_1(v) < 0$ , only one borrowing is needed to make the vertex positive. Moreover, the resulting positive number will be less than  $\deg(v)$ . This fact, together with Proposition 5.4, guarantees that the output of Step 2 satisfies  $0 \leq D_1(v) < \deg(v)$  for all  $v \neq q$ . The result now follows from Lemma 4.8 and Proposition 4.5 (b); the value of  $b_q(\cdot)$  is at least  $\sum_v \sum_p j_q(p, v) D_1(v)$  on the input of Step 2, and is at most  $\sum_v \sum_p j_q(p, v) (\deg(v) - 1)$ . Moreover, with each borrowing  $b_q(\cdot)$  increases by 1.

(b) This again follows from Lemma 4.8 and Proposition 4.5 (b). Note that  $D_2(v) < \deg(v)$ , and that no vertex  $v \neq q$  can become negative in Step 3.

(c) This follows from parts (a) and (b) and the inequalities

$$b_q(K^+ - D_1) < 2 \sum_v \sum_p j_q(p, v) \deg(v) ,$$

$$b_q(D_2) < \sum_v \sum_p j_q(p, v) \deg(v) .$$

□

**5.4. Comparison to other techniques in the literature.** By basic properties of the  $j$ -function (see § 3.2), we have

$$3 \sum_v \sum_p j_q(p, v) \deg(v) \leq 3(n-1) \sum_v r(v, q) \deg(v) .$$

By Proposition 5.6 (specifically (5.7)), it follows that Algorithm 4 terminates in fewer than

$$(5.8) \quad 3(n-1) \sum_v r(v, q) \deg(v)$$

chip-firing moves.

The bound (5.8) can be computed in matrix multiplication time  $O(n^\omega)$  (currently  $\omega = 2.376$ ) because  $j_q(p, v)$  is simply the  $(p, v)$ -entry of the matrix  $L_{(q)}$  in Construction 3.1.

*Remark 5.9.* There are several ways to bound the expression in (5.8) in terms of more familiar invariants of the graph. For example:

- (1) Let  $R_{\max}$  be the maximum effective resistance between vertices of  $G$  and let  $\Delta_{\max}$  be the maximum degree of a vertex in  $G$ . Then (5.8) is bounded above by

$$3(n-1) R_{\max} \sum_{v \neq q} \deg(v) ,$$

which is, in turn, bounded above by

$$3(n-1)^2 R_{\max} \Delta_{\max} .$$

These estimates give a factor  $n$  improvement over the bound for the running time of Algorithm 4 which could be derived using the technique in [34] by Holroyd, Levine, Mészáros, Peres, Propp, and Wilson.

- (2) Using Foster's network theorem, one can show that

$$r(v, q) < 3 \sum_{v \in V(G)} (\deg(v) + 1)^{-1}$$

(see, e.g., [24, proof of Theorem 6]). So another upper bound for the running time of Algorithm 4 is

$$9(n-1) \sum_{v \in V(G)} (\deg(v) + 1)^{-1} \sum_{v \neq q} \deg(v) .$$

This is a good bound when the graph is close (on average) to being regular. If one uses the fact that degree of a vertex is at least the edge-connectivity  $\lambda$  of the graph, one gets a bound of the form  $O(n^2 m / \lambda)$  for the running time of Algorithm 4. This is (up to constant factors) the bound that one can derive from the techniques of van den Heuvel [32].

- (3) Let  $\lambda_1$  (called the *algebraic connectivity* of  $G$ ) be the smallest non-zero eigenvalue of  $Q$ . Then  $r(p, q) \leq \frac{2}{\lambda_1}$  for every  $p, q \in V(G)$ . Indeed, the proof of the Cauchy-Schwarz inequality shows that for any positive semidefinite matrix  $L$  with largest eigenvalue  $\eta$ , and for all vectors  $x$  and  $y$ ,  $|x^T L y| \leq \eta \|x\|_2 \|y\|_2$ ; if we apply this estimate with  $x = y = (p) - (q)$ ,  $L = Q^+$  (cf. Construction 3.2), then  $\eta = 1/\lambda_1$ , and we obtain

$$r(p, q) = \langle (p) - (q), (p) - (q) \rangle \leq \frac{2}{\lambda_1} .$$

Therefore an upper bound for the running time of Algorithm 4 is

$$\frac{6(n-1)}{\lambda_1} \sum_{v \neq q} \deg(v) .$$

This is the bound that one can derive from the techniques of Björner–Lovász–Shor [17] or Chung–Ellis [22].

- (4) By Rayleigh's monotonicity law, we have  $R_{\max} \leq \text{diam}(G)$ , where  $\text{diam}(G)$  denotes the *diameter* of  $G$ . Equality holds if and only if  $G$  is a path. In fact  $R_{\max}$  is much smaller than  $\text{diam}(G)$  in a general graph. Another upper bound for the running time of Algorithm 4 is

$$3(n-1) \text{diam}(G) \sum_{v \neq q} \deg(v) .$$

This is the bound that one can derive from the techniques of Tardos [53].

*Remark 5.10.* Items (3) and (4) in the previous remark clarify the relationship between the seemingly different approaches of Tardos and Björner–Lovász–Shor.

## 6. SOME APPLICATIONS OF THE ALGORITHMS

**6.1. Bijective matrix-tree theorem.** Kirchhoff's celebrated *matrix-tree theorem* is usually formulated as follows. Let  $G$  be a (connected) graph. Following the terminology from Construction 3.1, fix an integer  $1 \leq i \leq n$  and let  $Q_i$  be the invertible  $(n-1) \times$

$(n - 1)$  matrix obtained from the Laplacian matrix  $Q$  of  $G$  by deleting  $i^{\text{th}}$  row and  $i^{\text{th}}$  column from  $Q$ .

**Theorem 6.1** (Kirchhoff’s matrix-tree theorem [37]). *The number of spanning trees in  $G$  is equal to  $|\det(Q_i)|$ .*

Our aim in this section is to give an “efficient bijective” proof of Theorem 6.1. In order to make sense of this goal, it is useful to reformulate Kirchhoff’s theorem in a more natural way in terms of the Jacobian group<sup>8</sup>  $\text{Jac}(G) = \text{Div}^0(G)/\text{Prin}(G)$ , where  $\text{Div}^0(G)$  is the subgroup of  $\text{Div}(G)$  consisting of divisors of degree zero and  $\text{Prin}(G)$  (the group of principal divisors) is the image of the Laplacian operator  $\Delta : \mathcal{M}(G) \rightarrow \text{Div}(G)$ .

By elementary group theory (e.g. the theory of the *Smith normal form*), one sees that  $\text{Jac}(G)$  is the torsion part<sup>9</sup> of the cokernel of  $Q : \mathbb{Z}^n \rightarrow \mathbb{Z}^n$ , and the order of  $\text{Jac}(G)$  is equal to  $|\det(Q_i)|$ . We may thus reformulate Kirchhoff’s theorem as follows:

**Theorem 6.2** (Kirchhoff’s matrix-tree theorem, canonical formulation). *The number of spanning trees in  $G$  is equal to  $|\text{Jac}(G)|$ . Moreover, there exists an efficiently computable bijection between elements of  $\text{Jac}(G)$  and spanning trees of  $G$ .*

Note that such a bijection cannot be canonical, as that would imply the existence of a distinguished spanning tree in  $G$  corresponding to the identity element of  $\text{Jac}(G)$ , but it is clear (think of the case where  $G$  is an  $n$ -cycle) that there is in general no distinguished spanning tree. Therefore, one needs to make some choices to write down a bijection.

*Proof of Theorem 6.2.* By Corollary 4.13, if we fix a vertex  $q$  of  $G$ , there is a unique  $q$ -reduced divisor representing each class in  $\text{Div}^0(G)$ . In particular, there is an explicit bijection between  $\text{Div}^0(G)$  and the set of  $q$ -reduced divisors of degree 0. If in addition we choose a total order on  $E(G)$ , then the algorithms in §5.2 show that there is a bijection between  $q$ -reduced divisors of degree 0 and spanning trees of  $G$ .

We have shown in §5.3 that the unique  $q$ -reduced representative for each class in  $\text{Div}^0(G)/\text{Prin}(G)$  can be computed efficiently. And in §5.2 we proved that the bijection between  $q$ -reduced divisors of degree 0 and spanning trees of  $G$  is also efficient.  $\square$

Kirchhoff’s matrix-tree theorem is of course a classical result. The main new contribution here is to observe that the bijections between reduced divisors and spanning trees (as described in §5.2 and Remark 5.3), in conjunction with Corollary 4.13, furnish a simple bijective proof of Kirchhoff’s theorem, and moreover this bijection is efficiently computable.

**6.2. Random spanning trees.** The *random spanning tree* problem has been extensively studied in the literature and there are two known types of algorithms: determinant based algorithms (e.g. [31, 23, 38]) and random walk based algorithms (e.g. [19, 1, 55, 36] and [43, Chapter 4]).

<sup>8</sup>Although we do not need this here, it is worth mentioning that the energy pairing descends to a non-degenerate  $\mathbb{Q}/\mathbb{Z}$ -valued bilinear form on  $\text{Jac}(G)$ ; it is called the “monodromy pairing” in [51].

<sup>9</sup>The full cokernel  $\text{Pic}(G) = \text{Div}(G)/\text{Prin}(G)$  is isomorphic to  $\mathbb{Z} \oplus \text{Jac}(G)$ .

Here we give a new deterministic polynomial time algorithm for choosing a random spanning tree in a graph  $G$ .

Although the bound we obtain for the running time of our algorithm does not beat the current best known running time  $O(n^\omega)$  of [23], we believe that our algebraic method has some advantages. For example, it is trivial that the output of our algorithm is a uniformly random spanning tree, whereas in other algorithms (e.g. [23]) this fact is non-trivial and requires proof. See Remark 6.3 for another advantage.

The idea behind our algorithm is very simple. Fix a vertex  $q \in V(G)$  and a total ordering of  $E(G)$ . The first step in the algorithm is to compute a presentation of  $\text{Jac}(G)$  as a direct sum of cyclic groups; this can be done efficiently by computing the Smith normal form for  $Q$ . Once  $\text{Jac}(G)$  is presented in this way, it is clear how to select a random element. Having done so, one computes the corresponding  $q$ -reduced divisor and then the corresponding spanning tree.

This procedure is formalized in Algorithm 5 below.

**Input:** A graph  $G$ .

**Output:** A uniformly random spanning tree of  $G$ .

- (1) Fix a vertex  $q \in V(G)$  and a total ordering of  $E(G)$ .
- (2) Compute the Smith normal form of the Laplacian matrix of  $G$  to find:
  - invariant factors  $\{n_1, \dots, n_s\}$ ,
  - generators  $\{\mathbf{g}_1, \dots, \mathbf{g}_s\}$  for  $\text{Jac}(G)$  (thought of as elements of  $\text{Div}^0(G)$ ).
- (3) Choose a random integer  $0 \leq a_i \leq n_i - 1$  for  $(1 \leq i \leq s)$ .
- (4) Compute the divisor  $D = \sum_{i=1}^s a_i \mathbf{g}_i$ .
- (5) Use Algorithm 4 to find the unique  $q$ -reduced divisor  $D'$  equivalent to  $D$ .
- (6) Use Algorithm 2 to find the spanning tree corresponding to  $D'$ .

**Algorithm 5:** Choosing a uniformly random spanning tree

To our knowledge, the fastest known Smith normal form algorithm (Step (2)) is given in [35] and has running time  $(n^{2.697263} \log \|Q\|)^{1+o(1)}$ , where  $\|Q\|$ , for our application, means the maximal degree of a vertex  $\Delta_{\max}$ . See also [50] for a fast and practical Smith normal form algorithm. For the running time of Step (5) see (5.7), (5.8), and Remark 5.9. Step (6) can be done in  $O(n^2)$  steps.

*Remark 6.3.* Note that for repeated sampling of random spanning trees in  $G$ , one has to perform steps (1) and (2) of Algorithm 5 only once. Note also that if there are  $N$  spanning trees in  $G$ , our algorithm uses only  $\log_2 N$  random bits for generating each random spanning tree. Thus our algorithm may have some advantages over existing methods for sampling multiple spanning trees. For example, very few random bits are required in our algorithm to generate pairwise independent spanning trees; to generate  $k$  pairwise independent spanning trees, the naive approach would use  $k \cdot \log_2 N$  random bits. But one can use standard methods to pick pairwise independent elements of the group using only  $O(\log_2 N)$  random bits. Also, it is possible with our method to sample multiple spanning trees according to joint distributions other than the uniform distribution. (We thank Richard Lipton for these observations; see [40]).

**6.3. Other applications.** We list briefly some other applications of our algorithms:

- (1) (The group law attached to chip-firing games) If we fix a vertex  $q \in V(G)$ , then  $\text{Jac}(G)$  induces a group structure on the set of  $q$ -reduced divisors ( $G$ -parking functions) or  $q$ -critical divisors of  $G$ . The latter is called *critical group* (or sandpile group) of  $G$ . Adding two elements in one of these groups requires first adding the given divisors as elements of  $\text{Div}(G)$ , and then finding the unique  $q$ -reduced or  $q$ -critical divisor equivalent to the sum. Our algorithm for finding  $q$ -reduced divisors can be used to efficiently compute the group law in these groups. A different approach for performing the group operation is given in [32] using “oil games”. The problem of finding a “purely algebraic” method for computing the sum of two elements of the critical group (and analyzing the running time of the resulting algorithm) was posed as an open problem by Chung and Ellis in [22].
- (2) (Determining whether the dollar game is winnable) In [7], the authors consider a dollar game played on the vertices of  $G$ . Given a divisor  $D$ , thought of as a configuration of dollars on  $G$ , the goal of the game is to get all the vertices out of debt via borrowing and lending moves, i.e., to find an effective divisor  $D'$  linearly equivalent to  $D$ . By the proof of Theorem 3.3 in [7], the game is winnable iff the unique  $q$ -reduced divisor equivalent to  $D$  is effective. As a corollary, once we can efficiently compute the  $q$ -reduced divisor associated to a given configuration, we can efficiently decide whether or not there is a winning strategy, and when there is one we can efficiently compute a sequence of winning moves.

These considerations are related to the Riemann-Roch theorem for graphs from [7]. To any divisor  $D \in \text{Div}(G)$  one associates an integer  $r(D) \geq -1$ , called the *rank* of  $D$ , such that  $r(D) \geq 0$  iff the unique  $q$ -reduced divisor equivalent to  $D$  is effective. Our algorithm for finding  $q$ -reduced divisors can therefore be used to efficiently check whether or not  $r(D) \geq 0$ . More generally, we can efficiently check whether  $r(D) \geq c$  for any fixed constant  $c$ . It is an open problem to determine whether or not one can compute  $r(D)$  itself in polynomial time. For a study of this problem, see [45].

## APPENDIX A. METRIC GRAPHS

Our goal in this section is to extend some of the considerations from §3.3, as well as Theorem 4.12 and Corollary 4.13, to the setting of *metric graphs*. The metric graph analogue of Corollary 4.13 has already been proved in [46] and [33], but since this is the main ingredient needed to prove the Riemann-Roch theorem for tropical curves (an important result in tropical geometry), it seems worthwhile to present the new proof which follows. Our main new result is Theorem A.7 (the metric graph analogue of Theorem 4.12).

**A.1. Background on metric graphs and potential theory.** We assume that the reader is familiar with some basic facts concerning potential theory on metric graphs; see for example [6, 8]. We recall here the main facts and terminology which we will use.

Let  $\Gamma$  be a metric graph. Let  $\mathcal{C}(\Gamma)$  denote the  $\mathbb{R}$ -algebra of continuous real-valued functions on  $\Gamma$ , let  $\text{CPA}(\Gamma) \subset \mathcal{C}(\Gamma)$  be the vector space consisting of all continuous piecewise affine functions on  $\Gamma$ , and let  $R(\Gamma)$  be the subgroup of  $\text{CPA}(\Gamma)$  consisting of continuous piecewise affine functions with integer slopes (this can be viewed as the space of *tropical rational functions* on  $\Gamma$ , cf. [30, 46]).

Let  $\Delta$  be the Laplacian operator on  $\Gamma$  (see [6, 8]), which takes a certain subspace  $\text{BDV}(\Gamma)$  of  $\mathcal{C}(\Gamma)$  into the space of measures of total mass zero on  $\Gamma$ . (The abbreviation  $\text{BDV}$  stands for “bounded differential variation”.) More precisely, let  $\text{Meas}^0(\Gamma)$  be the vector space of finite signed Borel measures of total mass zero on  $\Gamma$  and let  $\mathbb{R} \subset \mathcal{C}(\Gamma)$  denote the space of constant functions on  $\Gamma$ . Then the space  $\text{BDV}(\Gamma)$  is characterized by the property that  $f \mapsto \Delta(f)$  induces an isomorphism of vector spaces

$$\text{BDV}(\Gamma)/\mathbb{R} \longleftrightarrow \text{Meas}^0(\Gamma) .$$

For a fixed  $q \in \Gamma$ , an inverse to  $\Delta$  on  $\{f \in \text{BDV}(\Gamma) \mid f(q) = 0\}$  is given by

$$(A.1) \quad \nu \mapsto \int_{\Gamma} j_q(x, y) d\nu(y) \in \text{BDV}(\Gamma)$$

where  $j_q(x, y)$  is the fundamental potential kernel on  $\Gamma$  (defined below, see also [8]). It is shown in [8] that  $\text{CPA}(\Gamma) \subset \text{BDV}(\Gamma)$  and that  $\Delta(f)$  is a discrete measure if and only if  $f \in \text{CPA}(\Gamma)$ .

For  $f \in \text{CPA}(\Gamma)$  we have

$$\Delta(f) = \sum_{p \in \Gamma} \sigma_p(f)$$

where  $-\sigma_p(f)$  is the sum of the slopes of  $f$  in all tangent directions emanating from  $p$ . (Note that  $\Delta(f) = -\text{div}(f)$  with the conventions from [30, 33].) This formula uniquely characterizes  $\Delta$  on  $\text{BDV}(\Gamma)$ , because  $\text{CPA}(\Gamma)$  is dense in  $\mathcal{C}(\Gamma)$  and for  $f \in \text{CPA}(\Gamma)$  and  $g \in \text{BDV}(\Gamma)$ , we have

$$(A.2) \quad \int_{\Gamma} f \Delta(g) = \int_{\Gamma} g \Delta(f) .$$

(In fact, (A.2) holds for all  $f, g \in \text{BDV}(\Gamma)$ .)

For fixed  $q$  and  $y$ , one can define  $j_q(x, y)$  as the unique element of  $\text{CPA}(\Gamma)$  such that  $\Delta_x j_q(x, y) = \delta_y(x) - \delta_q(x)$ . In terms of electrical network theory, if we think of  $\Gamma$  as an electrical network with resistances given by the edge lengths, then  $j_q(x, y)$  is the potential at  $x$  when one unit of current enters the network at  $y$  and exits at  $q$ , with reference voltage zero at  $q$ . It is a basic fact, proved in [21] (see [6] for an alternate proof), that  $j_q(x, y)$  is jointly continuous in  $x$ ,  $y$ , and  $q$ .

There is a canonical bilinear form, called the *energy pairing*, on  $\text{Meas}^0(\Gamma)$ . It can be defined in many equivalent ways; if  $\mu = \Delta(f)$  and  $\nu = \Delta(g)$  are in  $\text{Meas}^0(\Gamma)$ , then

$$(A.3) \quad \langle \mu, \nu \rangle = \int f d\nu = \int g d\mu = \int f'(x) g'(x) dx = \int j_q(x, y) d\mu(x) d\nu(y) .$$

It is proved in [8, Theorem 10.4] that the energy pairing is *positive definite*. If  $\mu \in \text{Meas}^0(\Gamma)$  is a discrete measure, the energy pairing  $\langle \mu, \mu \rangle$  has a nice interpretation in terms of electrical networks. Write  $\mu = \sum a_i \delta_{p_i} - \sum b_j \delta_{q_j}$  with  $a_i, b_j \geq 0$ . If  $a_i$  units of current enter the network at each  $p_i$  and  $b_j$  units of current exit the network at each  $q_j$ , then  $\langle \mu, \mu \rangle$  is the total energy (or power) dissipated in the network. The different formulas for the energy pairing in (A.3) correspond to the different classical formulas for computing the power (as the sum of  $VI$ ,  $I^2R$ , etc.).

**A.2. Existence and uniqueness of reduced divisors.** Let  $\text{Div}(\Gamma)$  (the group of *divisors* on  $\Gamma$ ) be the free abelian group on  $\Gamma$ . We can identify a divisor  $D = \sum a_i(p_i)$  on  $\Gamma$  with the discrete measure  $\mu_D := \sum a_i \delta_{p_i}$ . We will frequently identify divisors and measures and will often not explicitly differentiate between  $D$  and  $\mu_D$ .

Let  $\text{Div}^0(\Gamma)$  be the subgroup of divisors of degree zero on  $\Gamma$ , and let  $\text{Prin}(\Gamma)$  be the subgroup  $\{\Delta(f) \mid f \in R(\Gamma)\}$  of  $\text{Div}^0(\Gamma)$  consisting of *principal divisors*. We write  $D \sim D'$  if  $D - D'$  belongs to  $\text{Prin}(\Gamma)$  and say that  $D$  and  $D'$  are *linearly equivalent*.

Fix  $q \in \Gamma$ . A divisor  $D = \sum a_p(p)$  is called *effective* if  $a_p \geq 0$  for all  $p$ , and is called *effective outside  $q$*  if  $a_p \geq 0$  for all  $p \neq q$ . For  $D \in \text{Div}(\Gamma)$ , we define the complete linear system  $|D|$  to be the set of all effective divisors  $E$  equivalent to  $D$ , i.e.,  $|D| = \{E \in \text{Div}(\Gamma) \mid E \geq 0, E \sim D\}$ . Similarly, we define  $|D|_q$  to be the set of divisors equivalent to  $D$  which are effective outside  $q$ :

$$|D|_q = \{E \in \text{Div}(\Gamma) \mid E(p) \geq 0, \forall p \neq q, E \sim D\}.$$

**Definition.** Fix  $q \in \Gamma$ . A divisor  $D$  on  $\Gamma$  is called  *$q$ -reduced* if it satisfies the following two conditions:

- (R1)  $D$  is effective outside  $q$ .
- (R2) If  $f \in R(\Gamma)$  is non-constant and has a global maximum at  $q$ , then

$$D + \Delta(f) \notin |D|_q.$$

Note that this is not the usual definition of reduced divisor on a metric graph, but it is easily seen to be equivalent to the definition used in [3, 33, 42]:

**Lemma A.4.** *A divisor  $D$  is  $q$ -reduced if and only if it satisfies (R1) and*

*(R2') For every closed connected set  $X \subseteq \Gamma$  not containing  $q$ , there exists a point  $p \in \partial X$  such that  $D(p) < \text{outdeg}_X(p)$ .*

*Proof.* Suppose  $D$  is effective outside  $q$  and satisfies (R2). Given a closed connected set  $X$  not containing  $q$ , construct a rational function  $f \in R(\Gamma)$  which is 0 on  $X$  and  $\epsilon$  outside of an  $\epsilon$ -neighborhood of  $X$  for some sufficiently small  $\epsilon > 0$ , with slope 1 in each outgoing direction from  $X$  and  $f(q) = \epsilon$  (so  $f$  achieves its maximum at  $q$ ). By (R2), there is a point  $p \in \Gamma \setminus \{q\}$  such that  $(D + \Delta(f))(p) < 0$ . Since  $(\Delta(f))(p) = -\text{outdeg}_X(p)$ , it follows that  $p \in \partial X$  and  $D(p) < \text{outdeg}_X(p)$ .

Conversely, suppose  $D$  is effective outside  $q$  and satisfies (R2'). Given a non-constant function  $f \in R(\Gamma)$  achieving its maximum value at  $q$ , let  $X$  be a connected component of the set of points where  $f$  achieves its minimum. By assumption (R2'), there exists  $p \in \partial X$  such that  $D(p) < \text{outdeg}_X(p)$ . Since  $(\Delta(f))(p) < -\text{outdeg}_X(p)$ , we have  $(D + \Delta(f))(p) \leq D(p) - \text{outdeg}_X(p) < 0$  and thus  $D + \Delta(f) \notin |D|_q$ .  $\square$

From the definition, one sees that if  $D$  is a  $q$ -reduced divisor and  $D + \Delta(f)$  is effective outside  $q$  for some non-constant function  $f \in R(\Gamma)$ , then  $f$  cannot have a global maximum at  $q$ . It turns out that such an  $f$  must in fact have a global *minimum* at  $q$ . The following lemma is the analogue of Lemma 4.11.

**Lemma A.5** (Principle of least action). *If  $D$  is  $q$ -reduced and  $f \in R(\Gamma)$  is a rational function with  $D + \Delta(f) \in |D|_q$ , then  $f$  has a global minimum at  $q$ .*

*Proof.* Suppose not, and let  $X$  be the set of points where  $f$  achieves its minimum value. Then  $X$  is a closed connected set not containing  $q$ . By Lemma A.4, there exists  $p \in \partial X$  such that  $D(p) < \text{outdeg}_X(p)$ . On the other hand, we have  $\Delta(f)(p) < -\text{outdeg}_X(p)$ , and thus  $(D + \Delta(f))(p) < 0$ , contradicting the assumption that  $D + \Delta(f) \in |D|_q$ .  $\square$

The importance of reduced divisors is given by the following theorem [33, 46], which is analogous to the corresponding result for finite graphs (Corollary 4.13 above) proved in [7]:

**Theorem A.6.** *Fix  $q \in \Gamma$ . There is a unique  $q$ -reduced divisor in each linear equivalence class of divisors on  $\Gamma$ .*

We will give a new proof of this theorem based on the following energy minimization result. For a divisor  $D \in \text{Div}(\Gamma)$  of degree  $d$ , define the  $q$ -energy of  $D$  by

$$\mathcal{E}_q(D) = \langle D - d(q), D - d(q) \rangle .$$

If  $D = \sum a_i(p_i)$  is effective and  $p_i \neq q$  for all  $i$ , then  $\mathcal{E}_q(D)$  is the total energy dissipated in an electrical network in which  $a_i$  units of current enter the network at  $p_i$  and  $d = \sum a_i$  units of current exit at  $q$ .

**Theorem A.7.** *Let  $D$  be a divisor on  $\Gamma$  which is effective outside  $q$ . Then  $D$  is  $q$ -reduced if and only if it has smaller  $q$ -energy than every other divisor in  $|D|_q$ .*

*Proof.* Suppose first that  $D$  has minimal  $q$ -energy among all divisors in  $|D|_q$ . We want to show that  $D$  is  $q$ -reduced. Let  $f \in R(\Gamma)$  be a non-constant rational function with a global maximum at  $q$ . Without loss of generality, we may assume that  $f(q) = 0$ . We claim that  $D' := D + \Delta(f)$  does not belong to  $|D|_q$ . Indeed, if  $D$  were in  $|D|_q$ , then we would have

$$\begin{aligned} \mathcal{E}_q(D) &= \mathcal{E}_q(D') - 2\langle D' - d(q), \Delta(f) \rangle + \mathcal{E}_q(\Delta(f)) \\ &> \mathcal{E}_q(D') - 2f(D') \\ &\geq \mathcal{E}_q(D') \end{aligned}$$

since  $f(D') \leq 0$  by our assumptions on  $f$  and  $D'$ . This contradicts the minimality of  $\mathcal{E}_q(D)$ , and thus  $D' \notin |D|_q$  as claimed.

For the other direction, we need to prove that if  $D$  is  $q$ -reduced and  $D' = D + \Delta(f) \in |D|_q$  with  $f \in R(\Gamma)$  non-constant, then  $\mathcal{E}_q(D') > \mathcal{E}_q(D)$ . As before we may assume that



$f(q) = 0$ . It follows from Lemma A.5 that  $f(p) \geq 0$  for all  $p \in \Gamma$ . We thus have

$$\begin{aligned}\mathcal{E}_q(D') &= \mathcal{E}_q(D) + 2\langle D' - d(q), \Delta(f) \rangle + \mathcal{E}_q(\Delta(f)) \\ &> \mathcal{E}_q(D) + 2f(D) \\ &\geq \mathcal{E}_q(D)\end{aligned}$$

as desired.  $\square$

**Corollary A.8.** *For every  $D \in \text{Div}(\Gamma)$ , there is at most one  $q$ -reduced divisor equivalent to  $D$ .*

In order to deduce Theorem A.6 from Theorem A.7, we need some auxiliary results.

**Proposition A.9.** *Every divisor  $D$  on  $\Gamma$  is equivalent to a divisor which is effective outside  $q$ .*

*Proof.* This is proved in [33] and [46]. Here is a simple variant of the proof in [33] (along the lines of a similar argument in [7]). Let  $S$  be the set of branch points of  $\Gamma$  together with  $q$  and all points in the support of  $D$ . If  $p \in \Gamma \setminus \{q\}$ , define  $\lambda_S(p)$  (the “level of  $p$ ” relative to  $S$ ) to be the minimal number of elements of  $S \setminus \{q\}$  contained in a path from  $q$  to  $p$ . We define  $\lambda_S(q)$  to be  $-1$ . Let  $k$  be the maximum value of  $\lambda_S(p)$  for  $p \in \Gamma$ . For  $i = 0, \dots, k$  let  $S_i = \{p \in \Gamma \mid \lambda_S(p) = i\}$ . For  $i = 0, \dots, k-1$ , define  $A_i$  to be the closure of  $\{p \in \Gamma \mid \lambda_S(p) < i\}$  and define  $B_i = \{p \in \Gamma \mid \lambda_S(p) > i\}$ . We have  $\partial A_i = S_i$  and  $\partial B_i = S_{i+1}$ . For  $i = 0, \dots, k-1$ , let  $f_i$  be the unique element of  $R(\Gamma)$  which is 0 on  $A_i$  and 1 on  $B_i$ . Define  $P_i = \Delta(f_i)$ . By definition,

$$P_i = \sum_{v \in S_{i+1}} b_v^{(i)}(v) - \sum_{v \in S_i} a_v^{(i)}(v)$$

where  $b_v^{(i)} = \text{outdeg}_{B_i}(v) \geq 1$  and  $a_v^{(i)} = \text{outdeg}_{A_i}(v) \geq 1$ . It is straightforward to verify that we can choose positive integers  $c_0, c_1, \dots, c_{k-1}$  inductively, starting from  $c_{k-1}$  and working backwards, in such a way that  $P := \sum_{i=0}^{k-1} c_i P_i$  satisfies  $D + P \in |D|_q$ .  $\square$

Let  $\text{Meas}_+^d(\Gamma)$  be the space of nonnegative Borel measures of total mass  $d$  on  $\Gamma$ . By Alaoglu’s theorem,  $\text{Meas}_+^d(\Gamma)$  is compact in the weak\* topology.

**Lemma A.10.** *Let  $\Gamma$  be a metric graph.*

- (a) *If  $f_n$  converges to  $f$  pointwise on  $\Gamma$  with  $f_n, f \in \text{BDV}(\Gamma)$ , then  $\Delta(f_n)$  converges to  $\Delta(f)$  in the weak\* topology on  $\text{Meas}^0(\Gamma)$ .*
- (b) *Conversely, fix  $q \in \Gamma$  and suppose that  $\Delta(f_n)$  converges to  $\Delta(f)$  in the weak\* topology on  $\text{Meas}^0(\Gamma)$ , with  $f_n, f \in \text{BDV}(\Gamma)$  and  $f_n(q) = f(q) = 0$  for all  $n$ . Then  $f_n$  converges uniformly to  $f$  on  $\Gamma$ .*

*Proof.* (1) Since  $\text{CPA}(\Gamma)$  is dense in  $\mathcal{C}(\Gamma)$ , we just need to prove that  $\int g \Delta(f_n) \rightarrow \int g \Delta(f)$  for every  $g \in \text{CPA}(\Gamma)$ . But this is equivalent to  $\int f_n \Delta(g) \rightarrow \int f \Delta(g)$ , which follows from pointwise convergence since  $\Delta(g)$  is a discrete measure.

(2) This follows from the continuity of  $j_q(x, y)$  together with the identities  $f_n(x) = \int j_q(x, y) (\Delta f_n)(y)$  and  $f(x) = \int j_q(x, y) (\Delta f)(y)$ .  $\square$

Let  $D$  be an effective divisor of degree  $d$  on  $\Gamma$ . We can topologize  $|D|$  by thinking of it as a subset of  $\text{Meas}_+^d(\Gamma)$  and giving it the subspace topology.

**Theorem A.11.** *The space  $|D|$ , with its weak\* topology as subset of  $\text{Meas}_+^d(\Gamma)$ , is compact.*

*Proof.* Since  $\text{Meas}_+^d(\Gamma)$  is a compact Hausdorff space by Alaoglu's theorem, it suffices to show that  $|D|$  is a closed subspace. Suppose  $\mu_n \in |D|$  converges to  $\mu \in \text{Meas}_+^d(\Gamma)$ . We want to show that  $\mu \in |D|$ . Since each  $\mu_n$  is a discrete measure of total mass  $d$  with nonnegative integer masses, it follows easily from Alaoglu's theorem and the "Portmanteau theorem" (see [9, Proof of Proposition 9.5]) that  $\mu$  is also a discrete measure. Therefore  $\mu = \mu_D + \Delta(f)$  for some  $f \in \text{CPA}(\Gamma)$ . Since  $\mu_n = \mu_D + \Delta(f_n)$  with  $f_n \in R(\Gamma)$  and  $\mu_n \rightarrow \mu$ , it suffices to prove that if  $f_n \in R(\Gamma)$ ,  $f \in \text{BDV}(\Gamma)$ , and  $\Delta(f_n)$  converges to  $\Delta(f)$ , then  $f \in R(\Gamma)$ . We may assume without loss of generality that  $f_n(q) = f(q) = 0$  for all  $n$ , so by Lemma A.10(b)  $f_n$  converges uniformly to  $f$  on  $\Gamma$ . But then every directional derivative of  $f_n$  converges to the corresponding directional derivative of  $f$ . Since the slopes of  $f_n$  are integers, it follows that the slopes of  $f$  are integers as well. Thus  $f \in R(\Gamma)$  and  $\mu \in |D|$  as desired.  $\square$

Alternatively, one can prove this using the results of [30] as follows. Gathmann and Kerber show that  $|D|$  can be given the structure of a compact polyhedral complex by thinking of it as a subset of the compact space  $\Gamma^{(d)} = \Gamma^d/S_d$  (here  $S_d$  denotes the symmetric group on  $d$  letters). The natural map from  $|D|$  (with the Gathmann-Kerber topology) to  $\text{Meas}_+^d(\Gamma)$  is a continuous injection whose image is  $|D|$  with its weak\* topology. As the continuous image of a compact space,  $|D|$  is therefore compact in the weak\* topology. (Note that since a continuous bijection between compact Hausdorff spaces is automatically a homeomorphism, it follows that the Gathmann-Kerber and weak\* topologies on  $|D|$  are the same.)  $\square$

*Remark A.12.* One can also deduce Theorem A.11 from the results in [46].

We now give the promised proof of Theorem A.6.

*Proof of Theorem A.6.* By Proposition A.9 and the definition of reduced divisors, we can assume without loss of generality that  $D$  is effective. We endow  $|D|$  with the weak\* topology as above. The function  $\mathcal{E}_q : |D| \rightarrow \mathbb{R}$  is continuous since

$$\mathcal{E}_q(D') = \int \int j_q(x, y) d\mu_{D'-d(q)}(x) d\mu_{D'-d(q)}(y)$$

and  $j_q(x, y)$  is jointly continuous in  $x$  and  $y$ . Since  $|D|$  is compact,  $\mathcal{E}_q$  attains its minimum value at some effective divisor  $D_q \in |D|$ . By Theorem A.7,  $D_q$  is  $q$ -reduced. This proves the existence of a  $q$ -reduced divisor equivalent to  $D$ . The uniqueness is Corollary A.8.  $\square$

*Remark A.13.* The continuity of  $\mathcal{E}_q : |D| \rightarrow \mathbb{R}$  played an important role in the Proof of Theorem A.6. More generally, the energy pairing is continuous as a function from

$\text{Meas}^0(\Gamma) \times \text{Meas}^0(\Gamma)$  to  $\mathbb{R}$ . If we fix a reference point  $q \in \Gamma$ , this follows from the joint continuity of  $j_q(x, y)$  as a function of  $x$  and  $y$  together with the formula

$$\langle \mu, \nu \rangle = \int \int j_q(x, y) d\mu(x) d\nu(y) .$$

*Remark A.14.* The continuity of the energy pairing, together with the principle of least action (Lemma A.5), can be used to give a new proof of Amini’s theorem [3] that for  $D$  effective, the map from  $\Gamma$  to  $|D|$  sending a point  $q \in \Gamma$  to the unique  $q$ -reduced divisor  $D_q$  equivalent to  $D$  is continuous.

We conclude this section with an analogue of Theorem 4.14 for metric graphs. For a divisor  $D \in \text{Div}(\Gamma)$  of degree  $d$ , define its “total potential”  $b_q(D)$  by

$$(A.15) \quad b_q(D) := \int \int j_q(x, y) d\mu_{D-d(q)}(x) dy ,$$

where  $dy$  denotes the measure on  $\Gamma$  whose restriction to each edge is Lebesgue measure.

**Theorem A.16.** *Let  $D$  be a divisor on  $\Gamma$  which is effective outside  $q$ . Then  $D$  is  $q$ -reduced if and only if  $b_q(D) < b_q(D')$  for every divisor  $D' \neq D$  in  $|D|_q$ .*

The proof is similar to the proofs of Theorem 4.14 and Theorem A.7, so we omit it.

**A.3. Dhar’s algorithm for metric graphs and Luo’s theorem.** In this section, we briefly discuss some results from Ye Luo’s paper [42] and their relation to the material in the present paper.

The idea behind Dhar’s algorithm for metric graphs [42, Algorithm 2.5] is the same as for non-metric graphs. Let  $\Gamma$  be a metric graph, fix a point  $q \in \Gamma$ , and let  $D$  be a divisor on  $\Gamma$  which is effective outside  $q$ . Think of the metric graph  $\Gamma$  as being made of a flammable material. At every point  $p \in \Gamma$  with  $p \neq q$ , there are  $D(p)$  firefighters, each of whom can control fires in a single direction leading into  $p$ . A fire starts at  $q$  and proceeds along each direction emanating from  $q$ ; whenever there are fires approaching  $p$  in more than  $D(p)$  directions, the fire burns through  $p$  and proceeds to burn in all directions emanating from  $p$ . (In particular, if  $e$  is a segment in  $\Gamma$  consisting entirely of points  $p$  with  $D(p) = 0$ , then a fire starting at one endpoint of  $e$  will burn through  $e$  unobstructed.) The divisor  $D$  is  $q$ -reduced iff the fire eventually burns through all of  $\Gamma$ .

For a formal statement of Dhar’s algorithm for metric graphs and a proof of correctness, see [42, Algorithm 2.5 and Lemma 2.6].

In order to compute the unique  $q$ -reduced divisor equivalent to a given divisor  $D$ , one can proceed along the lines of Algorithm 4 above. Starting with a divisor  $D$ , the first step is to replace  $D$  with an equivalent divisor which is effective outside  $q$ . This is accomplished by following the proof of Proposition A.9. The second step is to iterate Dhar’s algorithm until we reach a  $q$ -reduced divisor. More specifically, if  $D$  is not yet reduced then by running Dhar’s algorithm on  $D$  we obtain a proper connected subset  $Y \neq \Gamma$  of  $\Gamma$  which contains  $q$  (the “burned portion” of  $\Gamma$ ). Choose a connected

component  $X$  of the complement of  $Y$  and (as in the proof of Lemma A.4) choose  $\epsilon > 0$  maximal with respect to the property that there is a rational function  $f \in R(\Gamma)$  which is 0 on  $X$  and  $\epsilon$  outside of an  $\epsilon$ -neighborhood of  $X$ , with constant slope 1 in between. By construction, the divisor  $D + \Delta(f)$  is still effective outside  $q$ . Replacing  $D$  by  $D + \Delta(f)$  and iterating this procedure, one obtains (after finitely many iterations) the unique  $q$ -reduced divisor equivalent to  $D$ .

For a formal statement of this algorithm and a proof that it terminates after a finite number of iterations, see [42, Algorithm 2.12 and Theorem 2.14].

In (A.15) we defined a metric graph analogue of the functional  $b_q$  from §4. As in Proposition 4.5(b), this functional is a “monovariant” which decreases by an explicit function of  $\epsilon$  when  $D$  is replaced by  $D + \Delta(f)$  (with  $f$  as above):

$$\begin{aligned} b_q(D + \Delta(f)) - b_q(D) &= \int \int j_q(x, y) \Delta_x(f) dy \\ &= \int \int f(x) \Delta_x(j_q(x, y)) dy \\ &= \int \int f(x) (\delta_y(x) - \delta_q(x)) dy \\ &= \int (f(y) - f(q)) dy \\ &= -(l(X)\epsilon + \frac{\lambda(X)}{2}\epsilon^2), \end{aligned}$$

where  $l(X)$  is the total length of  $X$  and  $\lambda(X)$  is the size of the  $(X, \Gamma \setminus X)$ -cut.

It would be interesting to give explicit lower bounds for the  $\epsilon$ 's which can appear and thus obtain a running-time analysis of Luo's Algorithm 2.12 similar to our Proposition 5.6(c); we leave this as an open problem for future research.

## REFERENCES

- [1] D. J. Aldous. The random walk construction of uniform spanning trees and uniform labelled trees. *SIAM J. Discrete Math.*, 3(4):450–465, 1990.
- [2] N. Alon, I. Krasikov, and Y. Peres. Reflection sequences. *Amer. Math. Monthly*, 96(9):820–823, 1989.
- [3] O. Amini. Reduced divisors and embeddings of tropical curves. 2010. Preprint available at [arXiv:1007.5364v1](https://arxiv.org/abs/1007.5364v1).
- [4] R. Bacher, P. de la Harpe, and T. Nagnibeda. The lattice of integral flows and the lattice of integral cuts on a finite graph. *Bull. Soc. Math. France*, 125(2):167–198, 1997.
- [5] P. Bak, C. Tang, and K. Wiesenfeld. Self-organized criticality. *Phys. Rev. A* (3), 38(1):364–374, 1988.
- [6] M. Baker and X. Faber. Metrized graphs, Laplacian operators, and electrical networks. In *Quantum graphs and their applications*, volume 415 of *Contemp. Math.*, pages 15–33. Amer. Math. Soc., Providence, RI, 2006.
- [7] M. Baker and S. Norine. Riemann-Roch and Abel-Jacobi theory on a finite graph. *Adv. Math.*, 215(2):766–788, 2007.
- [8] M. Baker and R. Rumely. Harmonic analysis on metrized graphs. *Canad. J. Math.*, 59(2):225–275, 2007.

- [9] M. Baker and R. Rumely. *Potential theory and dynamics on the Berkovich projective line*, volume 159 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010.
- [10] A. Ben-Israel and T. N. E. Greville. *Generalized inverses*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 15. Springer-Verlag, New York, second edition, 2003. Theory and applications.
- [11] B. A. Benson, D. Chakrabarty, and P. Tetali.  $G$ -parking functions, acyclic orientations and spanning trees. *Discrete Mathematics*, 310(8):1340–1353, 2010.
- [12] N. Biggs. *Algebraic graph theory*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 1993.
- [13] N. Biggs. Algebraic potential theory on graphs. *Bull. London Math. Soc.*, 29(6):641–682, 1997.
- [14] N. Biggs. Chip-firing and the critical group of a graph. *J. Algebraic Combin.*, 9(1):25–45, 1999.
- [15] N. Biggs. The Tutte polynomial as a growth function. *J. Algebraic Combin.*, 10(2):115–133, 1999.
- [16] N. Biggs and P. Winkler. Chip-firing and the chromatic polynomial. Report LSE-CDAM-97-03, Centre for Discrete and Applicable Mathematics, London School of Economics, London, U.K., February 1997.
- [17] A. Björner, L. Lovász, and P. W. Shor. Chip-firing games on graphs. *European J. Combin.*, 12(4):283–291, 1991.
- [18] B. Bollobás. *Modern graph theory*, volume 184 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1998.
- [19] A. Broder. Generating random spanning trees. *Foundations of Computer Science, 30th Annual IEEE Symposium on*, pages 442–447, 1989.
- [20] D. Chebikin and P. Pylyavskyy. A family of bijections between  $G$ -parking functions and spanning trees. *J. Combin. Theory Ser. A*, 110(1):31–41, 2005.
- [21] T. Chinburg and R. Rumely. The capacity pairing. *J. Reine Angew. Math.*, 434:1–44, 1993.
- [22] F. Chung and R. B. Ellis. A chip-firing game and Dirichlet eigenvalues. *Discrete Math.*, 257(2-3):341–355, 2002. Kleitman and combinatorics: a celebration (Cambridge, MA, 1999).
- [23] C. J. Colbourn, W. J. Myrvold, and E. Neufeld. Two algorithms for unranking arborescences. *J. Algorithms*, 20(2):268–281, 1996.
- [24] D. Coppersmith, U. Feige, and J. Shearer. Random walks on regular and irregular graphs. *SIAM J. Discrete Math.*, 9(2):301–308, 1996.
- [25] D. Coppersmith and S. Winograd. Matrix multiplication via arithmetic progressions. *Journal of Symbolic Computation*, 9(3):251 – 280, 1990. Computational algebraic complexity editorial.
- [26] R. Cori and Y. Le Borgne. The sand-pile model and Tutte polynomials. *Adv. in Appl. Math.*, 30(1-2):44–52, 2003. Formal power series and algebraic combinatorics (Scottsdale, AZ, 2001).
- [27] R. Cori, D. Rossin, and B. Salvy. Polynomial ideals for sandpiles and their Gröbner bases. *Theoret. Comput. Sci.*, 276(1-2):1–15, 2002.
- [28] D. Dhar. Self-organized critical state of sandpile automaton models. *Phys. Rev. Lett.*, 64(14):1613–1616, Apr 1990.
- [29] A. Gabrielov. Abelian avalanches and Tutte polynomials. *Phys. A*, 195(1-2):253–274, 1993.
- [30] A. Gathmann and M. Kerber. A Riemann-Roch theorem in tropical geometry. *Math. Z.*, 259(1):217–230, 2008.
- [31] A. Guénoche. Random spanning tree. *J. Algorithms*, 4(3):214–220, 1983.
- [32] J. van den Heuvel. Algorithmic aspects of a chip-firing game. *Combin. Probab. Comput.*, 10(6):505–529, 2001.
- [33] J. Hladký, D. Král, and S. Norine. Rank of divisors on tropical curves. 2007. Preprint available at [arXiv:0709.4485](https://arxiv.org/abs/0709.4485).
- [34] A. E. Holroyd, L. Levine, K. Mészáros, Y. Peres, J. Propp, and D. B. Wilson. Chip-firing and rotor-routing on directed graphs. In *In and out of equilibrium. 2*, volume 60 of *Progr. Probab.*, pages 331–364. Birkhäuser, Basel, 2008.

- [35] E. Kaltofen and G. Villard. On the complexity of computing determinants. *Comput. Complexity*, 13(3-4):91–130, 2004.
- [36] J. A. Kelner and A. Madry. Faster generation of random spanning trees. In *Foundations of Computer Science, 2009., 50th Annual Symposium on*, pages 13–21, Oct 2009.
- [37] G. Kirchhoff. Über die Auflösung der Gleichungen, auf welche man bei der Untersuchung der linearen Verteilung galvanischer Ströme geführt wird. *Ann. Phys. Chem.*, (72):497–508, 1847.
- [38] V. G. Kulkarni. Generating random combinatorial objects. *J. Algorithms*, 11(2):185–207, 1990.
- [39] L. Levine and J. Propp. What is ... a sandpile? *Notices Amer. Math. Soc.*, 57(8):976–979, 2010.
- [40] R. Lipton. *A New Approach To Random Spanning Trees*. Blog post (2009), available at <http://rjlipton.wordpress.com/2009/07/15/a-new-approach-to-random-spanning-trees/>
- [41] D. Lorenzini. Arithmetical graphs. *Math. Ann.*, 285(3):481–501, 1989.
- [42] Y. Luo. Rank-determining sets of metric graphs. *J. Comb. Theory Ser. A*, 118:1775–1793, August 2011.
- [43] R. Lyons and Y. Peres. *Probability on Trees and Networks*. Cambridge University Press, 2011. In preparation. Current version available at <http://mypage.iu.edu/~rdlyons/>.
- [44] S.N. Majumdar and D. Dhar. Equivalence between the abelian sandpile model and the  $q \rightarrow 0$  limit of the potts model. *Physica A: Statistical Mechanics and its Applications*, 185(1-4):129–145, 1992.
- [45] M. Manjunath. The rank of a divisor on a finite graph: geometry and computation. 2011. Preprint available at [arXiv:1111.7251v1](https://arxiv.org/abs/1111.7251v1).
- [46] G. Mikhalkin and I. Zharkov. Tropical curves, their Jacobians and theta functions. In *Curves and abelian varieties*, volume 465 of *Contemp. Math.*, pages 203–230. Amer. Math. Soc., Providence, RI, 2008.
- [47] S. Mozes. Reflection processes on graphs and Weyl groups. *J. Combin. Theory Ser. A*, 53(1):128–142, 1990.
- [48] A. Postnikov and B. Shapiro. Trees, parking functions, syzygies, and deformations of monomial ideals. *Trans. Amer. Math. Soc.*, 356(8):3109–3142 (electronic), 2004.
- [49] M. Raynaud. Spécialisation du foncteur de Picard. *Inst. Hautes Études Sci. Publ. Math.*, (38):27–76, 1970.
- [50] D. Saunders and Z. Wan. Smith normal form of dense integer matrices fast algorithms into practice. In *Proceedings of the 2004 international symposium on Symbolic and algebraic computation, ISSAC '04*, pages 274–281, New York, NY, USA, 2004. ACM.
- [51] F. Shokrieh. The monodromy pairing and discrete logarithm on the Jacobian of finite graphs. *J. Math. Cryptol.*, 4(1):43–56, 2010.
- [52] S. Smirnov. How do research problems compare with IMO problems? In D. Schleicher and M. Lackmann, editors, *An Invitation to Mathematics*, pages 71–83. Springer Berlin Heidelberg, 2011.
- [53] G. Tardos. Polynomial bound for a chip firing game on graphs. *SIAM J. Discrete Math.*, 1(3):397–398, 1988.
- [54] E. Wegert and C. Reiher. Relaxation procedures on graphs. *Discrete Appl. Math.*, 157(9):2207–2216, 2009.
- [55] D. B. Wilson. Generating random spanning trees more quickly than the cover time. In *Proceedings of the Twenty-eighth Annual ACM Symposium on the Theory of Computing (Philadelphia, PA, 1996)*, pages 296–303, New York, 1996. ACM.
- [56] P. Winkler. *Mathematical puzzles: a connoisseur's collection*. A K Peters Ltd., Natick, MA, 2004.

*E-mail address:* mbaker@math.gatech.edu

*E-mail address:* shokrieh@math.gatech.edu

GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GEORGIA 30332-0160, USA