

# Relations for ideal gas

March 23, 2025

## 1 Ideal gas equation

$$P\nu = RT$$

## 2 Isentropic process

$$s = \text{constant}$$

$$\frac{P_2}{P_1} = \left(\frac{\nu_1}{\nu_2}\right)^k$$

$$\frac{T_2}{T_1} = \left(\frac{\nu_1}{\nu_2}\right)^{k-1}$$

$$\frac{T_2}{T_1} = \left(\frac{\nu_1}{\nu_2}\right)^{(k-1)/k}$$

## 3 Isobaric process

$$P = \text{constant}$$

$$s_2 = s_1 + c_\nu \ln \frac{T_2}{T_1} + R \ln \frac{\nu_2}{\nu_1}$$

## 4 Isochoric process

$$\nu = \text{constant}$$

$$s_2 = s_1 + c_\nu \ln \frac{T_2}{T_1} + R \ln \frac{\nu_2}{\nu_1}$$

## 5 Isothermal process

$$T = \text{constant}$$

$$P\nu = \text{constant}$$

$$s_2 = s_1 + c_\nu \ln \frac{T_2}{T_1} + R \ln \frac{\nu_2}{\nu_1}$$

## 6 var specific heat

$$s_2 - s_1 = \int_1^2 c_{v0} \frac{dT}{T} + R \ln \frac{\nu_2}{\nu_1}$$

$$c_{p0} = C_0 + C_1\theta + C_2\theta^2 + C_3\theta^3$$

where,  $\theta = T/1000$  and  $T$  is in Kelvin.

$$c_{p0} = C_0 + C_1 \frac{T}{1000} + C_2 \left( \frac{T}{1000} \right)^2 + C_3 \left( \frac{T}{1000} \right)^3$$

Note theta

$$c_{v0} = c_{p0} - R$$

$$\int_1^2 c_{v0} \frac{dT}{T} = \int_1^2 (c_{p0} - R) \frac{dT}{T} = \int_1^2 c_{p0} \frac{dT}{T} - \int_1^2 R \frac{dT}{T} = \int_1^2 c_{p0} \frac{dT}{T} - R \ln \frac{T_2}{T_1}$$

$$\begin{aligned} \int_1^2 c_{p0} \frac{dT}{T} &= \int_1^2 \left( C_0 + C_1 \frac{T}{1000} + C_2 \left( \frac{T}{1000} \right)^2 + C_3 \left( \frac{T}{1000} \right)^3 \right) \frac{dT}{T} \\ &= C_0 \int_1^2 \frac{dT}{T} + \frac{C_1}{1000} \int_1^2 dT + \frac{C_2}{10^6} \int_1^2 T dT + \frac{C_3}{10^9} \int_1^2 T^2 dT \\ &= C_0 \ln \frac{T_2}{T_1} + \frac{C_1}{1000} (T_2 - T_1) + \frac{C_2}{2 \cdot 10^6} (T_2^2 - T_1^2) + \frac{C_3}{3 \cdot 10^9} (T_2^3 - T_1^3) \end{aligned}$$

As a result,

$$\int_1^2 c_{v0} \frac{dT}{T} = (C_0 - R) \ln \frac{T_2}{T_1} + \frac{C_1}{1000} (T_2 - T_1) + \frac{C_2}{2 \cdot 10^6} (T_2^2 - T_1^2) + \frac{C_3}{3 \cdot 10^9} (T_2^3 - T_1^3)$$

Finally,

$$s_2 - s_1 = (C_0 - R) \ln \frac{T_2}{T_1} + \frac{C_1}{1000} (T_2 - T_1) + \frac{C_2}{2 \cdot 10^6} (T_2^2 - T_1^2) + \frac{C_3}{3 \cdot 10^9} (T_2^3 - T_1^3) + R \ln \frac{\nu_2}{\nu_1}$$

For isentropic process,  $s_2 = s_1$ .

$$0 = (C_0 - R) \ln \frac{T_2}{T_1} + \frac{C_1}{1000} (T_2 - T_1) + \frac{C_2}{2 \cdot 10^6} (T_2^2 - T_1^2) + \frac{C_3}{3 \cdot 10^9} (T_2^3 - T_1^3) + R \ln \frac{\nu_2}{\nu_1}$$

If specific volume  $\nu_2$  is given, the unknown is  $T_2$ . Writing the function in terms of  $T$ ,

$$f(T) = (C_0 - R) \ln \frac{T}{T_1} + \frac{C_1}{1000} (T - T_1) + \frac{C_2}{2 \cdot 10^6} (T^2 - T_1^2) + \frac{C_3}{3 \cdot 10^9} (T^3 - T_1^3) + R \ln \frac{\nu_2}{\nu_1}$$

## 7 Newton Raphson

For NR, derivative of  $f(T)$  with respect to  $T$  is needed.

$$f'(T) = \frac{C_0 - R}{T} + \frac{C_1}{1000} + \frac{C_2}{10^6} T + \frac{C_3}{10^9} T^2$$

Iterate

$$T_{\text{new}} = T - \frac{f(T)}{f'(T)}$$

until  $T_{\text{new}} - T < \epsilon$

## 8 var specific heat 2

$$s_2 - s_1 = \int_1^2 c_{p0} \frac{dT}{T} - R \ln \frac{P_2}{P_1}$$

$$c_{p0} = C_0 + C_1\theta + C_2\theta^2 + C_3\theta^3$$

where,  $\theta = T/1000$  and  $T$  is in Kelvin.

$$c_{p0} = C_0 + C_1 \frac{T}{1000} + C_2 \left( \frac{T}{1000} \right)^2 + C_3 \left( \frac{T}{1000} \right)^3$$

$$\begin{aligned} \int_1^2 c_{p0} \frac{dT}{T} &= \int_1^2 \left( C_0 + C_1 \frac{T}{1000} + C_2 \left( \frac{T}{1000} \right)^2 + C_3 \left( \frac{T}{1000} \right)^3 \right) \frac{dT}{T} \\ &= C_0 \int_1^2 \frac{dT}{T} + \frac{C_1}{1000} \int_1^2 dT + \frac{C_2}{10^6} \int_1^2 T dT + \frac{C_3}{10^9} \int_1^2 T^2 dT \\ &= C_0 \ln \frac{T_2}{T_1} + \frac{C_1}{1000} (T_2 - T_1) + \frac{C_2}{2 \cdot 10^6} (T_2^2 - T_1^2) + \frac{C_3}{3 \cdot 10^9} (T_2^3 - T_1^3) \end{aligned}$$

Finally,

$$s_2 - s_1 = C_0 \ln \frac{T_2}{T_1} + \frac{C_1}{1000} (T_2 - T_1) + \frac{C_2}{2 \cdot 10^6} (T_2^2 - T_1^2) + \frac{C_3}{3 \cdot 10^9} (T_2^3 - T_1^3) - R \ln \frac{P_2}{P_1}$$

For isentropic process,  $s_2 = s_1$ .

$$0 = C_0 \ln \frac{T_2}{T_1} + \frac{C_1}{1000} (T_2 - T_1) + \frac{C_2}{2 \cdot 10^6} (T_2^2 - T_1^2) + \frac{C_3}{3 \cdot 10^9} (T_2^3 - T_1^3) - R \ln \frac{P_2}{P_1}$$

If pressure  $P_2$  is given, the unknown is  $T_2$ . Writing the function in terms of  $T$ ,

$$f(T) = C_0 \ln \frac{T_2}{T_1} + \frac{C_1}{1000} (T_2 - T_1) + \frac{C_2}{2 \cdot 10^6} (T_2^2 - T_1^2) + \frac{C_3}{3 \cdot 10^9} (T_2^3 - T_1^3) - R \ln \frac{P_2}{P_1}$$

## 9 Newton Raphson

For NR, derivative of  $f(T)$  with respect to  $T$  is needed.

$$f'(T) = \frac{C_0}{T} + \frac{C_1}{1000} + \frac{C_2}{10^6} T + \frac{C_3}{10^9} T^2$$

Iterate

$$T_{\text{new}} = T - \frac{f(T)}{f'(T)}$$

until  $T_{\text{new}} - T < \epsilon$

## 10 $T$ is given find $\nu$

Transpose  $\nu_2$  in the following equation

$$0 = (C_0 - R) \ln \frac{T}{T_1} + \frac{C_1}{1000}(T - T_1) + \frac{C_2}{2 \cdot 10^6}(T^2 - T_1^2) + \frac{C_2}{3 \cdot 10^9}(T^3 - T_1^3) + R \ln \frac{\nu_2}{\nu_1}$$

such that

$$-R(\ln \nu_2 - \ln \nu_1) = (C_0 - R) \ln \frac{T}{T_1} + \frac{C_1}{1000}(T - T_1) + \frac{C_2}{2 \cdot 10^6}(T^2 - T_1^2) + \frac{C_2}{3 \cdot 10^9}(T^3 - T_1^3)$$

$$-R \ln \nu_2 + R \ln \nu_1 = (C_0 - R) \ln \frac{T}{T_1} + \frac{C_1}{1000}(T - T_1) + \frac{C_2}{2 \cdot 10^6}(T^2 - T_1^2) + \frac{C_2}{3 \cdot 10^9}(T^3 - T_1^3)$$

$$-R \ln \nu_2 = (C_0 - R) \ln \frac{T}{T_1} + \frac{C_1}{1000}(T - T_1) + \frac{C_2}{2 \cdot 10^6}(T^2 - T_1^2) + \frac{C_2}{3 \cdot 10^9}(T^3 - T_1^3) - R \ln \nu_1$$

$$\ln \nu_2 = \frac{R - C_0}{R} \ln \frac{T}{T_1} - \frac{C_1}{1000R}(T - T_1) - \frac{C_2}{2 \cdot 10^6 R}(T^2 - T_1^2) - \frac{C_2}{3 \cdot 10^9 R}(T^3 - T_1^3) - \ln \nu_1$$

$$\nu_2 = \exp \left( \frac{R - C_0}{R} \ln \frac{T}{T_1} - \frac{C_1}{1000R}(T - T_1) - \frac{C_2}{2 \cdot 10^6 R}(T^2 - T_1^2) - \frac{C_2}{3 \cdot 10^9 R}(T^3 - T_1^3) - \ln \nu_1 \right)$$

## 11 qs and qr isochoric

$$q_s = \int_1^2 c_{\nu 0} dT$$

$$c_{p0} = C_0 + C_1\theta + C_2\theta^2 + C_3\theta^3$$

$$c_{\nu 0} = c_{p0} - R$$

$$c_{\nu 0} = C_0 + C_1\theta + C_2\theta^2 + C_3\theta^3 - R$$

$$c_{\nu 0} = C_0 + \frac{C_1}{10^3}T + \frac{C_2}{10^6}T^2 + \frac{C_3}{10^9}T^3 - R$$

$$q_s = \int_1^2 \left( C_0 + \frac{C_1}{10^3}T + \frac{C_2}{10^6}T^2 + \frac{C_3}{10^9}T^3 - R \right) dT$$

$$q_s = (C_0 - R) \int_1^2 dT + \frac{C_1}{10^3} \int_1^2 T dT + \frac{C_2}{10^6} \int_1^2 T^2 dT + \frac{C_3}{10^9} \int_1^2 T^3 dT$$

$$q_s = (C_0 - R)(T_2 - T_1) + \frac{C_1}{2 \cdot 10^3}(T_2^2 - T_1^2) + \frac{C_2}{3 \cdot 10^6}(T_2^3 - T_1^3) + \frac{C_3}{4 \cdot 10^9}(T_2^4 - T_1^4)$$

Unknown is  $T_2$ .

$$q_s = (C_0 - R)(T - T_1) + \frac{C_1}{2 \cdot 10^3}(T^2 - T_1^2) + \frac{C_2}{3 \cdot 10^6}(T^3 - T_1^3) + \frac{C_3}{4 \cdot 10^9}(T^4 - T_1^4)$$

$$f(T) = (C_0 - R)(T - T_1) + \frac{C_1}{2 \cdot 10^3}(T^2 - T_1^2) + \frac{C_2}{3 \cdot 10^6}(T^3 - T_1^3) + \frac{C_3}{4 \cdot 10^9}(T^4 - T_1^4) - q_s$$

Derivative

$$f'(T) = (C_0 - R) + \frac{C_1}{10^3}T + \frac{C_2}{10^6}T^2 + \frac{C_3}{10^9}T^3$$