

Compact Answers to Temporal Regular Path Queries (Supplementary Material)

ACM Reference Format:

. 2023. Compact Answers to Temporal Regular Path Queries (Supplementary Material). In *Proceedings of ACM Conference (Conference'17)*. ACM, New York, NY, USA, 6 pages. <https://doi.org/10.1145/nnnnnnn.nnnnnnn>

1 INTRODUCTION

This document provides detailed definitions and proofs for the article *Compact answers to Temporal Regular Path Queries*, submitted at CIKM 2023.

As opposed to the structure adopted in the article, the result here are grouped by topic (inductive representation, finiteness, complexity, etc) rather than representation ($\mathcal{U}^{[t]}$, $\mathcal{U}^{[d]}$, etc.). This allows us to emphasize which proofs differ from one representation to the other.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

Conference'17, July 2017, Washington, DC, USA

© 2023 Association for Computing Machinery.

ACM ISBN 978-x-xxxx-xxxx-x/YY/MM...\$15.00

<https://doi.org/10.1145/nnnnnnn.nnnnnnn>

2 INDUCTIVE REPRESENTATION

2.1 $\langle q \rangle_G^{[t,d],b,e}$

2.1.1 Definition.

2.1.2 Correctness.

Operator $\text{path}_1/\text{path}_2$.

Let $\mathbf{u}_1 = \langle o_1, o_2, \tau_1, \delta_1 \rangle$ and $\mathbf{u}_2 = \langle o_3, o_4, \tau_2, \delta_2 \rangle$, with $o_2 = o_3$.

Let also $\mathbf{u}_1 \bowtie \mathbf{u}_2 = \langle o_1, o_4, \tau'_1, \delta_1 + \delta_2, b, e \rangle$, with

$$\begin{aligned}\tau &= (((\tau_1 + \delta_1) \cap \tau_2) \ominus \delta_1) \cap \tau_1 \\ b &= \max(b_1, b_2 - b_{\delta_1}) \\ e &= \min(e_1, e_2 - e_{\delta_1})\end{aligned}$$

For $i \in \{1, 2\}$ and $t \in \tau_i$, we use $\delta_i(t)$ for the interval

$$\delta_i \downarrow b_{\delta_i} + \max(0, b_i - t), e_{\delta_i} - \max(0, t - e_i) \downarrow \delta_i$$

And similarly to what we did for $\mathcal{U}^{[t,d]}$, we use R_i for be the binary relation over \mathcal{T} specified by the time points and distances in \mathbf{u}_i , i.e. $R_i = \{(t, t + d) \mid t \in \tau_i, d \in \delta_i(t)\}$.

Then the intervals in the set $\mathbf{u}_1 \bowtie \mathbf{u}_2$ should intuitively represent this relation $R_1 \bowtie R_2$, i.e. we need to prove that (i) $\tau = \text{dom}(R_1 \bowtie R_2)$ and that (ii) for each $t \in \tau$,

$$t + \delta(t) = \{t' \mid (t, t') \in R_1 \bowtie R_2\}$$

For (ii), let $t \in \tau$. We use

- a for the least value s.t. $(t, a) \in \text{range}(R_1) \cap \text{dom}(R_2)$, and
- a' for the least value s.t. $(a, a') \in R_2$

Then a' is also the least value s.t. $(t, a') \in R_1 \bowtie R_2$.

Analogously, we use z for the greatest value s.t. $(t, z) \in \text{range}(R_1) \cap \text{dom}(R_2)$, and z' for the greatest value s.t. $(z, z') \in R_2$. Then z' is also the greatest value s.t. $(t, z') \in R_1 \bowtie R_2$.

From Lemma ??:

- $\{t\} \times (t + [a, z]) \subseteq R_1$, and
- $[a, b] \times [a', z'] \subseteq R_2$

Therefore $[a', z'] = \{c \mid (t, c) \in R_1 \bowtie R_2\}$.

To conclude the proof, we show that $t + \delta_t = [a, z]$.

We only prove that $t + b_{\delta_t} = a$ (the proof that $t + e_{\delta_t} = z$ is symmetric).

Following the definition of b , we consider 2 cases:

- (1) $b_1 < b_2 - b_{\delta_1}$
- (2) $b_1 \geq b_2 - b_{\delta_1}$

For Case (1), we get

$$b_1 < b_2 - b_{\delta_1} \tag{1}$$

$$\max(b_1, b_2 - b_{\delta_1}) = b_2 - b_{\delta_1} \tag{2}$$

$$b = b_2 - b_{\delta_1} \quad \text{from the definition of } b \tag{3}$$

And in Case (1) still, we get:

$$b_1 < b_2 - b_{\delta_1} \tag{4}$$

$$0 < b_2 - b_{\delta_1} - b_1 \tag{5}$$

$$\max(0, b_2 - b_{\delta_1} - b_1) = b_2 - b_{\delta_1} - b_1 \tag{6}$$

Next, we consider two subcases:

- (i) $t < b_2 - b_{\delta_1}$
- (ii) $t \geq b_2 - b_{\delta_1}$

In Case (i), we get

$$t < b_2 - b_{\delta_1} \quad (7)$$

$$0 < b_2 - b_{\delta_1} - t \quad (8)$$

$$\max(0, b_2 - b_{\delta_1} - t) = b_2 - b_{\delta_1} - t \quad (9)$$

Now from the definition of δ_t ,

$$b_{\delta_t} = b_{\delta_1} + b_{\delta_2} + \max(0, b - t) \quad (10)$$

$$= b_{\delta_1} + b_{\delta_2} + \max(0, b_2 - b_{\delta_1} - t) \quad \text{from (3)} \quad (11)$$

$$= b_{\delta_1} + b_{\delta_2} + b_2 - b_{\delta_1} - t \quad \text{from (9)} \quad (12)$$

$$= b_{\delta_2} + b_2 - t \quad (13)$$

$$b_{\delta_t} + t = b_{\delta_2} + b_2 - t + t \quad (14)$$

$$= b_{\delta_2} + b_2 \quad (15)$$

Next, from the definition of a'

$$a' = b_{\delta_2(a)} + a \quad (16)$$

$$= b_{\delta_2} + \max(0, b_2 - a) + a \quad (17)$$

And, from the definition of a

$$a = b_{\delta_1(t)} + t \quad (18)$$

$$= b_{\delta_1} + \max(0, b_1 - t) + t \quad (19)$$

Then we have two further subcases:

(I) $t \geq b_1$, or

(II) $t < b_1$

In case (I):

$$t \geq b_1 \quad (20)$$

$$0 \geq b_1 - t \quad (21)$$

$$\max(0, b_1 - t) = 0 \quad (22)$$

$$a = b_{\delta_1} + t \quad \text{from (19)} \quad (23)$$

$$\max(0, b_2 - a) = \max(0, b_2 - b_{\delta_1} - t) \quad (24)$$

$$= b_2 - b_{\delta_1} - t \quad \text{from (9)} \quad (25)$$

$$= b_2 - a \quad \text{from (23)} \quad (26)$$

In case (II):

$$t < b_1 \quad (27)$$

$$0 < b_1 - t \quad (28)$$

$$\max(0, b_1 - t) = b_1 - t \quad (29)$$

$$a = b_{\delta_1} + b_1 - t + t \quad \text{from (19)} \quad (30)$$

$$= b_{\delta_1} + b_1 \quad (31)$$

$$\max(0, b_2 - a) = \max(0, b_2 - b_{\delta_1} - b_1) \quad (32)$$

$$= b_2 - b_{\delta_1} - b_1 \quad \text{from (6)} \quad (33)$$

$$= b_2 - a \quad \text{from (31)} \quad (34)$$

$$(35)$$

So in both cases (I) and (II), we get

$$\max(0, b_2 - a) = b_2 - a$$

Therefore from (17)

$$a' = b_{\delta_2} + b_2 - a + a \quad (36)$$

$$= b_{\delta_2} + b_2 \quad (37)$$

$$= t + b_{\delta_t} \quad \text{from (15)} \quad (38)$$

which concludes the proof for Case (1)- (i).

We continue with Case (1)- (ii).

From Case (ii):

$$t \geq b_2 - b_{\delta_1} \quad (39)$$

$$0 \geq b_2 - b_{\delta_1} - t \quad (40)$$

$$\max(0, b_2 - b_{\delta_1} - t) = 0 \quad (41)$$

Now from the definition of δ_t :

$$b_{\delta_t} = b_{\delta_1} + b_{\delta_2} + \max(0, b - t) \quad (42)$$

$$= b_{\delta_1} + b_{\delta_2} + \max(0, b_2 - b_1 - t) \quad \text{from (3)} \quad (43)$$

$$= b_{\delta_1} + b_{\delta_2} \quad \text{from (41)} \quad (44)$$

$$b_{\delta_t} + t = b_{\delta_1} + b_{\delta_2} + t \quad (45)$$

Next, from Case (1) and Case (ii), by transitivity, we get

$$b_1 \leq t \quad (46)$$

$$\max(0, b_1 - t) = 0 \quad (47)$$

And from the definition of a

$$a = b_{\delta_1(t)} + t \quad (48)$$

$$= b_{\delta_1} + \max(0, b_1 - t) + t \quad (49)$$

$$= b_{\delta_1} + t \quad \text{from (47)} \quad (50)$$

$$\geq b_{\delta_1} + b_2 - b_{\delta_1} \quad \text{from Case (ii)} \quad (51)$$

$$\geq b_2 \quad (52)$$

$$0 \geq b_2 - a \quad (53)$$

$$\max(0, b_2 - a) = 0 \quad (54)$$

Therefore from (17) and (54)

$$a' = b_{\delta_2} + a \quad (55)$$

$$= b_{\delta_2} + b_{\delta_1} + t \quad \text{from (50)} \quad (56)$$

$$= b_{\delta_t} + t \quad \text{from (15)} \quad (57)$$

which concludes the proof for Case (1)- (ii).

We continue with Case (2).

In this case, we get

$$b_1 \geq b_2 - b_{\delta_1} \quad (58)$$

$$\max(b_1, b_2 - b_{\delta_1}) = b_1 \quad (59)$$

$$b = b_1 \quad \text{from the definition of } b \quad (60)$$

And from Case (2) still, we derive

$$b_1 \geq b_2 - b_{\delta_1} \quad (61)$$

$$0 \geq b_2 - b_{\delta_1} - b_1 \quad (62)$$

$$\max(0, -b_{\delta_1} - b_1) = 0 \quad (63)$$

As well as

$$b_1 \geq b_2 - b_{\delta_1} \quad (64)$$

$$b_1 + b_{\delta_1} \geq b_2 \quad (65)$$

Next, we distinguish two subcases, namely

(a) $t < b_1$ and

(b) $t \geq b_1$

We start with Case (a).

In this case,

$$t < b_1 \quad (66)$$

$$0 < b_1 - t \quad (67)$$

$$\max(0, b_1 - t) = b_1 - t \quad (68)$$

And from the definition of δ_t :

$$b_{\delta_t} = b_{\delta_1} + b_{\delta_2} + \max(0, b - t) \quad (69)$$

$$= b_{\delta_1} + b_{\delta_2} + \max(0, b_1 - t) \quad \text{from (60)} \quad (70)$$

$$= b_{\delta_1} + b_{\delta_2} + b_1 - t \quad \text{from (68)} \quad (71)$$

$$b_{\delta_t} + t = b_{\delta_1} + b_{\delta_2} + b_1 - t + t \quad (72)$$

$$= b_{\delta_1} + b_{\delta_2} + b_1 \quad (73)$$

Next, from the definition of a

$$a = b_{\delta_1(t)} + t \quad (74)$$

$$= \max(0, b_1 - t) + b_{\delta_1} + t \quad (75)$$

$$= b_1 - t + b_{\delta_1} + t \quad \text{from (68)} \quad (76)$$

$$= b_1 + b_{\delta_1} \quad (77)$$

So from (65)

$$a \geq b_2 \quad (78)$$

$$0 \geq b_2 - a \quad (79)$$

$$\max(0, b_2 - a) = 0 \quad (80)$$

$$b_{\delta_2} + \max(0, b_2 - a) = b_{\delta_2} \quad (81)$$

$$b_{\delta_2(a)} = b_{\delta_2} \quad (82)$$

$$b_{\delta_2(a)} + a = b_{\delta_2} + a \quad (83)$$

$$a' = b_{\delta_2} + a \quad \text{from the definition of } a' \quad (84)$$

$$a' = b_{\delta_2} + b_1 + b_{\delta_1} \quad \text{from (77)} \quad (85)$$

$$a' = b_{\delta_t} + t \quad \text{from (73)} \quad (86)$$

which concludes the proof for Case (2)- (a).

We end with Case (2)- (b). In this case,

$$t \geq b_1 \quad (87)$$

$$0 \geq b_1 - t \quad (88)$$

$$\max(0, b_1 - t) = 0 \quad (89)$$

And from the definition of δ_t :

$$b_{\delta_t} = b_{\delta_1} + b_{\delta_2} + \max(0, b - t) \quad (90)$$

$$= b_{\delta_1} + b_{\delta_2} + \max(0, b_1 - t) \quad \text{from (60)} \quad (91)$$

$$= b_{\delta_1} + b_{\delta_2} \quad \text{from (90)} \quad (92)$$

$$b_{\delta_t} + t = b_{\delta_1} + b_{\delta_2} + t \quad (93)$$

Next, from the definition of a

$$a = b_{\delta_1(t)} + t \quad (94)$$

$$= \max(0, b_1 - t) + b_{\delta_1} + t \quad (95)$$

$$= b_{\delta_1} + t \quad \text{from (90)} \quad (96)$$

Now from Case (b)

$$b_1 + \leq t \quad (98)$$

$$b_1 + b_{\delta_1} \leq t + b_{\delta_1} \quad (99)$$

$$b_1 + b_{\delta_1} \leq a \quad \text{from (97)} \quad (100)$$

$$b_2 \leq a \quad \text{from (65), by transitivity} \quad (101)$$

$$b_2 - a \leq 0 \quad (102)$$

$$\max(0, b_2 - a) = 0 \quad (103)$$

$$b_{\delta_2} + \max(0, b_2 - a) = b_{\delta_2} \quad (104)$$

$$b_{\delta_2(a)} = b_{\delta_2} \quad (105)$$

$$b_{\delta_2(a)} + a = b_{\delta_2} + a \quad (106)$$

$$a' = b_{\delta_2} + a \quad \text{from the defintiion of } a' \quad (107)$$

$$= b_{\delta_2} + b_{\delta_1} + t \quad \text{from (97)} \quad (108)$$

$$= b_{\delta_t} + t \quad \text{from (94)} \quad (109)$$

which concludes the proof for Case (2)- (b). \square

3 FINITENESS

4 COMPLEXITY OF QUERY ANSWERING

5 MINIMIZATION

6 SIZE OF COMPACT ANSWERS