

Compact Answers to Temporal Regular Path Queries (Supplementary Material)

ACM Reference Format:

. 2023. Compact Answers to Temporal Regular Path Queries (Supplementary Material). In *Proceedings of ACM Conference (Conference'17)*. ACM, New York, NY, USA, 10 pages. <https://doi.org/10.1145/nnnnnnn.nnnnnnn>

1 INTRODUCTION

This document provides detailed definitions and proofs for the article *Compact answers to Temporal Regular Path Queries*, submitted at CIKM 2023.

As opposed to the structure adopted in the article, the result here are grouped by topic (inductive representation, finiteness, complexity, etc) rather than representation ($\mathcal{U}^{[t]}$, $\mathcal{U}^{[d]}$, etc.). This allows us to emphasize which proofs differ from one representation to the other.

2 NOTATION

Let $\langle o_1, o_2, \tau, \delta, b, e \rangle \in \mathcal{U}^{[t,d],b,e}$, and let $t \in \tau$.

In the article, we defined the interval δ_t for each t as

$$\delta_t \mid b_\delta + \max(0, b - t) , e_\delta - \max(0, t - e) \rfloor_\delta$$

In this supplementary material, we will use $\delta(t)$ instead of δ_t . This notation will allow us to write $\delta_1(t)$ when several tuples are involved. Note that the time points b and e in this notation are still omitted, for conciseness, because they should be clear from the context.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

Conference'17, July 2017, Washington, DC, USA

© 2023 Association for Computing Machinery.

ACM ISBN 978-x-xxxx-xxxx-x/YY/MM...\$15.00

<https://doi.org/10.1145/nnnnnnn.nnnnnnn>

3 INDUCTIVE REPRESENTATION

Let q be a TRPQ and G a TG.

Then $\llbracket q \rrbracket_G$ is the set of answers to q over G (represented as tuples in \mathcal{U}).

In this section, we provide the full definition of the four inductive representations of $\llbracket q \rrbracket_G$ discussed in the article, in $\mathcal{U}^{[t]}$, $\mathcal{U}^{[d]}$, $\mathcal{U}^{[t,d]}$ and $\mathcal{U}^{[t,d],b,e}$ respectively, and prove that they are correct.

These representations are denoted as $\llbracket q \rrbracket_G^{[t]}$, $\llbracket q \rrbracket_G^{[d]}$, $\llbracket q \rrbracket_G^{[t,d]}$ and $\llbracket q \rrbracket_G^{[t,d],b,e}$ respectively.

3.1 Grammar

For convenience, we reproduce here the syntax of TRPQs (already provided in the article).

A TRPQ is an expression for the symbol path in the following grammar:

$$\begin{aligned} \text{path} &::= \text{test} \mid \text{axis} \mid (\text{path}/\text{path}) \mid (\text{path} + \text{path}) \mid \text{path}[m, n] \mid \text{path}[m, _] \\ \text{test} &::= \text{pred} \mid (? \text{path}) \mid \text{test} \vee \text{test} \mid \text{test} \wedge \text{test} \mid \neg \text{test} \\ \text{axis} &::= F \mid B \mid T_\delta \end{aligned}$$

with $\delta \in \text{intv}(\mathcal{T})$, $m, n \in \mathbb{N}^+$ and $m \leq n$.

3.2 In \mathcal{U}

For convenience still, we reproduce the full semantics of TRPQs, i.e. the inductive definition of the evaluation $\llbracket q \rrbracket_G$ of a query q over a graph G in \mathcal{U} (already provided in the article).

$$\begin{aligned} \llbracket \text{pred} \rrbracket_G &= \{ \langle o, o, t, 0 \rangle \mid t \in \tau \text{ for some } \tau \in \text{val}(o, \text{pred}) \} \\ \llbracket T_\delta \rrbracket_G &= \{ \langle o, o, t, d \rangle \mid o \in (N \cup E), t \in \mathcal{T}_G, d \in \delta, t + d \in \mathcal{T}_G \} \\ \llbracket F \rrbracket_G &= \{ \langle v, e, t, 0 \rangle \mid \text{src}(e) = v, t \in \mathcal{T}_G \} \cup \{ \langle e, v, t, 0 \rangle \mid \text{tgt}(e) = v, t \in \mathcal{T}_G \} \\ \llbracket B \rrbracket_G &= \{ \langle v, e, t, 0 \rangle \mid \text{tgt}(e) = v, t \in \mathcal{T}_G \} \cup \{ \langle e, v, t, 0 \rangle \mid \text{src}(e) = v, t \in \mathcal{T}_G \} \\ \llbracket ?\text{path} \rrbracket_G &= \{ \langle o, o, t, 0 \rangle \mid \langle o, o', t, t + d \rangle \in \llbracket \text{path} \rrbracket_G \text{ for some } o' \in N \cup E, d \in \mathcal{T} \} \\ \llbracket \text{test}_1 \vee \text{test}_2 \rrbracket_G &= \llbracket \text{test}_1 \rrbracket_G \cup \llbracket \text{test}_2 \rrbracket_G \\ \llbracket \text{test}_1 \wedge \text{test}_2 \rrbracket_G &= \llbracket \text{test}_1 \rrbracket_G \cap \llbracket \text{test}_2 \rrbracket_G \\ \llbracket \neg \text{test} \rrbracket_G &= (\{ \langle o, o \rangle \mid o \in N \cup E \} \times \mathcal{T}_G \times \{0\}) \setminus \llbracket \text{test} \rrbracket_G \\ \llbracket \text{path}_1 / \text{path}_2 \rrbracket_G &= \{ \langle o_1, o_3, t, d_1 + d_2 \rangle \mid \exists o_2: \langle o_1, o_2, t, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G \wedge \langle o_2, o_3, t + d_1, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G \} \\ \llbracket \text{path}_1 + \text{path}_2 \rrbracket_G &= \llbracket \text{path}_1 \rrbracket_G \cup \llbracket \text{path}_2 \rrbracket_G \\ \llbracket \text{path}[m, n] \rrbracket_G &= \bigcup_{k=m}^n \llbracket \text{path}^k \rrbracket_G \\ \llbracket \text{path}[m, _] \rrbracket_G &= \bigcup_{k \geq m} \llbracket \text{path}^k \rrbracket_G \end{aligned}$$

3.3 In $\mathcal{U}^{[t]}$

3.3.1 Definition.

The full definition of $\llbracket q \rrbracket_G^{[t]}$ is already provide in the article. We only reproduce it here for convenience.

$$\begin{aligned}
\langle pred \rangle_G^{[t]} &= \{ \langle o, o, \tau, 0 \rangle \mid o \in (N \cup E), \tau \in \text{val}(o, pred) \} \\
\langle T_\delta \rangle_G^{[t]} &= \{ \langle o, o, [t_1, t_1], t_2 - t_1 \rangle \mid o \in (N \cup E), t_1 \in \mathcal{T}_G, t_2 \in (\delta + t_1) \cap \mathcal{T}_G \} \\
\langle F \rangle_G^{[t]} &= \{ \langle v, e, \mathcal{T}_G, 0 \rangle \mid \text{src}(e) = v \} \cup \{ \langle e, v, \mathcal{T}_G, 0 \rangle \mid \text{tgt}(e) = v \} \\
\langle B \rangle_G^{[t]} &= \{ \langle v, e, \mathcal{T}_G, 0 \rangle \mid \text{tgt}(e) = v \} \cup \{ \langle e, v, \mathcal{T}_G, 0 \rangle \mid \text{src}(e) = v \} \\
\langle (?path) \rangle_G^{[t]} &= \{ \langle o_1, o_1, \tau, 0 \rangle \mid \exists o_2, d: \langle o_1, o_2, \tau, d \rangle \in \langle path \rangle_G^{[t]} \} \\
\langle test_1 \vee test_2 \rangle_G^{[t]} &= \langle test_1 \rangle_G^{[t]} \cup \langle test_2 \rangle_G^{[t]} \\
\langle test_1 \wedge test_2 \rangle_G^{[t]} &= \{ \langle o, o, \tau_1 \cap \tau_2, 0 \rangle \mid \langle o, o, \tau_1, 0 \rangle \in \langle test_1 \rangle_G^{[t]}, \langle o, o, \tau_2, 0 \rangle \in \langle test_2 \rangle_G^{[t]}, \tau_1 \cap \tau_2 \neq \emptyset \} \\
\langle \neg test \rangle_G^{[t]} &= \bigcup_{o \in N \cup E} \left\{ \langle o, o, \tau, 0 \rangle \mid \tau \in \text{compl} \left(\{ \tau' \mid \langle o, o, \tau', 0 \rangle \in \langle test \rangle_G^{[t]} \}, \mathcal{T}_G \right) \right\} \\
\langle path_1 / path_2 \rangle_G^{[t]} &= \left\{ \langle o_1, o_3, ((\tau_1 + d_1) \cap \tau_2) - d_1, d_1 + d_2 \rangle \mid \exists o_2: \langle o_1, o_2, \tau_1, d_1 \rangle \in \langle path_1 \rangle_G^{[t]} \wedge \langle o_2, o_3, \tau_2, d_2 \rangle \in \langle path_2 \rangle_G^{[t]} \wedge (\tau_1 + d_1) \cap \tau_2 \neq \emptyset \right\} \\
\langle path_1 + path_2 \rangle_G^{[t]} &= \langle path_1 \rangle_G^{[t]} \cup \langle path_2 \rangle_G^{[t]} \\
\langle path[m, n] \rangle_G^{[t]} &= \bigcup_{k=m}^n \langle path^k \rangle_G^{[t]} \\
\langle path[m, _] \rangle_G^{[t]} &= \bigcup_{k \geq m} \langle path^k \rangle_G^{[t]}
\end{aligned}$$

We observe that for the operators $(path_1 + path_2)$, $(path[m, _])$ and $(path[m, n])$, the definition of $\langle q \rangle_G^{[t]}$ is nearly identical to the one of $\llbracket q \rrbracket_G$. It will also be the case for the three representations below.

3.3.2 Correctness. The following result states that this representation is correct:

PROPOSITION 3.1. *Let $G = \langle N, E, \text{conn}, \mathcal{T}_G, \text{val} \rangle$ be a TPG and q a TRPQ. Then the unfolding of $\langle q \rangle_G^{[t]}$ is $\llbracket q \rrbracket_G$.*

PROOF. Let $G = \langle \Omega, N, E, \text{conn}, \mathcal{T}_G, \text{val} \rangle$ be a TG, and let q be a TRPQ.

We show below that:

- (I) for any $\langle o_1, o_2, t, d \rangle \in \llbracket q \rrbracket_G$, there is a $\tau \in \text{intv}(\mathcal{T})$ such that
 - (a) $\langle o_1, o_2, \tau, d \rangle \in \langle q \rangle_G^{[t]}$, and
 - (b) $t \in \tau$
- (II) for any $\langle o_1, o_2, \tau, d \rangle \in \langle q \rangle_G^{[t]}$ for any $t \in \tau$,
 - $\langle o_1, o_2, t, d \rangle \in \llbracket q \rrbracket_G$

By induction on the structure of q :

$q = F$

From the above definitions, we have:

$$\begin{aligned}
\llbracket F \rrbracket_G &= \{ \langle v, e, t, 0 \rangle \mid \text{src}(e) = v, t \in \mathcal{T}_G \} \cup \{ \langle e, v, t, 0 \rangle \mid \text{tgt}(e) = v, t \in \mathcal{T}_G \} \\
\langle F \rangle_G^{[t]} &= \{ \langle v, e, \mathcal{T}_G, 0 \rangle \mid \text{src}(e) = v \} \cup \{ \langle e, v, \mathcal{T}_G, 0 \rangle \mid \text{tgt}(e) = v \}
\end{aligned}$$

- For (I), let $\mathbf{u} = \langle o_1, o_2, t, d \rangle \in \llbracket F \rrbracket_G$.

Then $d = 0$.

And since $\mathbf{u} \in \llbracket F \rrbracket_G$, $\text{src}(o_2) = o_1$ or $\text{tgt}(o_1) = o_2$ must hold.

Therefore $\langle o_1, o_2, \mathcal{T}_G, 0 \rangle \in \langle F \rangle_G^{[t]}$.

So (Ia) is verified.

An trivially, $t \in \mathcal{T}_G$, so (Ib) is verified as well.

- For (II), let $\mathbf{v} = \langle o_1, o_2, \mathcal{T}_G, 0 \rangle \in \langle q \rangle_G^{[t]}$, and let $t \in \mathcal{T}_G$.

Because $\mathbf{v} \in \langle q \rangle_G^{[t]}$, $\text{src}(o_2) = o_1$ or $\text{tgt}(o_1) = o_2$ must hold.

Therefore $\langle o_1, o_2, t, t \rangle = \langle o_1, o_2, t, t + d \rangle \in \llbracket q \rrbracket_G$.

$q = B$

The proof is almost identical to the case $q = F$.

$q = T_\delta$

From the above definitions, we have:

$$\begin{aligned}
\llbracket T_\delta \rrbracket_G &= \{ \langle o, o, t, d \rangle \mid o \in (N \cup E), t \in \mathcal{T}_G, d \in \delta, t + d \in \mathcal{T}_G \} \\
\langle T_\delta \rangle_G^{[t]} &= \{ \langle o, o, [t_1, t_1], t_2 - t_1 \rangle \mid o \in (N \cup E), t_1 \in \mathcal{T}_G, t_2 \in (\delta + t_1) \cap \mathcal{T}_G \}
\end{aligned}$$

- For (I), let $\mathbf{u} = \langle o, o, t, t + d \rangle \in \llbracket q \rrbracket_G$.
Now let $\mathbf{v} = \langle o, o, [t, t], d \rangle$ in $\mathcal{U}^{[t]}$.
For (Ia) we show that $\mathbf{v} \in \langle T_\delta \rangle_G^{[t]}$.
Since $\mathbf{u} \in \llbracket q \rrbracket_G$, we have $o \in N \cup E$ and $t \in \mathcal{T}_G$.
Besides, because $\mathbf{u} \in \llbracket q \rrbracket_G$ still, $d \in \delta$ and $t + d \in \mathcal{T}_G$.
Therefore $t + d \in t + \delta \cap \mathcal{T}_G$.
So there is a t_2 (namely $t + d$) such that $d = t_2 - t$ and $t_2 \in t + \delta \cap \mathcal{T}_G$, which concludes the proof for (Ia).
An trivially, $t \in [t, t]$ so (Ib) is verified as well.
- For (II), let $\mathbf{v} = \langle o, o, [t, t], d \rangle \in \langle q \rangle_G^{[t]}$.
Because $\mathbf{v} \in \langle q \rangle_G^{[t]}$, we have $o \in N \cup E$ and $t \in \mathcal{T}_G$.
So to conclude the proof, it is sufficient to show that (i) $d \in \delta$ and (ii) $t + d \in \mathcal{T}_G$.
Because $\mathbf{v} \in \langle q \rangle_G^{[t]}$ still, we have

$$d = t_2 - t \text{ for some } t_2 \in (\delta + t) \cap \mathcal{T}_G \quad (1)$$

From (1), $t_2 = t + d$.

Therefore from (1) still,

$$t + d \in (\delta + t) \cap \mathcal{T}_G \quad (2)$$

which proves (ii).

And from (2), we also get

$$\begin{aligned} d + t &\in \delta + t \\ d + t - t &\in \delta + t - t \\ d &\in \delta \end{aligned}$$

which proves (i).

□

3.4 In $\mathcal{U}^{[d]}$

3.4.1 Definition.

We start with the case where q is an expression for the symbol test in the grammar of Section 3.1.

The definitions of $\langle pred \rangle_G^{[t]}$ and $\langle \neg \text{test} \rangle_G^{[t]}$ are already provided in the article, we reproduce them here for completeness:

$$\begin{aligned} \langle pred \rangle_G^{[d]} &= \{ \langle o, o, t, [0, 0] \rangle \mid \tau \in \text{val}(o, pred) \text{ and } t \in \tau \} \\ \langle F \rangle_G^{[d]} &= \{ \langle v, e, t, [0, 0] \rangle \mid \text{src}(e) = v, t \in \mathcal{T}_G \} \cup \{ \langle e, v, t, [0, 0] \rangle \mid \text{tgt}(e) = v, t \in \mathcal{T}_G \} \\ \langle B \rangle_G^{[d]} &= \{ \langle v, e, t, [0, 0] \rangle \mid \text{tgt}(e) = v, t \in \mathcal{T}_G \} \cup \{ \langle e, v, t, [0, 0] \rangle \mid \text{src}(e) = v, t \in \mathcal{T}_G \} \\ \langle (? \text{path}) \rangle_G^{[d]} &= \{ \langle o_1, o_1, t, [0, 0] \rangle \mid \exists o_2, \delta: \langle o_1, o_2, t, \delta \rangle \in \langle \text{path} \rangle_G^{[d]} \} \\ \langle \text{test}_1 \vee \text{test}_2 \rangle_G^{[d]} &= \langle \text{test}_1 \rangle_G^{[d]} \cup \langle \text{test}_2 \rangle_G^{[d]} \\ \langle \text{test}_1 \wedge \text{test}_2 \rangle_G^{[d]} &= \langle \text{test}_1 \rangle_G^{[d]} \cap \langle \text{test}_2 \rangle_G^{[d]} \\ \langle \neg \text{test} \rangle_G^{[d]} &= \left\{ \langle o, o, t, [0, 0] \rangle \mid o \in N \cup E, t \in \mathcal{T}_G \setminus \{ t' \mid \langle o, o, t', [0, 0] \rangle \in \langle \text{test} \rangle_G^{[d]} \} \right\} \end{aligned}$$

Next, we consider the operators $(\text{path}_1 + \text{path}_2)$, $(\text{path}[m, _])$ and $(\text{path}[m, n])$.

For these cases, $\langle q \rangle_G^{[t, d]}$ is once again defined analogously to $\llbracket q \rrbracket_G$, in terms of temporal join (a.k.a. $\text{path}_1/\text{path}_2$) and set union.

We only write the definitions here for the sake of completeness:

$$\begin{aligned} \langle \text{path}_1 + \text{path}_2 \rangle_G^{[t, d]} &= \langle \text{path}_1 \rangle_G^{[t, d]} \cup \langle \text{path}_2 \rangle_G^{[t, d]} \\ \llbracket \text{path}[m, n] \rrbracket_G &= \bigcup_{k=m}^n \langle \text{path}^k \rangle_G^{[t, d]} \\ \llbracket \text{path}[m, _] \rrbracket_G &= \bigcup_{k \geq m} \langle \text{path}^k \rangle_G^{[t, d]} \end{aligned}$$

The only remaining operators are temporal join $(\text{path}_1/\text{path}_2)$ and temporal navigation (T_δ) , already defined in the article.

We reproduce here these two definition for convenience:

$$\begin{aligned} \langle \text{path}_1/\text{path}_2 \rangle_G^{[d]} &= \left\{ \langle o_1, o_3, t_1, \delta_2 + t_2 - t_1 \rangle \mid \exists o_2: \langle o_1, o_2, t_1, \delta_1 \rangle \in \langle \text{path}_1 \rangle_G^{[d]} \wedge \langle o_2, o_3, t_2, \delta_2 \rangle \in \langle \text{path}_2 \rangle_G^{[d]} \wedge t_2 \in t_1 + \delta_1 \right\} \\ \langle T_\delta \rangle_G^{[d]} &= \{ \langle o, o, t, (\delta + t) \cap \mathcal{T}_G \rangle \mid o \in N \cup E, t \in \mathcal{T}_G \} \end{aligned}$$

We also reproduce the alternative characterization of $\langle T_\delta \rangle_G^{[d]}$ provided in the article, as a unary operator:

$$\langle q/T_\delta \rangle_G^{[d]} = \{ \langle o_1, o_2, t, (\delta' + \delta) \cap \mathcal{T}_G \rangle \mid \langle o_1, o_2, t, \delta' \rangle \in \langle q \rangle_G^{[d]} \}$$

3.5 In $\mathcal{U}^{[t,d]}$

3.5.1 Definition.

If q is an expression for the symbol test in the grammar of Section 3.1, then all tuples in $\llbracket q \rrbracket_G$ must have distance 0. As a result, $\langle q \rangle_G^{[t,d]}$ can be trivially defined out of $\langle q \rangle_G^{[t]}$ by replacing the distance 0 with the interval $[0, 0]$, i.e.

$$\langle \text{test} \rangle_G^{[t,d]} = \{ \langle o, o, \tau, [0, 0] \rangle \mid \langle o, o, \tau, 0 \rangle \in \langle \text{test} \rangle_G^{[t]} \}$$

Again, for the operators $(\text{path}_1 + \text{path}_2)$, $(\text{path}[m, _])$ and $(\text{path}[m, n])$, the definition of $\langle q \rangle_G^{[t,d]}$ is nearly identical to the one of $\llbracket q \rrbracket_G$:

$$\begin{aligned} \langle \text{path}_1 + \text{path}_2 \rangle_G^{[t,d]} &= \langle \text{path}_1 \rangle_G^{[t,d]} \cup \langle \text{path}_2 \rangle_G^{[t,d]} \\ \llbracket \text{path}[m, n] \rrbracket_G &= \bigcup_{k=m}^n \langle \text{path}^k \rangle_G^{[t,d]} \\ \llbracket \text{path}[m, _] \rrbracket_G &= \bigcup_{k \geq m} \langle \text{path}^k \rangle_G^{[t,d]} \end{aligned}$$

The only remaining operators are temporal join $(\text{path}_1/\text{path}_2)$ and temporal navigation (T_δ) , already defined in the article, and reproduced here for convenience:

$$\begin{aligned} \langle \text{path}_1/\text{path}_2 \rangle_G^{[t,d]} &= \bigcup \{ \mathbf{u}_1 \bowtie \mathbf{u}_2 \mid \mathbf{u}_1 \in \langle \text{path}_1 \rangle_G^{[t,d]}, \mathbf{u}_2 \in \langle \text{path}_2 \rangle_G^{[t,d]} \} \\ \langle T_\delta \rangle_G^{[t,d]} &= \bigcup_{o \in N \cup E} \{ \langle o, o, \mathcal{T}_G, \delta \rangle \bowtie \langle o, o, \mathcal{T}_G, [0, 0] \rangle \} \end{aligned}$$

where $\mathbf{u}_1 \bowtie \mathbf{u}_2$ is defined as follows.

Let $\mathbf{u}_1 = \langle o_1, o_2, \tau_1, \delta_1 \rangle$ and $\mathbf{u}_2 = \langle o_3, o_4, \tau_2, \delta_2 \rangle$.

If $o_2 \neq o_3$, then $\mathbf{u}_1 \bowtie \mathbf{u}_2 = \emptyset$.

Otherwise, each tuple in $\mathbf{u}_1 \bowtie \mathbf{u}_2$ is of the form $\langle o_1, o_4, \tau, \delta \rangle$ for some τ and δ .

For $i \in \{1, 2\}$, let R_i be the binary relation over \mathcal{T} specified by the time points and distances in \mathbf{u}_i , i.e., $R_i = \{ (t, t + d) \mid t \in \tau_i, d \in \delta_i \}$.

And let $R_1 \bowtie R_2$ denote $\{ (t_1, t_3) \mid (t_1, t_2) \in R_1 \text{ and } (t_2, t_3) \in R_2 \text{ for some } t_2 \}$. Then the intervals in the set $\mathbf{u}_1 \bowtie \mathbf{u}_2$ should intuitively represent this relation $R_1 \bowtie R_2$.

For each time point $t \in \text{dom}(R_1 \bowtie R_2)$, let $\delta(t)$ denote the maximal interval s.t. $(\delta(t) + t) \subseteq \tau_2$.

We define $\mathbf{u}_1 \bowtie \mathbf{u}_2$ as

$$\{ \langle o_1, o_4, [t, t], \delta(t) + \delta_2 \rangle \mid t \in \text{dom}(R_1 \bowtie R_2) \}$$

3.6 In $\mathcal{U}^{[t,d],b,e}$

3.6.1 Definition.

If q is an expression for the symbol test in the grammar of Section 3.1, then the definition of $\langle q \rangle_G^{[t,d],b,e}$ is nearly identical to the one of $\langle q \rangle_G^{[t,d]}$, extending each tuple $\{ \langle o, o, \tau, [0, 0] \rangle \}$ with b_τ and e_τ , i.e.

$$\langle \text{test} \rangle_G^{[t,d],b,e} = \{ \langle o, o, \tau, [0, 0], b_\tau, e_\tau \rangle \mid \langle o, o, \tau, [0, 0] \rangle \in \langle \text{test} \rangle_G^{[t,d]} \}$$

Next, for the operators $(\text{path}_1 + \text{path}_2)$, $(\text{path}[m, _])$ and $(\text{path}[m, n])$, the definition of $\langle q \rangle_G^{[t,d]}$ is once again nearly identical to the one of $\llbracket q \rrbracket_G$:

$$\begin{aligned} \langle \text{path}_1 + \text{path}_2 \rangle_G^{[t,d],b,e} &= \langle \text{path}_1 \rangle_G^{[t,d],b,e} \cup \langle \text{path}_2 \rangle_G^{[t,d],b,e} \\ \llbracket \text{path}[m, n] \rrbracket_G &= \bigcup_{k=m}^n \langle \text{path}^k \rangle_G^{[t,d],b,e} \\ \llbracket \text{path}[m, _] \rrbracket_G &= \bigcup_{k \geq m} \langle \text{path}^k \rangle_G^{[t,d],b,e} \end{aligned}$$

So the only remaining operator are temporal join $(\text{path}_1/\text{path}_2)$ and temporal navigation (T_δ) , already defined in the article. We reproduce here these two definition for convenience:

$$\begin{aligned} \langle \text{path}_1/\text{path}_2 \rangle_G^{[t,d],b,e} &= \{ \mathbf{u}_1 \bowtie \mathbf{u}_2 \mid \mathbf{u}_1 \in \langle \text{path}_1 \rangle_G^{[t,d],b,e}, \mathbf{u}_2 \in \langle \text{path}_2 \rangle_G^{[t,d],b,e}, \mathbf{u}_1 \sim \mathbf{u}_2 \} \\ \langle T_\delta \rangle_G^{[t,d],b,e} &= \{ \langle o, o, \mathcal{T}_G, \delta, b_{\mathcal{T}_G}, e_{\mathcal{T}_G} \rangle \bowtie \langle o, o, \mathcal{T}_G, [0, 0], b_{\mathcal{T}_G}, e_{\mathcal{T}_G} \rangle \mid o \in N \cup E \} \end{aligned}$$

where $\mathbf{u}_1 \sim \mathbf{u}_2$ and $\mathbf{u}_1 \bowtie \mathbf{u}_2$ are defined by:

$$\begin{aligned} \langle o_1, o_2, \tau_1, \delta_1, b_1, e_1 \rangle &\sim \langle o_3, o_4, \tau_2, \delta_2, e_2, b_2 \rangle \text{ iff } o_2 = o_3 \\ \langle o_1, o_3, \tau_1, \delta_1, b_1, e_1 \rangle &\bowtie \langle o_3, o_4, \tau_2, \delta_2, e_2, b_2 \rangle = \langle o_1, o_4, \tau, \delta_1 + \delta_2, b, e \rangle \\ &\text{with} \\ \tau &= (((\tau_1 + \delta_1) \cap \tau_2) \ominus \delta_1) \cap \tau_1 \\ b &= \max(b_1, b_2 - b_{\delta_1}) \\ e &= \min(e_1, e_2 - e_{\delta_1}) \end{aligned}$$

3.6.2 Correctness.

Operator $\text{path}_1/\text{path}_2$.

Let $\mathbf{u}_1 = \langle o_1, o_2, \tau_1, \delta_1 \rangle$ and $\mathbf{u}_2 = \langle o_3, o_4, \tau_2, \delta_2 \rangle$, with $o_2 = o_3$.

Let also $\mathbf{u}_1 \bowtie \mathbf{u}_2 = \langle o_1, o_4, \tau'_1, \delta_1 + \delta_2, b, e \rangle$, with

$$\begin{aligned} \tau &= (((\tau_1 + \delta_1) \cap \tau_2) \ominus \delta_1) \cap \tau_1 \\ b &= \max(b_1, b_2 - b_{\delta_1}) \\ e &= \min(e_1, e_2 - e_{\delta_1}) \end{aligned}$$

As explained in Section 2, for $i \in \{1, 2\}$ and $t \in \tau_i$, we use $\delta_i(t)$ for the interval

$$\delta_i \downarrow b_{\delta_i} + \max(0, b_i - t), e_{\delta_i} - \max(0, t - e_i) \downarrow_{\delta_i}$$

And similarly to what we did for $\mathcal{U}^{[t,d]}$, we use R_i for be the binary relation over \mathcal{T} specified by the time points and distances in \mathbf{u}_i , i.e. $R_i = \{(t, t + d) \mid t \in \tau_i, d \in \delta_i(t)\}$.

Then the intervals in the set $\mathbf{u}_1 \bowtie \mathbf{u}_2$ should intuitively represent this relation $R_1 \bowtie R_2$, i.e. we need to prove that (i) $\tau = \text{dom}(R_1 \bowtie R_2)$ and that (ii) for each $t \in \tau$,

$$t + \delta(t) = \{t' \mid (t, t') \in R_1 \bowtie R_2\}$$

For (ii), let $t \in \tau$. We use

- a for the least value s.t. $(t, a) \in \text{range}(R_1) \cap \text{dom}(R_2)$, and
- a' for the least value s.t. $(a, a') \in R_2$

Then a' is also the least value s.t. $(t, a') \in R_1 \bowtie R_2$.

Analogously, we use z for the greatest value s.t. $(t, z) \in \text{range}(R_1) \cap \text{dom}(R_2)$, and z' for the greatest value s.t. $(z, z') \in R_2$. Then z' is also the greatest value s.t. $(t, z') \in R_1 \bowtie R_2$.

From Lemma ??:

- $\{t\} \times (t + [a, z]) \subseteq R_1$, and
- $[a, b] \times [a', z'] \subseteq R_2$

Therefore $[a', z'] = \{c \mid (t, c) \in R_1 \bowtie R_2\}$.

To conclude the proof, we show that $t + \delta_t = [a, z]$.

We only prove that $t + b_{\delta_t} = a$ (the proof that $t + e_{\delta_t} = z$ is symmetric).

Following the definition of b , we consider 2 cases:

- (1) $b_1 < b_2 - b_{\delta_1}$
- (2) $b_1 \geq b_2 - b_{\delta_1}$

For Case (1), we get

$$b_1 < b_2 - b_{\delta_1} \tag{3}$$

$$\max(b_1, b_2 - b_{\delta_1}) = b_2 - b_{\delta_1} \tag{4}$$

$$b = b_2 - b_{\delta_1} \tag{5}$$

from the definition of b

And in Case (1) still, we get:

$$b_1 < b_2 - b_{\delta_1} \tag{6}$$

$$0 < b_2 - b_{\delta_1} - b_1 \tag{7}$$

$$\max(0, b_2 - b_{\delta_1} - b_1) = b_2 - b_{\delta_1} - b_1 \tag{8}$$

Next, we consider two subcases:

- (i) $t < b_2 - b_{\delta_1}$
- (ii) $t \geq b_2 - b_{\delta_1}$

In Case (i), we get

$$t < b_2 - b_{\delta_1} \quad (9)$$

$$0 < b_2 - b_{\delta_1} - t \quad (10)$$

$$\max(0, b_2 - b_{\delta_1} - t) = b_2 - b_{\delta_1} - t \quad (11)$$

Now from the definition of δ_t ,

$$b_{\delta_t} = b_{\delta_1} + b_{\delta_2} + \max(0, b - t) \quad (12)$$

$$= b_{\delta_1} + b_{\delta_2} + \max(0, b_2 - b_{\delta_1} - t) \quad \text{from (5)} \quad (13)$$

$$= b_{\delta_1} + b_{\delta_2} + b_2 - b_{\delta_1} - t \quad \text{from (11)} \quad (14)$$

$$= b_{\delta_2} + b_2 - t \quad (15)$$

$$b_{\delta_t} + t = b_{\delta_2} + b_2 - t + t \quad (16)$$

$$= b_{\delta_2} + b_2 \quad (17)$$

Next, from the definition of a'

$$a' = b_{\delta_2(a)} + a \quad (18)$$

$$= b_{\delta_2} + \max(0, b_2 - a) + a \quad (19)$$

And, from the definition of a

$$a = b_{\delta_1(t)} + t \quad (20)$$

$$= b_{\delta_1} + \max(0, b_1 - t) + t \quad (21)$$

Then we have two further subcases:

(I) $t \geq b_1$, or

(II) $t < b_1$

In case (I):

$$t \geq b_1 \quad (22)$$

$$0 \geq b_1 - t \quad (23)$$

$$\max(0, b_1 - t) = 0 \quad (24)$$

$$a = b_{\delta_1} + t \quad \text{from (21)} \quad (25)$$

$$\max(0, b_2 - a) = \max(0, b_2 - b_{\delta_1} - t) \quad (26)$$

$$= b_2 - b_{\delta_1} - t \quad \text{from (11)} \quad (27)$$

$$= b_2 - a \quad \text{from (25)} \quad (28)$$

In case (II):

$$t < b_1 \quad (29)$$

$$0 < b_1 - t \quad (30)$$

$$\max(0, b_1 - t) = b_1 - t \quad (31)$$

$$a = b_{\delta_1} + b_1 - t + t \quad \text{from (21)} \quad (32)$$

$$= b_{\delta_1} + b_1 \quad (33)$$

$$\max(0, b_2 - a) = \max(0, b_2 - b_{\delta_1} - b_1) \quad (34)$$

$$= b_2 - b_{\delta_1} - b_1 \quad \text{from (8)} \quad (35)$$

$$= b_2 - a \quad \text{from (33)} \quad (36)$$

$$(37)$$

So in both cases (I) and (II), we get

$$\max(0, b_2 - a) = b_2 - a$$

Therefore from (19)

$$a' = b_{\delta_2} + b_2 - a + a \quad (38)$$

$$= b_{\delta_2} + b_2 \quad (39)$$

$$= t + b_{\delta_t} \quad \text{from (17)} \quad (40)$$

which concludes the proof for Case (1)- (i).

We continue with Case (1)- (ii).

From Case (ii):

$$t \geq b_2 - b_{\delta_1} \quad (41)$$

$$0 \geq b_2 - b_{\delta_1} - t \quad (42)$$

$$\max(0, b_2 - b_{\delta_1} - t) = 0 \quad (43)$$

Now from the definition of δ_t :

$$b_{\delta_t} = b_{\delta_1} + b_{\delta_2} + \max(0, b - t) \quad (44)$$

$$= b_{\delta_1} + b_{\delta_2} + \max(0, b_2 - b_1 - t) \quad \text{from (5)} \quad (45)$$

$$= b_{\delta_1} + b_{\delta_2} \quad \text{from (43)} \quad (46)$$

$$b_{\delta_t} + t = b_{\delta_1} + b_{\delta_2} + t \quad (47)$$

Next, from Case (1) and Case (ii), by transitivity, we get

$$b_1 \leq t \quad (48)$$

$$\max(0, b_1 - t) = 0 \quad (49)$$

And from the definition of a

$$a = b_{\delta_1(t)} + t \quad (50)$$

$$= b_{\delta_1} + \max(0, b_1 - t) + t \quad (51)$$

$$= b_{\delta_1} + t \quad \text{from (49)} \quad (52)$$

$$\geq b_{\delta_1} + b_2 - b_{\delta_1} \quad \text{from Case (ii)} \quad (53)$$

$$\geq b_2 \quad (54)$$

$$0 \geq b_2 - a \quad (55)$$

$$\max(0, b_2 - a) = 0 \quad (56)$$

Therefore from (19) and (56)

$$a' = b_{\delta_2} + a \quad (57)$$

$$= b_{\delta_2} + b_{\delta_1} + t \quad \text{from (52)} \quad (58)$$

$$= b_{\delta_t} + t \quad \text{from (17)} \quad (59)$$

which concludes the proof for Case (1)- (ii).

We continue with Case (2).

In this case, we get

$$b_1 \geq b_2 - b_{\delta_1} \quad (60)$$

$$\max(b_1, b_2 - b_{\delta_1}) = b_1 \quad (61)$$

$$b = b_1 \quad \text{from the definition of } b \quad (62)$$

And from Case (2) still, we derive

$$b_1 \geq b_2 - b_{\delta_1} \quad (63)$$

$$0 \geq b_2 - b_{\delta_1} - b_1 \quad (64)$$

$$\max(0, -b_{\delta_1} - b_1) = 0 \quad (65)$$

As well as

$$b_1 \geq b_2 - b_{\delta_1} \quad (66)$$

$$b_1 + b_{\delta_1} \geq b_2 \quad (67)$$

Next, we distinguish two subcases, namely

(a) $t < b_1$ and

(b) $t \geq b_1$

We start with Case (a).

In this case,

$$t < b_1 \quad (68)$$

$$0 < b_1 - t \quad (69)$$

$$\max(0, b_1 - t) = b_1 - t \quad (70)$$

And from the definition of δ_t :

$$b_{\delta_t} = b_{\delta_1} + b_{\delta_2} + \max(0, b - t) \quad (71)$$

$$= b_{\delta_1} + b_{\delta_2} + \max(0, b_1 - t) \quad \text{from (62)} \quad (72)$$

$$= b_{\delta_1} + b_{\delta_2} + b_1 - t \quad \text{from (70)} \quad (73)$$

$$b_{\delta_t} + t = b_{\delta_1} + b_{\delta_2} + b_1 - t + t \quad (74)$$

$$= b_{\delta_1} + b_{\delta_2} + b_1 \quad (75)$$

Next, from the definition of a

$$a = b_{\delta_1(t)} + t \quad (76)$$

$$= \max(0, b_1 - t) + b_{\delta_1} + t \quad (77)$$

$$= b_1 - t + b_{\delta_1} + t \quad \text{from (70)} \quad (78)$$

$$= b_1 + b_{\delta_1} \quad (79)$$

So from (67)

$$a \geq b_2 \quad (80)$$

$$0 \geq b_2 - a \quad (81)$$

$$\max(0, b_2 - a) = 0 \quad (82)$$

$$b_{\delta_2} + \max(0, b_2 - a) = b_{\delta_2} \quad (83)$$

$$b_{\delta_2(a)} = b_{\delta_2} \quad (84)$$

$$b_{\delta_2(a)} + a = b_{\delta_2} + a \quad (85)$$

$$a' = b_{\delta_2} + a \quad \text{from the definition of } a' \quad (86)$$

$$a' = b_{\delta_2} + b_1 + b_{\delta_1} \quad \text{from (79)} \quad (87)$$

$$a' = b_{\delta_t} + t \quad \text{from (75)} \quad (88)$$

which concludes the proof for Case (2)- (a).

We end with Case (2)- (b). In this case,

$$t \geq b_1 \quad (89)$$

$$0 \geq b_1 - t \quad (90)$$

$$\max(0, b_1 - t) = 0 \quad (91)$$

And from the definition of δ_t :

$$b_{\delta_t} = b_{\delta_1} + b_{\delta_2} + \max(0, b - t) \quad (92)$$

$$= b_{\delta_1} + b_{\delta_2} + \max(0, b_1 - t) \quad \text{from (62)} \quad (93)$$

$$= b_{\delta_1} + b_{\delta_2} \quad \text{from (92)} \quad (94)$$

$$b_{\delta_t} + t = b_{\delta_1} + b_{\delta_2} + t \quad (95)$$

Next, from the definition of a

$$a = b_{\delta_1(t)} + t \quad (96)$$

$$= \max(0, b_1 - t) + b_{\delta_1} + t \quad (97)$$

$$= b_{\delta_1} + t \quad \text{from (92)} \quad (98)$$

Now from Case (b)

$$b_1 + \leq t \quad (100)$$

$$b_1 + b_{\delta_1} \leq t + b_{\delta_1} \quad (101)$$

$$b_1 + b_{\delta_1} \leq a \quad \text{from (99)} \quad (102)$$

$$b_2 \leq a \quad \text{from (67), by transitivity} \quad (103)$$

$$b_2 - a \leq 0 \quad (104)$$

$$\max(0, b_2 - a) = 0 \quad (105)$$

$$b_{\delta_2} + \max(0, b_2 - a) = b_{\delta_2} \quad (106)$$

$$b_{\delta_2(a)} = b_{\delta_2} \quad (107)$$

$$b_{\delta_2(a)} + a = b_{\delta_2} + a \quad (108)$$

$$a' = b_{\delta_2} + a \quad \text{from the definition of } a' \quad (109)$$

$$= b_{\delta_2} + b_{\delta_1} + t \quad \text{from (99)} \quad (110)$$

$$= b_{\delta_t} + t \quad \text{from (96)} \quad (111)$$

which concludes the proof for Case (2)- (b). \square

4 FINITENESS

5 COMPLEXITY OF QUERY ANSWERING

5.1 Problem

We define in this section a decision problem for each representation, similar to the problem $\text{COMPACT ANSWER}^{[t]}$ defined in the article. First, we define a (possibly partial) order over tuples of each representation, in the expected way, i.e.:

- over $\mathcal{U}^{[t]}$: $\langle o_1, o_2, \tau_1, d_1 \rangle \sqsubseteq_{[t]} \langle o_3, o_4, \tau_2, d_2 \rangle$ iff $\langle o_1, o_2, d_1 \rangle = \langle o_3, o_4, d_2 \rangle$ and $\tau_1 \subseteq \tau_2$,
- over $\mathcal{U}^{[d]}$: $\langle o_1, o_2, t_1, \delta_1 \rangle \sqsubseteq_{[d]} \langle o_3, o_4, t_2, \delta_2 \rangle$ iff $\langle o_1, o_2, t_1 \rangle = \langle o_3, o_4, t_2 \rangle$ and $\delta_1 \subseteq \delta_2$,
- over $\mathcal{U}^{[t,d]}$: $\langle o_1, o_2, \tau_1, \delta_1 \rangle \sqsubseteq_{[t,d]} \langle o_3, o_4, \tau_2, \delta_2 \rangle$ iff $\langle o_1, o_2 \rangle = \langle o_3, o_4 \rangle$, $\tau_1 \subseteq \tau_2$ and $\delta_1 \subseteq \delta_2$,
- over $\mathcal{U}^{[t,d],b,e}$: $\langle o_1, o_2, \tau_1, \delta_1, b_1, b_2 \rangle \sqsubseteq_{[t,d],b,e} \langle o_3, o_4, \tau_2, \delta_2, b_2, e_2 \rangle$ iff $\langle o_1, o_2 \rangle = \langle o_3, o_4 \rangle$, $\tau_1 \subseteq \tau_2$ and $\delta_1(t) \subseteq \delta_2(t)$ for all $t \in \tau_1 \cap \tau_2$ (the notation of $\delta_i(t)$ is explained above, in Section 2).

Now let x be one of $[t]$, $[d]$, $[t, d]$ or $[t, d], b, e$.

We decline the notion of compact answer defined in Section XXX in four flavors, as follows:

Definition 5.1. Let G be a TG, let q be a TRPQ and let $\mathbf{u} \in \mathcal{U}^x$.

We say that \mathbf{u} is a *compact answer* to q over G (in \mathcal{U}^x) if $\mathbf{u} \in \max_{\sqsubseteq_x} \{\mathbf{u}' \in \mathcal{U}^x \mid \text{unfold}(\mathbf{u}') = \llbracket q \rrbracket_G\}$

And similary, we get four decision problems:

COMPACT ANSWER^x

Input: TG G , TRPQ q , tuple $\mathbf{u} \in \mathcal{U}^x$

Decide: \mathbf{u} is a compact answer to q over G (in \mathcal{U}^x)

5.2 Hardness

5.3 Membership

6 MINIMIZATION

7 SIZE OF COMPACT ANSWERS