

# Compact Answers to Temporal Regular Path Queries (Supplementary Material)

## ACM Reference Format:

. 2023. Compact Answers to Temporal Regular Path Queries (Supplementary Material). In *Proceedings of 32nd ACM International Conference on Information and Knowledge Management (CIKM'23)*. ACM, New York, NY, USA, 16 pages. <https://doi.org/10.1145/nnnnnnnn.nnnnnnnn>

## 1 INTRODUCTION

This document provides detailed definitions and proofs for the article *Compact answers to Temporal Regular Path Queries*, submitted at CIKM 2023.

As opposed to the structure adopted in the article, the result here are grouped by topic (inductive representation, finiteness, complexity, etc) rather than representation ( $\mathcal{U}^{[t]}$ ,  $\mathcal{U}^{[d]}$ , etc.). This allows us to emphasize which proofs differ from one representation to the other.

## 2 NOTATION

Let  $\langle o_1, o_2, \tau, \delta, b, e \rangle \in \mathcal{U}^{[t,d],b,e}$ , and let  $t \in \tau$ .

In the article, we defined the interval  $\delta_t$  for each  $t$  as

$$\delta_t [b_\delta + \max(0, b - t), e_{\delta_t} - \max(0, t - e)]_\delta$$

In this supplementary material, we will use  $\delta(t)$  instead of  $\delta_t$ . This notation will allow us to write  $\delta_1(t)$  when several tuples are involved. Note that the time points  $b$  and  $e$  in this notation are still omitted, for conciseness, because they should be clear from the context.

---

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from [permissions@acm.org](mailto:permissions@acm.org).

CIKM'23, October 21–25, 2023, Birmingham, U.K.

© 2023 Association for Computing Machinery.

ACM ISBN 978-x-xxxx-xxxx-x/YY/MM...\$15.00

<https://doi.org/10.1145/nnnnnnnn.nnnnnnnn>

### 3 INDUCTIVE REPRESENTATION

Let  $q$  be a TRPQ and  $G$  a TG.

Then  $\llbracket q \rrbracket_G$  is the set of answers to  $q$  over  $G$  (represented as tuples in  $\mathcal{U}$ ).

In this section, we provide the full definition of the four inductive representations of  $\llbracket q \rrbracket_G$  discussed in the article, in  $\mathcal{U}^{[t]}$ ,  $\mathcal{U}^{[d]}$ ,  $\mathcal{U}^{[t,d]}$  and  $\mathcal{U}^{[t,d],b,e}$  respectively, and prove that they are correct.

These representations are denoted as  $\langle q \rangle_G^{[t]}$ ,  $\langle q \rangle_G^{[d]}$ ,  $\langle q \rangle_G^{[t,d]}$  and  $\langle q \rangle_G^{[t,d],b,e}$  respectively.

#### 3.1 Grammar

For convenience, we reproduce here the syntax of TRPQs (already provided in the article).

*Definition 3.1 (TRPQ).* A TRPQ is an expression for the symbol path in the following grammar:

$$\begin{aligned} \text{path} &::= \text{test} \mid \text{axis} \mid (\text{path}/\text{path}) \mid (\text{path} + \text{path}) \mid \text{path}[m, n] \mid \text{path}[m, \_] \\ \text{test} &::= \text{pred} \mid (? \text{path}) \mid \text{test} \vee \text{test} \mid \text{test} \wedge \text{test} \mid \neg \text{test} \\ \text{axis} &::= F \mid B \mid T_\delta \end{aligned}$$

with  $\delta \in \text{intv}(\mathcal{T})$ ,  $m, n \in \mathbb{N}^+$  and  $m \leq n$ .

#### 3.2 In $\mathcal{U}$

For convenience still, we reproduce the full semantics of TRPQs, i.e. the inductive definition of the evaluation  $\llbracket q \rrbracket_G$  of a query  $q$  over a graph  $G$  in  $\mathcal{U}$  (already provided in the article).

$$\begin{aligned} \llbracket \text{pred} \rrbracket_G &= \{ \langle o, o, t, 0 \rangle \mid t \in \tau \text{ for some } \tau \in \text{val}(o, \text{pred}) \} \\ \llbracket T_\delta \rrbracket_G &= \{ \langle o, o, t, d \rangle \mid o \in (N \cup E), t \in \mathcal{T}_G, d \in \delta, t + d \in \mathcal{T}_G \} \\ \llbracket F \rrbracket_G &= \{ \langle v, e, t, 0 \rangle \mid \text{src}(e) = v, t \in \mathcal{T}_G \} \cup \{ \langle e, v, t, 0 \rangle \mid \text{tgt}(e) = v, t \in \mathcal{T}_G \} \\ \llbracket B \rrbracket_G &= \{ \langle v, e, t, 0 \rangle \mid \text{tgt}(e) = v, t \in \mathcal{T}_G \} \cup \{ \langle e, v, t, 0 \rangle \mid \text{src}(e) = v, t \in \mathcal{T}_G \} \\ \llbracket ? \text{path} \rrbracket_G &= \{ \langle o, o, t, 0 \rangle \mid \langle o, o', t, t + d \rangle \in \llbracket \text{path} \rrbracket_G \text{ for some } o' \in N \cup E, d \in \mathcal{T} \} \\ \llbracket \text{test}_1 \vee \text{test}_2 \rrbracket_G &= \llbracket \text{test}_1 \rrbracket_G \cup \llbracket \text{test}_2 \rrbracket_G \\ \llbracket \text{test}_1 \wedge \text{test}_2 \rrbracket_G &= \llbracket \text{test}_1 \rrbracket_G \cap \llbracket \text{test}_2 \rrbracket_G \\ \llbracket \neg \text{test} \rrbracket_G &= (\{ \langle o, o \rangle \mid o \in N \cup E \} \times \mathcal{T}_G \times \{0\}) \setminus \llbracket \text{test} \rrbracket_G \\ \llbracket \text{path}_1 / \text{path}_2 \rrbracket_G &= \{ \langle o_1, o_3, t, d_1 + d_2 \rangle \mid \exists o_2: \langle o_1, o_2, t, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G \wedge \langle o_2, o_3, t + d_1, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G \} \\ \llbracket \text{path}_1 + \text{path}_2 \rrbracket_G &= \llbracket \text{path}_1 \rrbracket_G \cup \llbracket \text{path}_2 \rrbracket_G \\ \llbracket \text{path}[m, n] \rrbracket_G &= \bigcup_{k=m}^n \llbracket \text{path}^k \rrbracket_G \\ \llbracket \text{path}[m, \_] \rrbracket_G &= \bigcup_{k \geq m} \llbracket \text{path}^k \rrbracket_G \end{aligned}$$

#### 3.3 In $\mathcal{U}^{[t]}$

##### 3.3.1 Definition.

The full definition of  $\langle q \rangle_G^{[t]}$  is already provided in the article. We only reproduce it here for convenience.

$$\begin{aligned} \langle \text{pred} \rangle_G^{[t]} &= \{ \langle o, o, \tau, 0 \rangle \mid o \in (N \cup E), \tau \in \text{val}(o, \text{pred}) \} \\ \langle T_\delta \rangle_G^{[t]} &= \{ \langle o, o, [t_1, t_1], t_2 - t_1 \rangle \mid o \in (N \cup E), t_1 \in \mathcal{T}_G, t_2 \in (\delta + t_1) \cap \mathcal{T}_G \} \\ \langle F \rangle_G^{[t]} &= \{ \langle v, e, \mathcal{T}_G, 0 \rangle \mid \text{src}(e) = v \} \cup \{ \langle e, v, \mathcal{T}_G, 0 \rangle \mid \text{tgt}(e) = v \} \\ \langle B \rangle_G^{[t]} &= \{ \langle v, e, \mathcal{T}_G, 0 \rangle \mid \text{tgt}(e) = v \} \cup \{ \langle e, v, \mathcal{T}_G, 0 \rangle \mid \text{src}(e) = v \} \\ \langle (? \text{path}) \rangle_G^{[t]} &= \{ \langle o, o, \tau, 0 \rangle \mid \langle o, o', \tau, d \rangle \in \langle \text{path} \rangle_G^{[t]} \text{ for some } o' \in N \cup E, d \in \mathcal{T} \} \\ \langle \text{test}_1 \vee \text{test}_2 \rangle_G^{[t]} &= \langle \text{test}_1 \rangle_G^{[t]} \cup \langle \text{test}_2 \rangle_G^{[t]} \\ \langle \text{test}_1 \wedge \text{test}_2 \rangle_G^{[t]} &= \{ \langle o, o, \tau_1 \cap \tau_2, 0 \rangle \mid \langle o, o, \tau_1, 0 \rangle \in \langle \text{test}_1 \rangle_G^{[t]}, \langle o, o, \tau_2, 0 \rangle \in \langle \text{test}_2 \rangle_G^{[t]}, \tau_1 \cap \tau_2 \neq \emptyset \} \\ \langle \neg \text{test} \rangle_G^{[t]} &= \bigcup_{o \in N \cup E} \left\{ \langle o, o, \tau, 0 \rangle \mid \tau \in \text{compl} \left( \{ \tau' \mid \langle o, o, \tau', 0 \rangle \in \langle \text{test} \rangle_G^{[t]}, \mathcal{T}_G \} \right) \right\} \\ \langle \text{path}_1 / \text{path}_2 \rangle_G^{[t]} &= \left\{ \langle o_1, o_3, ((\tau_1 + d_1) \cap \tau_2) - d_1, d_1 + d_2 \rangle \mid \exists o_2: \langle o_1, o_2, \tau_1, d_1 \rangle \in \langle \text{path}_1 \rangle_G^{[t]} \wedge \langle o_2, o_3, \tau_2, d_2 \rangle \in \langle \text{path}_2 \rangle_G^{[t]} \wedge (\tau_1 + d_1) \cap \tau_2 \neq \emptyset \right\} \\ \langle \text{path}_1 + \text{path}_2 \rangle_G^{[t]} &= \langle \text{path}_1 \rangle_G^{[t]} \cup \langle \text{path}_2 \rangle_G^{[t]} \\ \langle \text{path}[m, n] \rangle_G^{[t]} &= \bigcup_{k=m}^n \langle \text{path}^k \rangle_G^{[t]} \\ \langle \text{path}[m, \_] \rangle_G^{[t]} &= \bigcup_{k \geq m} \langle \text{path}^k \rangle_G^{[t]} \end{aligned}$$

We observe that when  $q$  is of the form  $(\text{path}_1 + \text{path}_2)$ ,  $(\text{path}[m, \_])$  and  $(\text{path}[m, n])$ , the definition of  $\llbracket q \rrbracket_G^{[t]}$  is nearly identical to the one of  $\llbracket q \rrbracket_G$ . This will also be the case for the three representations below.

### 3.3.2 Correctness.

We start with a lemma:

LEMMA 3.2. *Let  $G = \langle N, E, \text{conn}, \mathcal{T}_G, \text{val} \rangle$  be a TPG and let  $q$  an expression for the symbol test in the grammar of Definition 3.1. Then:*

- *each tuples in  $\llbracket q \rrbracket_G$  is of the form  $\langle o_1, o_2, t, 0 \rangle$  for some  $o_1, o_2$  and  $t$ ,*
- *each tuples in  $\llbracket q \rrbracket_G^{[t]}$  is of the form  $\langle o_1, o_2, \tau, 0 \rangle$  for some  $o_1, o_2$  and  $\tau$ .*

PROOF. Immediate from the definitions of  $\llbracket q \rrbracket_G$  and  $\llbracket q \rrbracket_G^{[t]}$ . □

The following result states that the representation  $\llbracket q \rrbracket_G^{[t]}$  is correct:

PROPOSITION 3.3. *Let  $G = \langle N, E, \text{conn}, \mathcal{T}_G, \text{val} \rangle$  be a TPG and  $q$  a TRPQ. Then the unfolding of  $\llbracket q \rrbracket_G^{[t]}$  is  $\llbracket q \rrbracket_G$ .*

PROOF.

Let  $G = \langle N, E, \text{conn}, \mathcal{T}_G, \text{val} \rangle$  be a TG, and let  $q$  be a TRPQ.

We show below that:

- (I) for any  $\langle o_1, o_2, t, d \rangle \in \llbracket q \rrbracket_G$ , there is a  $\tau \in \text{intv}(\mathcal{T})$  such that
  - (a)  $\langle o_1, o_2, \tau, d \rangle \in \llbracket q \rrbracket_G^{[t]}$ , and
  - (b)  $t \in \tau$ ,
- (II) for any  $\langle o_1, o_2, \tau, d \rangle \in \llbracket q \rrbracket_G^{[t]}$  for any  $t \in \tau$ ,
  - $\langle o_1, o_2, t, d \rangle$  is in  $\llbracket q \rrbracket_G$ .

We proceed by induction on the structure of  $q$ .

If  $q$  is of the form  $\text{pred}$ ,  $F$ ,  $B$ ,  $(\text{test} \vee \text{test})$ ,  $(\text{path} + \text{path})$ ,  $\text{path}[m, n]$  or  $\text{path}[m, \_]$ , then (I) and (II) immediately follow from the definitions of  $\llbracket q \rrbracket_G$  and  $\llbracket q \rrbracket_G^{[t]}$ .

So we focus below on the five remaining cases:

- $q = T_\delta$ .

From the above definitions, we have:

$$\begin{aligned} \llbracket q \rrbracket_G &= \{ \langle o, o, t, d \rangle \mid o \in (N \cup E), t \in \mathcal{T}_G, d \in \delta, t + d \in \mathcal{T}_G \} \\ \llbracket q \rrbracket_G^{[t]} &= \{ \langle o, o, [t_1, t_1], t_2 - t_1 \rangle \mid o \in (N \cup E), t_1 \in \mathcal{T}_G, t_2 \in (\delta + t_1) \cap \mathcal{T}_G \} \end{aligned}$$

- For (I), let  $\mathbf{v} = \langle o, o, t, t + d \rangle \in \llbracket q \rrbracket_G$ .

And let  $\mathbf{u} = \langle o, o, [t, t], d \rangle$  in  $\mathcal{U}^{[t]}$ .

For (Ia) we show that  $\mathbf{u} \in \llbracket q \rrbracket_G^{[t]}$ .

From  $\mathbf{v} \in \llbracket q \rrbracket_G$ , we get  $o \in N \cup E$  and  $t \in \mathcal{T}_G$ .

Besides, because  $\mathbf{v} \in \llbracket q \rrbracket_G$  still,

$$t + d \in \mathcal{T}_G \tag{1}$$

and

$$d \in \delta \tag{2}$$

$$t + d \in t + \delta \tag{3}$$

So from (1) and (3)

$$t + d \in (\delta + t) \cap \mathcal{T}_G \tag{4}$$

So there is a  $t_2$  (namely  $t + d$ ) such that  $d = t_2 - t$  and  $t_2 \in t + \delta \cap \mathcal{T}_G$ .

Together with the definition of  $\llbracket q \rrbracket_G^{[t]}$ , this implies  $\mathbf{u} \in \llbracket q \rrbracket_G^{[t]}$ , which concludes the proof for (Ia).

And trivially,  $t \in [t, t]$ , so (Ib) is verified as well.

- For (II), let  $\mathbf{u} = \langle o, o, [t, t], d \rangle \in \llbracket q \rrbracket_G^{[t]}$ .

From  $\mathbf{u} \in \llbracket q \rrbracket_G^{[t]}$ , we get  $o \in N \cup E$  and  $t \in \mathcal{T}_G$ .

So to conclude the proof, it is sufficient to show that (i)  $d \in \delta$  and (ii)  $t + d \in \mathcal{T}_G$ .

Because  $\mathbf{u} \in \llbracket q \rrbracket_G^{[t]}$  still, we have

$$d = t_2 - t \text{ for some } t_2 \in (\delta + t) \cap \mathcal{T}_G \quad (5)$$

From (5), we get  $t_2 = t + d$ .

Therefore from (5) still,

$$t + d \in (\delta + t) \cap \mathcal{T}_G \quad (6)$$

which proves (ii).

And from (6), we also get

$$\begin{aligned} t + d &\in \delta + t \\ t + d - t &\in (\delta + t) - t \\ d &\in \delta \end{aligned}$$

which proves (i).

- $q = \text{test}_1 \wedge \text{test}_2$ .

From the above definitions, we have:

$$\begin{aligned} \llbracket q \rrbracket_G &= \llbracket \text{test}_1 \rrbracket_G \cap \llbracket \text{test}_2 \rrbracket_G \\ \llbracket q \rrbracket_G^{[t]} &= \{ \langle o, o, \tau_1 \cap \tau_2, 0 \rangle \mid \langle o, o, \tau_1, 0 \rangle \in \llbracket \text{test}_1 \rrbracket_G^{[t]}, \langle o, o, \tau_2, 0 \rangle \in \llbracket \text{test}_2 \rrbracket_G^{[t]}, \tau_1 \cap \tau_2 \neq \emptyset \} \end{aligned}$$

- For (I), let  $\mathbf{v} = \langle o, o, t, d \rangle \in \llbracket q \rrbracket_G$ .

From Lemma 3.2,  $d = 0$ .

And from the definition of  $\llbracket q \rrbracket_G$ ,  $\mathbf{v} \in \llbracket \text{test}_1 \rrbracket_G \cap \llbracket \text{test}_2 \rrbracket_G$ .

So by IH, there are intervals  $\tau_1$  and  $\tau_2$  s.t.  $\langle o, o, \tau_i, 0 \rangle \in \llbracket \text{test}_i \rrbracket_G^{[t]}$  for  $i \in \{1, 2\}$  and  $t \in \tau_1 \cap \tau_2$ .

Together with the definition of  $\llbracket q \rrbracket_G^{[t]}$ , this proves (I).

- For (II), let  $\langle o, o, \tau, d \rangle \in \llbracket q \rrbracket_G^{[t]}$ .

Then from Lemma 3.2,  $d = 0$ .

And from the definition of  $\llbracket q \rrbracket_G^{[t]}$ , there are two intervals  $\tau_1$  and  $\tau_2$  s.t.  $\tau = \tau_1 \cap \tau_2$  and  $\langle o, o, \tau_i, 0 \rangle \in \llbracket \text{test}_i \rrbracket_G^{[t]}$  for  $i \in \{1, 2\}$ .

Now take any  $t \in \tau$ .

Then  $t \in \tau_i$  for  $i \in \{1, 2\}$ .

So by IH,  $\langle o, o, t, 0 \rangle \in \llbracket \text{test}_i \rrbracket_G$  for each  $i \in \{1, 2\}$ .

Together with the definition of  $\llbracket q \rrbracket_G$ , this proves (II).

- $q = (? \text{path})$ .

From the above definitions, we have:

$$\begin{aligned} \llbracket q \rrbracket_G &= \{ \langle o, o, t, 0 \rangle \mid \langle o, o', t, t + d \rangle \in \llbracket \text{path} \rrbracket_G \text{ for some } o' \in N \cup E, d \in \mathcal{T} \} \\ \llbracket q \rrbracket_G^{[t]} &= \{ \langle o, o, \tau, 0 \rangle \mid \langle o, o', \tau, d \rangle \in \llbracket \text{path} \rrbracket_G^{[t]} \text{ for some } o' \in N \cup E, d \in \mathcal{T} \} \end{aligned}$$

- For (I), let  $\langle o, o, t, 0 \rangle \in \llbracket q \rrbracket_G$ .

From the definition of  $\llbracket q \rrbracket_G$ , there are  $o'$  and  $d$  such that  $\langle o, o', t, t + d \rangle \in \llbracket \text{path} \rrbracket_G$ .

So by IH, there is a  $\tau$  s.t.  $t \in \tau$  and  $\langle o, o', \tau, d \rangle \in \llbracket \text{path} \rrbracket_G^{[t]}$ .

Therefore  $\langle o, o, \tau, 0 \rangle \in \llbracket q \rrbracket_G^{[t]}$ , from the definition of  $\llbracket q \rrbracket_G^{[t]}$ .

- For (II), let  $\langle o, o, \tau, 0 \rangle \in \llbracket q \rrbracket_G^{[t]}$ .

From the definition of  $\llbracket q \rrbracket_G^{[t]}$ , there are  $o'$  and  $d$  s.t.  $\langle o, o', \tau, d \rangle \in \llbracket \text{path} \rrbracket_G^{[t]}$ .

Now take any  $t \in \tau$ .

By IH,  $\langle o, o', t, t + d \rangle \in \llbracket \text{path} \rrbracket_G$ .

Therefore  $\langle o, o, t, 0 \rangle \in \llbracket q \rrbracket_G$ , from the definition of  $\llbracket q \rrbracket_G$ .

- $q = \neg \text{test}$ .

From the above definitions, we have:

$$\begin{aligned} \llbracket q \rrbracket_G &= (\{\langle o, o \rangle \mid o \in N \cup E\} \times \mathcal{T}_G \times \{0\}) \setminus \llbracket \text{test} \rrbracket_G \\ \langle q \rangle_G^{[t]} &= \bigcup_{o \in N \cup E} \left\{ \langle o, o, \tau, 0 \rangle \mid \tau \in \text{compl} \left( \{\tau' \mid \langle o, o, \tau', 0 \rangle \in \langle \text{test} \rangle_G^{[t]} \}, \mathcal{T}_G \right) \right\} \end{aligned}$$

- For (I), let  $\mathbf{v} = \langle o, o, t, 0 \rangle \in \llbracket q \rrbracket_G$ .  
From the definition of  $\llbracket q \rrbracket_G$ ,  $\mathbf{v} \notin \llbracket \text{test} \rrbracket_G$ .  
So

$$t \notin \{\tau' \mid \langle o, o, \tau', 0 \rangle \in \llbracket \text{test} \rrbracket_G\} \quad (7)$$

Now by IH, together with Lemma 3.2, we get:

$$\langle o, o, t, 0 \rangle \in \llbracket \text{test} \rrbracket_G \text{ iff } t' \in \tau' \text{ for some } \tau' \text{ s.t. } \langle o, o, \tau', 0 \rangle \in \langle \text{test} \rangle_G^{[t]} \quad (8)$$

So from (7) and (8):

$$t \notin \bigcup \{\tau' \mid \langle o, o, \tau', 0 \rangle \in \langle \text{test} \rangle_G^{[t]}\}$$

Therefore

$$t \in \mathcal{T}_G \setminus \bigcup \{\tau' \mid \langle o, o, \tau', 0 \rangle \in \langle \text{test} \rangle_G^{[t]}\} \quad (9)$$

So  $t \in \tau$  for some  $\tau \in \text{compl} \left( \bigcup \{\tau' \mid \langle o, o, \tau', 0 \rangle \in \langle \text{test} \rangle_G^{[t]} \}, \mathcal{T}_G \right)$ .

And  $\langle o, o, \tau, 0 \rangle \in \langle q \rangle_G^{[t]}$ , from the definition of  $\langle q \rangle_G^{[t]}$ .

- For (II), let  $\langle o, o, \tau, 0 \rangle \in \langle q \rangle_G^{[t]}$ .  
And take any  $t \in \tau$ .

From the definition of  $\langle q \rangle_G^{[t]}$ :

$$t \in \mathcal{T}_G \setminus \bigcup \{\tau' \mid \langle o, o, \tau', 0 \rangle \in \langle \text{test} \rangle_G^{[t]}\}$$

Together with (8), this implies

$$\langle o, o, t, 0 \rangle \notin \llbracket \text{test} \rrbracket_G$$

Therefore  $\langle o, o, t, 0 \rangle \in \llbracket q \rrbracket_G$ , from the definition of  $\llbracket q \rrbracket_G$ .

- $q = \text{path}_1 / \text{path}_2$ .

From the above definitions, we have:

$$\begin{aligned} \llbracket q \rrbracket_G &= \{ \langle o_1, o_3, t, d_1 + d_2 \rangle \mid \exists o_2: \langle o_1, o_2, t, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G \wedge \langle o_2, o_3, t + d_1, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G \} \\ \langle q \rangle_G^{[t]} &= \left\{ \langle o_1, o_3, ((\tau_1 + d_1) \cap \tau_2) - d_1, d_1 + d_2 \rangle \mid \exists o_2: \langle o_1, o_2, \tau_1, d_1 \rangle \in \langle \text{path}_1 \rangle_G^{[t]} \wedge \langle o_2, o_3, \tau_2, d_2 \rangle \in \langle \text{path}_2 \rangle_G^{[t]} \wedge \right. \\ &\quad \left. (\tau_1 + d_1) \cap \tau_2 \neq \emptyset \right\} \end{aligned}$$

- For (I), let  $\mathbf{v} = \langle o_1, o_3, t, d \rangle \in \llbracket q \rrbracket_G$ .  
From the definition of  $\llbracket q \rrbracket_G$ , there are  $o_2, d_1$  and  $d_2$  such that  $\langle o_1, o_2, t, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G$ ,  $\langle o_2, o_3, t + d_1, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G$  and  $d = d_1 + d_2$ .  
By IH, because  $\langle o_1, o_2, t, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G$ , there is a  $\tau_1$  such that  $t \in \tau_1$  and

$$\langle o_1, o_2, \tau_1, d_1 \rangle \in \langle \text{path}_1 \rangle_G^{[t]} \quad (10)$$

And similarly, because  $\langle o_2, o_3, t + d_1, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G$ , there is a  $\tau_2$  such that  $t + d_1 \in \tau_2$  and

$$\langle o_2, o_3, \tau_2, d_2 \rangle \in \langle \text{path}_2 \rangle_G^{[t]} \quad (11)$$

From  $t \in \tau_1$ , we get

$$t + d_1 \in \tau_1 + d_1 \quad (12)$$

Together with the fact that  $t + d_1 \in \tau_2$ , this implies

$$\tau_1 + d_1 \cap \tau_2 \neq \emptyset \quad (13)$$

So from (10), (11), (13) and the definition of  $\langle q \rangle_G^{[t]}$ ,

$$\langle o_1, o_2, ((\tau_1 + d_1) \cap \tau_2) - d_1, d_1 + d_2 \rangle \in \langle q \rangle_G^{[t]}$$

which proves (Ia).

And in order to prove (Ib), we only need to show that

$$t \in ((\tau_1 + d_1) \cap \tau_2) - d_1$$

We know that  $t \in \tau_1$ , therefore

$$t + d_1 \in \tau_1 + d_1$$

Together with the fact that  $t + d_1 \in \tau_2$ , this yields

$$\begin{aligned} t + d_1 &\in (\tau_1 + d_1) \cap \tau_2 \\ t &\in ((\tau_1 + d_1) \cap \tau_2) - d_1 \end{aligned}$$

– For (II), let  $\mathbf{u} = \langle o_1, o_3, \tau, d \rangle \in \llbracket q \rrbracket_G^{[t]}$ , and let  $t \in \tau$ .

We show that  $\langle o_1, o_3, t, t + d \rangle \in \llbracket q \rrbracket_G$ .

Because  $\mathbf{u} \in \llbracket q \rrbracket_G^{[t]}$ , from the definition of  $\llbracket q \rrbracket_G^{[t]}$ , there are  $\tau_1, \tau_2, d_1, d_2$  and  $o_2$  s.t.:

- (i)  $d = d_1 + d_2$
- (ii)  $\tau = ((\tau_1 + d_1) \cap \tau_2) - d_1$
- (iii)  $\langle o_1, o_2, \tau_1, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G^{[t]}$
- (iv)  $\langle o_2, o_3, \tau_2, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G^{[t]}$

Since  $t \in \tau$ , from (ii), we have

$$t \in ((\tau_1 + d_1) \cap \tau_2) - d_1 \quad (14)$$

$$t + d_1 \in (((\tau_1 + d_1) \cap \tau_2) - d_1) + d_1 \quad (15)$$

$$t + d_1 \in (\tau_1 + d_1) \cap \tau_2 \quad (16)$$

$$t + d_1 \in \tau_1 + d_1 \quad (17)$$

$$t \in \tau_1 \quad (18)$$

From (iii), by IH, for any  $t' \in \tau_1$

$$\langle o_1, o_2, t' + d_1 \rangle \in \llbracket q \rrbracket_G$$

In particular, from (18)

$$\langle o_1, o_2, t, t + d_1 \rangle \in \llbracket q \rrbracket_G \quad (19)$$

And from (iv), by IH, for any  $t'' \in \tau_2$

$$\langle o_2, o_3, t'', t'' + d_2 \rangle \in \llbracket q \rrbracket_G$$

In particular, from (16)

$$\langle o_2, o_3, t + d_1, (t + d_1) + d_2 \rangle \in \llbracket q \rrbracket_G \quad (20)$$

So from (19), (20) and the definition of  $\llbracket q \rrbracket_G$

$$\langle o_1, o_3, t, t + d_1 + d_2 \rangle \in \llbracket q \rrbracket_G$$

□

### 3.4 In $\mathcal{U}^{[d]}$

#### 3.4.1 Definition.

We start with the case where  $q$  is an expression for the symbol test in the grammar of Definition 3.1.

The definitions of  $\llbracket \text{pred} \rrbracket_G^{[t]}$  and  $\llbracket \neg \text{test} \rrbracket_G^{[t]}$  are already provided in the article, we reproduce them here for completeness:

$$\begin{aligned} \llbracket \text{pred} \rrbracket_G^{[d]} &= \{ \langle o, o, t, [0, 0] \rangle \mid t \in \tau \text{ for some } \tau \in \text{val}(o, \text{pred}) \} \\ \llbracket \text{F} \rrbracket_G^{[d]} &= \{ \langle v, e, t, [0, 0] \rangle \mid \text{src}(e) = v, t \in \mathcal{T}_G \} \cup \{ \langle e, v, t, [0, 0] \rangle \mid \text{tgt}(e) = v, t \in \mathcal{T}_G \} \\ \llbracket \text{B} \rrbracket_G^{[d]} &= \{ \langle v, e, t, [0, 0] \rangle \mid \text{tgt}(e) = v, t \in \mathcal{T}_G \} \cup \{ \langle e, v, t, [0, 0] \rangle \mid \text{src}(e) = v, t \in \mathcal{T}_G \} \\ \llbracket (? \text{path}) \rrbracket_G^{[d]} &= \{ \langle o, o, t, [0, 0] \rangle \mid \exists o', \delta: \langle o, o', t, \delta \rangle \in \llbracket \text{path} \rrbracket_G^{[d]} \} \\ \llbracket \text{test}_1 \vee \text{test}_2 \rrbracket_G^{[d]} &= \llbracket \text{test}_1 \rrbracket_G^{[d]} \cup \llbracket \text{test}_2 \rrbracket_G^{[d]} \\ \llbracket \text{test}_1 \wedge \text{test}_2 \rrbracket_G^{[d]} &= \llbracket \text{test}_1 \rrbracket_G^{[d]} \cap \llbracket \text{test}_2 \rrbracket_G^{[d]} \\ \llbracket \neg \text{test} \rrbracket_G^{[d]} &= \left\{ \langle o, o, t, [0, 0] \rangle \mid o \in N \cup E, t \in \mathcal{T}_G \setminus \{ t' \mid \langle o, o, t', [0, 0] \rangle \in \llbracket \text{test} \rrbracket_G^{[d]} \} \right\} \end{aligned}$$

Next, we consider the operators  $(\text{path}_1 + \text{path}_2)$ ,  $(\text{path}[m, \_])$  and  $(\text{path}[m, n])$ .

For these cases,  $\llbracket q \rrbracket_G^{[t, d]}$  is once again defined analogously to  $\llbracket q \rrbracket_G$ , in terms of temporal join (a.k.a.  $\text{path}_1/\text{path}_2$ ) and set union.

We only write the definitions here for the sake of completeness:

$$\begin{aligned} \llbracket \text{path}_1 + \text{path}_2 \rrbracket_G^{[t,d]} &= \llbracket \text{path}_1 \rrbracket_G^{[t,d]} \cup \llbracket \text{path}_2 \rrbracket_G^{[t,d]} \\ \llbracket \text{path}[m, n] \rrbracket_G &= \bigcup_{k=m}^n \llbracket \text{path}^k \rrbracket_G^{[t,d]} \\ \llbracket \text{path}[m, \_] \rrbracket_G &= \bigcup_{k \geq m} \llbracket \text{path}^k \rrbracket_G^{[t,d]} \end{aligned}$$

The only remaining operators are temporal join ( $\text{path}_1/\text{path}_2$ ) and temporal navigation ( $T_\delta$ ), already defined in the article. We reproduce here these two definition for convenience:

$$\begin{aligned} \llbracket \text{path}_1/\text{path}_2 \rrbracket_G^{[d]} &= \left\{ \langle o_1, o_3, t_1, \delta_2 + t_2 - t_1 \rangle \mid \exists o_2: \langle o_1, o_2, t_1, \delta_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G^{[d]} \wedge \langle o_2, o_3, t_2, \delta_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G^{[d]} \wedge t_2 \in t_1 + \delta_1 \right\} \\ \llbracket T_\delta \rrbracket_G^{[d]} &= \left\{ \langle o, o, t, ((\delta + t) \cap \mathcal{T}_G) - t \rangle \mid o \in N \cup E, t \in \mathcal{T}_G, (\delta + t) \cap \mathcal{T}_G \neq \emptyset \right\} \end{aligned}$$

We also reproduce the alternative characterization of  $\llbracket T_\delta \rrbracket_G^{[d]}$  provided in the article, as a unary operator:

$$\llbracket q/T_\delta \rrbracket_G^{[d]} = \{ \langle o_1, o_2, t, (\delta' + \delta) \cap \mathcal{T}_G \rangle \mid \langle o_1, o_2, t, \delta' \rangle \in \llbracket q \rrbracket_G^{[d]}, (t + (\delta' + \delta)) \cap \mathcal{T}_G \neq \emptyset \}$$

### 3.4.2 Correctness.

The following result states that the representation  $\llbracket q \rrbracket_G^{[d]}$  is correct:

**PROPOSITION 3.4.** *Let  $G = \langle N, E, \text{conn}, \mathcal{T}_G, \text{val} \rangle$  be a TPG and  $q$  a TRPQ. Then the unfolding of  $\llbracket q \rrbracket_G^{[d]}$  is  $\llbracket q \rrbracket_G$ .*

**PROOF.**

Let  $G = \langle N, E, \text{conn}, \mathcal{T}_G, \text{val} \rangle$  be a TG, and let  $q$  be a TRPQ.

We show below that:

- (I) for any  $\langle o_1, o_2, t, d \rangle \in \llbracket q \rrbracket_G$ , there is a  $\delta \in \text{intv}(\mathcal{T})$  such that
  - (a)  $\langle o_1, o_2, t, \delta \rangle \in \llbracket q \rrbracket_G^{[d]}$ , and
  - (b)  $d \in \delta$ ,
- (II) for any  $\langle o_1, o_2, t, \delta \rangle \in \llbracket q \rrbracket_G^{[d]}$  for any  $d \in \delta$ ,  $\langle o_1, o_2, t, d \rangle$  is in  $\llbracket q \rrbracket_G$ .

We proceed once again by induction on the structure of  $q$ .

If  $q$  is of the form *pred*, F, B, (test  $\vee$  test), (test  $\wedge$  test),  $\neg$ test, (path + path), path[m, n] or path[m, \_], then (I) and (II) immediately follow from the definitions of  $\llbracket q \rrbracket_G$  and  $\llbracket q \rrbracket_G^{[d]}$ .

If  $q$  is of the form (?path), then the proof is nearly identical to one already provided for  $\llbracket (?path) \rrbracket_G^{[t]}$ .

So we focus below on the two remaining cases:

- $q = T_\delta$ .

From the above definitions, we have:

$$\begin{aligned} \llbracket q \rrbracket_G &= \{ \langle o, o, t, d \rangle \mid o \in (N \cup E), t \in \mathcal{T}_G, d \in \delta, t + d \in \mathcal{T}_G \} \\ \llbracket q \rrbracket_G^{[d]} &= \{ \langle o, o, t, ((\delta + t) \cap \mathcal{T}_G) - t \rangle \mid o \in N \cup E, t \in \mathcal{T}_G, (\delta + t) \cap \mathcal{T}_G \neq \emptyset \} \end{aligned}$$

- For (I), let  $\mathbf{v} = \langle o, o, t, d \rangle \in \llbracket q \rrbracket_G$ .  
And let  $\mathbf{u} = \langle o, o, t, ((\delta + t) \cap \mathcal{T}_G) - t \rangle$  in  $\mathcal{U}^{[d]}$ .  
For (Ia) we show that  $\mathbf{u} \in \llbracket q \rrbracket_G^{[d]}$ .  
From  $\mathbf{v} \in \llbracket q \rrbracket_G$ , we get  $o \in N \cup E$  and  $t \in \mathcal{T}_G$ .  
Besides, because  $\mathbf{v} \in \llbracket q \rrbracket_G$  still,

$$t + d \in \mathcal{T}_G \tag{21}$$

and

$$d \in \delta \tag{22}$$

$$t + d \in t + \delta \tag{23}$$

So from (21) and (23)

$$t + d \in (\delta + t) \cap \mathcal{T}_G \tag{24}$$

$$(\delta + t) \cap \mathcal{T}_G \neq \emptyset \tag{25}$$

Together with the definition of  $\langle q \rangle_G^{[d]}$ , this implies  $\mathbf{u} \in \langle q \rangle_G^{[d]}$ , which concludes the proof for (Ia).  
Finally, from (24), we get

$$t + d - t \in ((\delta + t) \cap \mathcal{T}_G) - t \quad (26)$$

$$d \in ((\delta + t) \cap \mathcal{T}_G) - t \quad (27)$$

which proves (Ib).

- For (II), let  $\mathbf{u} = \langle o, o, t, ((\delta + t) \cap \mathcal{T}_G) - t \rangle \in \langle q \rangle_G^{[d]}$ , and let  $d \in ((\delta + t) \cap \mathcal{T}_G) - t$ .

From  $\mathbf{u} \in \langle q \rangle_G^{[d]}$ , we get  $o \in N \cup E$  and  $t \in \mathcal{T}_G$ .

So to conclude the proof, it is sufficient to show that (i)  $d \in \delta$  and (ii)  $t + d \in \mathcal{T}_G$ .

By assumption, we have

$$d \in ((\delta + t) \cap \mathcal{T}_G) - t \quad (28)$$

$$d + t \in (\delta + t) \cap \mathcal{T}_G \quad (29)$$

$$d + t \in \mathcal{T}_G \quad (30)$$

which proves (ii).

And from (29), we also get

$$d + t \in \delta + t$$

$$d + t - t \in (\delta + t) - t$$

$$d \in \delta$$

which proves (i).

- $q = \text{path}_1 / \text{path}_2$ .

From the above definitions, we have:

$$\begin{aligned} \llbracket q \rrbracket_G &= \{ \langle o_1, o_3, t, d_1 + d_2 \rangle \mid \exists o_2: \langle o_1, o_2, t, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G \wedge \langle o_2, o_3, t + d_1, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G \} \\ \langle q \rangle_G^{[d]} &= \left\{ \langle o_1, o_3, t_1, \delta_2 + t_2 - t_1 \rangle \mid \exists o_2: \langle o_1, o_2, t_1, \delta_1 \rangle \in \langle \text{path}_1 \rangle_G^{[d]} \wedge \langle o_2, o_3, t_2, \delta_2 \rangle \in \langle \text{path}_2 \rangle_G^{[d]} \wedge t_2 \in t_1 + \delta_1 \right\} \end{aligned}$$

- For (I), let  $\mathbf{v} = \langle o_1, o_3, t, d \rangle \in \llbracket q \rrbracket_G$ .

From the definition of  $\llbracket q \rrbracket_G$ , there are  $o_2, d_1$  and  $d_2$  such that  $\langle o_1, o_2, t, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G$ ,  $\langle o_2, o_3, t + d_1, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G$  and  $d = d_1 + d_2$ .

By IH, because  $\langle o_1, o_2, t, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G$ , there is a  $\delta_1$  such that  $d_1 \in \delta_1$  and

$$\langle o_1, o_2, t, \delta_1 \rangle \in \langle \text{path}_1 \rangle_G^{[d]} \quad (31)$$

And similarly, because  $\langle o_2, o_3, t + d_1, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G$ , there is a  $\delta_2$  such that  $d_2 \in \delta_2$  and

$$\langle o_2, o_3, t + d_1, \delta_2 \rangle \in \langle \text{path}_2 \rangle_G^{[d]} \quad (32)$$

Next, since  $d \in \delta_1$

$$t + d_1 \in t + \delta_1 \quad (33)$$

So from (31), (32), (33) and the definition of  $\langle q \rangle_G^{[d]}$  (replacing  $t_1$  with  $t$  and  $t_2$  with  $t + d_1$ ), we get

$$\langle o_1, o_2, t, \delta_2 + (t + d_1) - t \rangle \in \langle q \rangle_G^{[d]}$$

which proves (Ia).

And in order to prove (Ib), we only need to show that  $d \in \delta_2 + (t + d_1) - t$ , or in other words that

$$d \in \delta_2 + d_1$$

We know that

$$d_2 \in \delta_2 \quad (34)$$

$$d_2 + d_1 \in \delta_2 + d_1 \quad (35)$$

Together with the fact that  $d = d_1 + d_2$ , this concludes the proof for (Ib).

- For (II), let  $\mathbf{u} = \langle o_1, o_3, t_1, \delta \rangle \in \langle q \rangle_G^{[d]}$ , and let  $d \in \delta$ .

Because  $\mathbf{u} \in \langle q \rangle_G^{[d]}$ , from the definition of  $\langle q \rangle_G^{[d]}$ , there are  $\delta_1, \delta_2, t_2$  and  $o_2$  s.t.:

$$(i) \quad \delta = \delta_2 + t_2 - t_1$$

$$(ii) \quad t_2 \in t_1 + \delta_1$$

$$(iii) \quad \langle o_1, o_2, t_1, \delta_1 \rangle \in \langle \text{path}_1 \rangle_G^{[d]}$$



(iv)  $\langle o_2, o_3, t_2, \delta_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G^{[d]}$   
 From (i) and (ii), we get

$$\begin{aligned}\delta &= \delta_2 + (t_1 + \delta_1) - t_1 \\ &= \delta_2 + \delta_1\end{aligned}$$

Together with  $d \in \delta$ , this implies that there are  $d_1 \in \delta_1$  and  $d_2 \in \delta_2$  such that  $d = d_1 + d_2$ .

Next, because  $d_1 \in \delta_1$ , from (iii), by IH

$$\langle o_1, o_2, t_1, t_1 + d_1 \rangle \in \llbracket q \rrbracket_G \quad (36)$$

And similarly, because  $d_2 \in \delta_2$ , from (iv)

$$\langle o_2, o_3, t_2, t_2 + d_2 \rangle \in \llbracket q \rrbracket_G \quad (37)$$

So from (36), (37) and the definition of  $\llbracket q \rrbracket_G$

$$\langle o_1, o_3, t_1, d_1 + d_2 \rangle \in \llbracket q \rrbracket_G \quad (38)$$

Together with the fact that  $d = d_1 + d_2$ , this concludes the proof for (II).  $\square$

### 3.5 In $\mathcal{U}^{[t,d]}$

#### 3.5.1 Definition.

We start with the case where  $q$  is an expression for the symbol test in the grammar of Definition 3.1.

As a consequence of Lemma 3.2,  $\llbracket q \rrbracket_G^{[t,d]}$  can be trivially defined out of  $\llbracket q \rrbracket_G^{[t]}$  by replacing the distance 0 with the interval  $[0, 0]$ , i.e.

$$\llbracket \text{test} \rrbracket_G^{[t,d]} = \{ \langle o, o, \tau, [0, 0] \rangle \mid \langle o, o, \tau, 0 \rangle \in \llbracket \text{test} \rrbracket_G^{[t]} \}$$

Next, if  $q$  is of the form  $(\text{path}_1 + \text{path}_2)$ ,  $(\text{path}[m, \_])$  or  $(\text{path}[m, n])$ , then the definition of  $\llbracket q \rrbracket_G^{[t,d]}$  is once again nearly identical to the one of  $\llbracket q \rrbracket_G$ :

$$\begin{aligned}\llbracket \text{path}_1 + \text{path}_2 \rrbracket_G^{[t,d]} &= \llbracket \text{path}_1 \rrbracket_G^{[t,d]} \cup \llbracket \text{path}_2 \rrbracket_G^{[t,d]} \\ \llbracket \text{path}[m, n] \rrbracket_G &= \bigcup_{k=m}^n \llbracket \text{path}^k \rrbracket_G^{[t,d]} \\ \llbracket \text{path}[m, \_] \rrbracket_G &= \bigcup_{k \geq m} \llbracket \text{path}^k \rrbracket_G^{[t,d]}\end{aligned}$$

The only remaining operators are temporal join  $(\text{path}_1 / \text{path}_2)$  and temporal navigation  $(T_\delta)$ , already defined in the article, and reproduced here for convenience:

$$\begin{aligned}\llbracket \text{path}_1 / \text{path}_2 \rrbracket_G^{[t,d]} &= \bigcup \{ \mathbf{u}_1 \bowtie \mathbf{u}_2 \mid \mathbf{u}_1 \in \llbracket \text{path}_1 \rrbracket_G^{[t,d]}, \mathbf{u}_2 \in \llbracket \text{path}_2 \rrbracket_G^{[t,d]} \} \\ \llbracket T_\delta \rrbracket_G^{[t,d]} &= \bigcup_{o \in N \cup E} \{ \langle o, o, \mathcal{T}_G, \delta \rangle \bowtie \langle o, o, \mathcal{T}_G, [0, 0] \rangle \}\end{aligned}$$

where  $\mathbf{u}_1 \bowtie \mathbf{u}_2$  is defined as follows.

Let  $\mathbf{u}_1 = \langle o_1, o_2, \tau_1, \delta_1 \rangle$  and  $\mathbf{u}_2 = \langle o_3, o_4, \tau_2, \delta_2 \rangle$ .

If  $o_2 \neq o_3$ , then  $\mathbf{u}_1 \bowtie \mathbf{u}_2 = \emptyset$ .

Otherwise, let:

$$\tau = ((\tau_1 + \delta_1) \cap \tau_2) \ominus \delta_1 \cap \tau_1$$

And for every  $t \in \tau$ , let

$$\delta(t) = \delta_1 \downarrow b_{\delta_1} + \max(0, b - t), \quad e_{\delta_1} - \max(0, t - e) \downarrow_{\delta_1}$$

Then

$$\mathbf{u}_1 \bowtie \mathbf{u}_2 = \{ \langle o_1, o_4, [t, t], \delta(t) + \delta_2 \rangle \mid t \in \tau \}$$

#### 3.5.2 Correctness.

If  $\mathbf{u} = \langle o_1, o_2, \tau, \delta \rangle \in \mathcal{U}^{[t,d]}$ , we call *temporal relation induced by  $\mathbf{u}$*  the set  $\{ \langle t, t + d \rangle \mid t \in \tau, d \in \delta \}$ .

We also define the binary operator  $\bowtie$ :  $(\mathcal{T} \times \mathcal{T}) \times (\mathcal{T} \times \mathcal{T}) \rightarrow (\mathcal{T} \times \mathcal{T})$  as in the article, i.e.

$$R_1 \bowtie R_2 = \{ t_1, t_3 \mid (t_1, t_2) \in R_1 \text{ and } (t_2, t_3) \in R_2 \text{ for some } t_2 \}$$

We can now formulate the following lemma:

LEMMA 3.5. Let  $\mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}^{[t,d]}$ , and for  $i \in \{1, 2\}$ , let  $R_i$  denote the temporal relation induced by  $\mathbf{u}_i$ . If  $\mathbf{u}_1 \bowtie \mathbf{u}_2 \neq \emptyset$ , then

$$R_1 \bowtie R_2 = \bigcup_{\langle o_1, o_2, \tau, \delta \rangle \in \mathbf{u}_1 \bowtie \mathbf{u}_2} \{ \langle t, t + d \rangle \mid t \in \tau, d \in \delta \}$$

The following result states that the representation  $\llbracket q \rrbracket_G^{[t,d]}$  is correct:

PROPOSITION 3.6. Let  $G = \langle N, E, \text{conn}, \mathcal{T}_G, \text{val} \rangle$  be a TPG and  $q$  a TRPQ. Then the unfolding of  $\langle q \rangle_G^{[t,d]}$  is  $\llbracket q \rrbracket_G$ .

PROOF.

Let  $G = \langle N, E, \text{conn}, \mathcal{T}_G, \text{val} \rangle$  be a TG, and let  $q$  be a TRPQ.

We show below that:

- (I) for any  $\langle o_1, o_2, t, d \rangle \in \llbracket q \rrbracket_G$ , there are  $\tau, \delta \in \text{intv}(\mathcal{T})$  such that
  - (a)  $\langle o_1, o_2, \tau, \delta \rangle \in \langle q \rangle_G^{[t,d]}$ ,
  - (b)  $t \in \tau$ , and
  - (c)  $d \in \delta$ .
- (II) for any  $\langle o_1, o_2, \tau, \delta \rangle \in \langle q \rangle_G^{[t,d]}$  for any  $(t, d) \in \tau \times \delta$ ,  $\langle o_1, o_2, t, d \rangle$  is in  $\llbracket q \rrbracket_G$ .

We proceed once again by induction on the structure of  $q$ .

If  $q$  is of the form *pred*, *F*, *B*, (*test*  $\vee$  *test*), (*path* + *path*), *path*[ $m, n$ ] or *path*[ $m, \_$ ], then (I) and (II) immediately follow from the definitions of  $\llbracket q \rrbracket_G$  and  $\langle q \rangle_G^{[t,d]}$ .

If  $q$  is of the form *test*  $\wedge$  *test*,  $\neg$ *test* or (*?path*), then the proof is nearly identical to the one already provided for  $\langle q \rangle_G^{[t]}$ .

So we focus below on the two remaining cases:

- $q = \text{path}_1 / \text{path}_2$ .

From the above definitions, we have:

$$\begin{aligned} \llbracket q \rrbracket_G &= \{ \langle o_1, o_3, t, d_1 + d_2 \rangle \mid \exists o_2: \langle o_1, o_2, t, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G \wedge \langle o_2, o_3, t + d_1, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G \} \\ \langle \text{path}_1 / \text{path}_2 \rangle_G^{[t,d]} &= \bigcup \{ \mathbf{u}_1 \bowtie \mathbf{u}_2 \mid \mathbf{u}_1 \in \langle \text{path}_1 \rangle_G^{[t,d]}, \mathbf{u}_2 \in \langle \text{path}_2 \rangle_G^{[t,d]} \} \end{aligned}$$

- For (I), let  $\mathbf{v} = \langle o_1, o_3, t, d \rangle \in \llbracket q \rrbracket_G$ .

From the definition of  $\llbracket q \rrbracket_G$ , there are  $o_2, d_1$  and  $d_2$  such that  $\langle o_1, o_2, t, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G$ ,  $\langle o_2, o_3, t + d_1, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G$  and  $d = d_1 + d_2$ .

By IH, because  $\langle o_1, o_2, t, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G$ , there are  $\tau_1$  and  $\delta_1$  such that  $t \in \tau_1$ ,  $d_1 \in \delta_1$  and

$$\langle o_1, o_2, \tau_1, \delta_1 \rangle \in \langle \text{path}_1 \rangle_G^{[t,d]} \quad (39)$$

Let  $R_1$  be the temporal relation induced by this tuple  $\langle o_1, o_2, \tau_1, \delta_1 \rangle$ .

Since  $t \in \tau_1$  and  $d_1 \in \delta_1$ , we have

$$(t, t + d_1) \in R_1 \quad (40)$$

Similarly, because  $\langle o_2, o_3, t + d_1, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G$ , there are  $\tau_2$  and  $\delta_2$  such that  $t + d_1 \in \tau_2$ ,  $d_2 \in \delta_2$  and

$$\langle o_2, o_3, \tau_2, \delta_2 \rangle \in \langle \text{path}_2 \rangle_G^{[t,d]} \quad (41)$$

Let  $R_2$  be the temporal relation induced by this tuple  $\langle o_2, o_3, \tau_2, \delta_2 \rangle$ .

Since  $t + d_1 \in \tau_2$  and  $d_2 \in \delta_2$ , we have

$$(t + d_1, t + d_1 + d_2) \in R_2 \quad (42)$$

So from (40), (42) and Lemma 3.5, there are  $\tau$  and  $\delta$  such that  $\langle o_1, o_3, \tau, \delta \rangle \in \mathbf{u}_1 \bowtie \mathbf{u}_2$ ,  $t \in \tau$  and  $d_1 + d_2 = d \in \delta$ , which concludes the proof for (I).

- For (II), let  $\mathbf{u} = \langle o_1, o_3, t_1, \delta \rangle \in \langle q \rangle_G^{[t,d]}$ , and let  $(t, d) \in \tau \times \delta$ .

Because  $\mathbf{u} \in \langle q \rangle_G^{[t,d]}$ , from the definition of  $\langle q \rangle_G^{[t,d]}$ , there are  $\mathbf{u}_1$  and  $\mathbf{u}_2$  s.t.:

- (i)  $\mathbf{u} \in \mathbf{u}_1 \bowtie \mathbf{u}_2$
- (ii)  $\mathbf{u}_1 \in \langle \text{path}_1 \rangle_G^{[t,d]}$
- (iii)  $\mathbf{u}_2 \in \langle \text{path}_2 \rangle_G^{[t,d]}$

Let  $R_i$  be the temporal relation induced by  $\mathbf{u}_i$  for  $i \in \{1, 2\}$ .

From (i), and Lemma 3.5,

$$(t, t + d) \in R_1 \bowtie R_2 \quad (43)$$

Now let  $\mathbf{u}_1 = \langle o_1, o_2, \tau_1, \delta_1 \rangle$  and  $\mathbf{u}_2 = \langle o_2, o_3, \tau_2, \delta_2 \rangle$  for some  $o_2, \tau_1, \tau_2, \delta_1$  and  $\delta_2$ .

From (43) and the definition of  $\bowtie$ , there must be  $d_1$  and  $d_2$  s.t.  $d = d_1 + d_2$ ,  $t \in \tau_1$ ,  $d_1 \in \delta_1$ ,  $t + d_1 \in \tau_2$  and  $d_2 \in \delta_2$ .

So from ii, and iii, by IH

$$\langle o_1, o_2, t, d_1 \rangle \in \llbracket \text{path}_1 \rrbracket_G \quad (44)$$

$$\langle o_2, o_3, t + d_1, d_2 \rangle \in \llbracket \text{path}_2 \rrbracket_G \quad (45)$$

So from (44), (45) and the definition of  $\llbracket q \rrbracket_G$

$$\langle o_1, o_3, t, d_1 + d_2 \rangle \in \llbracket q \rrbracket_G,$$

which concludes the proof for (II).  $\square$

### 3.6 In $\mathcal{U}^{[t,d],b,e}$

#### 3.6.1 Definition.

If  $q$  is an expression for the symbol test in the grammar of Definition ??, then the definition of  $\langle q \rangle_G^{[t,d],b,e}$  is nearly identical to the one of  $\langle q \rangle_G^{[t,d]}$ , extending each tuple  $\{\langle o, o, \tau, [0, 0] \rangle$  with  $b_\tau$  and  $e_\tau$ , i.e.

$$\langle \text{test} \rangle_G^{[t,d],b,e} = \{ \langle o, o, \tau, [0, 0], b_\tau, e_\tau \rangle \mid \{ \langle o, o, \tau, [0, 0] \rangle \in \langle \text{test} \rangle_G^{[t,d]} \}$$

Next, if  $q$  is of the form  $(\text{path}_1 + \text{path}_2)$ ,  $(\text{path}[m, \_])$  or  $(\text{path}[m, n])$ , then the definition of  $\langle q \rangle_G^{[t,d]}$  is once again nearly identical to the one of  $\llbracket q \rrbracket_G$ :

$$\begin{aligned} \langle \text{path}_1 + \text{path}_2 \rangle_G^{[t,d],b,e} &= \langle \text{path}_1 \rangle_G^{[t,d],b,e} \cup \langle \text{path}_2 \rangle_G^{[t,d],b,e} \\ \llbracket \text{path}[m, n] \rrbracket_G &= \bigcup_{k=m}^n \langle \text{path}^k \rangle_G^{[t,d],b,e} \\ \llbracket \text{path}[m, \_] \rrbracket_G &= \bigcup_{k \geq m} \langle \text{path}^k \rangle_G^{[t,d],b,e} \end{aligned}$$

So the only remaining operator are temporal join  $(\text{path}_1 / \text{path}_2)$  and temporal navigation  $(T_\delta)$ , already defined in the article. We reproduce here these two definition for convenience:

$$\begin{aligned} \langle \text{path}_1 / \text{path}_2 \rangle_G^{[t,d],b,e} &= \{ \mathbf{u}_1 \bowtie \mathbf{u}_2 \mid \mathbf{u}_1 \in \langle \text{path}_1 \rangle_G^{[t,d],b,e}, \mathbf{u}_2 \in \langle \text{path}_2 \rangle_G^{[t,d],b,e}, \mathbf{u}_1 \sim \mathbf{u}_2 \} \\ \langle T_\delta \rangle_G^{[t,d],b,e} &= \{ \langle o, o, \mathcal{T}_G, \delta, b_{\mathcal{T}_G}, e_{\mathcal{T}_G} \rangle \bowtie \langle o, o, \mathcal{T}_G, [0, 0], b_{\mathcal{T}_G}, e_{\mathcal{T}_G} \rangle \mid o \in N \cup E \} \end{aligned}$$

where  $\mathbf{u}_1 \sim \mathbf{u}_2$  and  $\mathbf{u}_1 \bowtie \mathbf{u}_2$  are defined by:

$$\begin{aligned} \langle o_1, o_2, \tau_1, \delta_1, b_1, e_1 \rangle &\sim \langle o_3, o_4, \tau_2, \delta_2, e_2, b_2 \rangle \text{ iff } o_2 = o_3 \\ \langle o_1, o_3, \tau_1, \delta_1, b_1, e_1 \rangle &\bowtie \langle o_3, o_4, \tau_2, \delta_2, e_2, b_2 \rangle = \langle o_1, o_4, \tau, \delta_1 + \delta_2, b, e \rangle \\ &\text{with} \\ \tau &= ((\tau_1 + \delta_1) \cap \tau_2) \ominus \delta_1 \cap \tau_1 \\ b &= \max(b_1, b_2 - b_{\delta_1}) \\ e &= \min(e_1, e_2 - e_{\delta_1}) \end{aligned}$$

**3.6.2 Correctness.** Similarly to what we did above for  $\mathcal{U}^{[t,d]}$ , if  $\mathbf{u} = \langle o_1, o_2, \tau, \delta, b, e \rangle \in \mathcal{U}^{[t,d],b,e}$ , we call *temporal relation induced by  $\mathbf{u}$*  the set  $\{(t, t + d) \mid t \in \tau, d \in \delta(t)\}$ .

We can now formulate a lemma analogoud to Lemma 3.5:

**LEMMA 3.7.** *Let  $\mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}^{[t,d],b,e}$ , and for  $i \in \{1, 2\}$ , let  $R_i$  denote the temporal relation induced by  $\mathbf{u}_i$ . If  $\mathbf{u}_1 \sim \mathbf{u}_2$  and  $\mathbf{u}_1 \bowtie \mathbf{u}_2 = \langle o_1, o_3, \tau, \delta, b, e \rangle$ , then*

$$R_1 \bowtie R_2 = \{(t, t + d) \mid t \in \tau, d \in \delta(t)\}$$

**PROOF.** Let  $\mathbf{u}_1 = \langle o_1, o_2, \tau_1, \delta_1 \rangle$  and  $\mathbf{u}_2 = \langle o_2, o_3, \tau_2, \delta_2 \rangle$ . As explained in Section 2, for  $i \in \{1, 2\}$  and  $t \in \tau_i$ , we use  $\delta_i(t)$  for the interval

$$\delta_i \upharpoonright b_{\delta_i} + \max(0, b_i - t), e_{\delta_i} - \max(0, t - e_i) \downharpoonright \delta_i$$

We need to prove that (i)  $\tau = \text{dom}(R_1 \bowtie R_2)$  and that (ii) for each  $t \in \tau$ ,

$$t + \delta(t) = \{t' \mid (t, t') \in R_1 \bowtie R_2\}$$

The proof of (i) is nearly identical to the one provided above for Lemma 3.5.

For (ii), let  $t \in \tau$ . We use

- $a$  for the least value s.t.  $(t, a) \in \text{range}(R_1) \cap \text{dom}(R_2)$ , and
- $a'$  for the least value s.t.  $(a, a') \in R_2$

Then  $a'$  is also the least value s.t.  $(t, a') \in R_1 \bowtie R_2$ .

Analogously, we use  $z$  for the greatest value s.t.  $(t, z) \in \text{range}(R_1) \cap \text{dom}(R_2)$ , and  $z'$  for the greatest value s.t.  $(z, z') \in R_2$ .

Then  $z'$  is also the greatest value s.t.  $(t, z') \in R_1 \bowtie R_2$ .

From Lemma ??:

- $\{t\} \times (t + [a, z]) \subseteq R_1$ , and
- $[a, b] \times [a', z'] \subseteq R_2$

Therefore  $[a', z'] = \{c \mid (t, c) \in R_1 \bowtie R_2\}$ .

To conclude the proof, we show that  $t + \delta_t = [a, z]$ .

We only prove that  $t + b_{\delta_t} = a$  (the proof that  $t + e_{\delta_t} = z$  is symmetric).

Following the definition of  $b$ , we consider 2 cases:

- (1)  $b_1 < b_2 - b_{\delta_1}$
- (2)  $b_1 \geq b_2 - b_{\delta_1}$

For Case (1), we get

$$b_1 < b_2 - b_{\delta_1} \quad (46)$$

$$\max(b_1, b_2 - b_{\delta_1}) = b_2 - b_{\delta_1} \quad (47)$$

$$b = b_2 - b_{\delta_1} \quad \text{from the definition of } b \quad (48)$$

And in Case (1) still, we get:

$$b_1 < b_2 - b_{\delta_1} \quad (49)$$

$$0 < b_2 - b_{\delta_1} - b_1 \quad (50)$$

$$\max(0, b_2 - b_{\delta_1} - b_1) = b_2 - b_{\delta_1} - b_1 \quad (51)$$

Next, we consider two subcases:

- (i)  $t < b_2 - b_{\delta_1}$
- (ii)  $t \geq b_2 - b_{\delta_1}$

In Case (i), we get

$$t < b_2 - b_{\delta_1} \quad (52)$$

$$0 < b_2 - b_{\delta_1} - t \quad (53)$$

$$\max(0, b_2 - b_{\delta_1} - t) = b_2 - b_{\delta_1} - t \quad (54)$$

Now from the definition of  $\delta_t$ ,

$$b_{\delta_t} = b_{\delta_1} + b_{\delta_2} + \max(0, b - t) \quad (55)$$

$$= b_{\delta_1} + b_{\delta_2} + \max(0, b_2 - b_{\delta_1} - t) \quad \text{from (48)} \quad (56)$$

$$= b_{\delta_1} + b_{\delta_2} + b_2 - b_{\delta_1} - t \quad \text{from (54)} \quad (57)$$

$$= b_{\delta_2} + b_2 - t \quad (58)$$

$$b_{\delta_t} + t = b_{\delta_2} + b_2 - t + t \quad (59)$$

$$= b_{\delta_2} + b_2 \quad (60)$$

Next, from the definition of  $a'$

$$a' = b_{\delta_2(a)} + a \quad (61)$$

$$= b_{\delta_2} + \max(0, b_2 - a) + a \quad (62)$$

And, from the definition of  $a$

$$a = b_{\delta_1(t)} + t \quad (63)$$

$$= b_{\delta_1} + \max(0, b_1 - t) + t \quad (64)$$

Then we have two further subcases:

- (I)  $t \geq b_1$ , or
- (II)  $t < b_1$

In case (I):

$$t \geq b_1 \quad (65)$$

$$0 \geq b_1 - t \quad (66)$$

$$\max(0, b_1 - t) = 0 \quad (67)$$

$$a = b_{\delta_1} + t \quad \text{from (64)} \quad (68)$$

$$\max(0, b_2 - a) = \max(0, b_2 - b_{\delta_1} - t) \quad (69)$$

$$= b_2 - b_{\delta_1} - t \quad \text{from (54)} \quad (70)$$

$$= b_2 - a \quad \text{from (68)} \quad (71)$$

In case (II):

$$t < b_1 \quad (72)$$

$$0 < b_1 - t \quad (73)$$

$$\max(0, b_1 - t) = b_1 - t \quad (74)$$

$$a = b_{\delta_1} + b_1 - t + t \quad \text{from (64)} \quad (75)$$

$$= b_{\delta_1} + b_1 \quad (76)$$

$$\max(0, b_2 - a) = \max(0, b_2 - b_{\delta_1} - b_1) \quad (77)$$

$$= b_2 - b_{\delta_1} - b_1 \quad \text{from (51)} \quad (78)$$

$$= b_2 - a \quad \text{from (76)} \quad (79)$$

$$(80)$$

So in both cases (I) and (II), we get

$$\max(0, b_2 - a) = b_2 - a$$

Therefore from (62)

$$a' = b_{\delta_2} + b_2 - a + a \quad (81)$$

$$= b_{\delta_2} + b_2 \quad (82)$$

$$= t + b_{\delta_t} \quad \text{from (60)} \quad (83)$$

which concludes the proof for Case (1)- (i).

We continue with Case (1)- (ii).

From Case (ii):

$$t \geq b_2 - b_{\delta_1} \quad (84)$$

$$0 \geq b_2 - b_{\delta_1} - t \quad (85)$$

$$\max(0, b_2 - b_{\delta_1} - t) = 0 \quad (86)$$

Now from the definition of  $\delta_t$ :

$$b_{\delta_t} = b_{\delta_1} + b_{\delta_2} + \max(0, b - t) \quad (87)$$

$$= b_{\delta_1} + b_{\delta_2} + \max(0, b_2 - b_1 - t) \quad \text{from (48)} \quad (88)$$

$$= b_{\delta_1} + b_{\delta_2} \quad \text{from (86)} \quad (89)$$

$$b_{\delta_t} + t = b_{\delta_1} + b_{\delta_2} + t \quad (90)$$

Next, from Case (1) and Case (ii), by transitivity, we get

$$b_1 \leq t \quad (91)$$

$$\max(0, b_1 - t) = 0 \quad (92)$$

And from the definition of  $a$

$$a = b_{\delta_1(t)} + t \quad (93)$$

$$= b_{\delta_1} + \max(0, b_1 - t) + t \quad (94)$$

$$= b_{\delta_1} + t \quad \text{from (92)} \quad (95)$$

$$\geq b_{\delta_1} + b_2 - b_{\delta_1} \quad \text{from Case (ii)} \quad (96)$$

$$\geq b_2 \quad (97)$$

$$0 \geq b_2 - a \quad (98)$$

$$\max(0, b_2 - a) = 0 \quad (99)$$

Therefore from (62) and (99)

$$a' = b_{\delta_2} + a \quad (100)$$

$$= b_{\delta_2} + b_{\delta_1} + t \quad \text{from (95)} \quad (101)$$

$$= b_{\delta_t} + t \quad \text{from (60)} \quad (102)$$

which concludes the proof for Case (1)- (ii).

We continue with Case (2).

In this case, we get

$$b_1 \geq b_2 - b_{\delta_1} \quad (103)$$

$$\max(b_1, b_2 - b_{\delta_1}) = b_1 \quad (104)$$

$$b = b_1 \quad \text{from the definition of } b \quad (105)$$

And from Case (2) still, we derive

$$b_1 \geq b_2 - b_{\delta_1} \quad (106)$$

$$0 \geq b_2 - b_{\delta_1} - b_1 \quad (107)$$

$$\max(0, -b_{\delta_1} - b_1) = 0 \quad (108)$$

As well as

$$b_1 \geq b_2 - b_{\delta_1} \quad (109)$$

$$b_1 + b_{\delta_1} \geq b_2 \quad (110)$$

Next, we distinguish two subcases, namely

(a)  $t < b_1$  and

(b)  $t \geq b_1$

We start with Case (a).

In this case,

$$t < b_1 \quad (111)$$

$$0 < b_1 - t \quad (112)$$

$$\max(0, b_1 - t) = b_1 - t \quad (113)$$

And from the definition of  $\delta_t$ :

$$b_{\delta_t} = b_{\delta_1} + b_{\delta_2} + \max(0, b - t) \quad (114)$$

$$= b_{\delta_1} + b_{\delta_2} + \max(0, b_1 - t) \quad \text{from (105)} \quad (115)$$

$$= b_{\delta_1} + b_{\delta_2} + b_1 - t \quad \text{from (113)} \quad (116)$$

$$b_{\delta_t} + t = b_{\delta_1} + b_{\delta_2} + b_1 - t + t \quad (117)$$

$$= b_{\delta_1} + b_{\delta_2} + b_1 \quad (118)$$

Next, from the definition of  $a$

$$a = b_{\delta_1(t)} + t \quad (119)$$

$$= \max(0, b_1 - t) + b_{\delta_1} + t \quad (120)$$

$$= b_1 - t + b_{\delta_1} + t \quad \text{from (113)} \quad (121)$$

$$= b_1 + b_{\delta_1} \quad (122)$$

$$\text{So from (110)} \quad (123)$$

$$a \geq b_2 \quad (124)$$

$$0 \geq b_2 - a \quad (125)$$

$$\max(0, b_2 - a) = 0 \quad (126)$$

$$b_{\delta_2} + \max(0, b_2 - a) = b_{\delta_2} \quad (127)$$

$$b_{\delta_2(a)} = b_{\delta_2} \quad (128)$$

$$b_{\delta_2(a)} + a = b_{\delta_2} + a \quad (129)$$

$$a' = b_{\delta_2} + a \quad \text{from the defintiion of } a' \quad (130)$$

$$a' = b_{\delta_2} + b_1 + b_{\delta_1} \quad \text{from (122)} \quad (131)$$

$$a' = b_{\delta_t} + t \quad \text{from (118)} \quad (132)$$

which concludes the proof for Case (2)- (a).

We end with Case (2)- (b). In this case,

$$t \geq b_1 \quad (133)$$

$$0 \geq b_1 - t \quad (134)$$

$$\max(0, b_1 - t) = 0 \quad (135)$$

And from the definition of  $\delta_t$ :

$$b_{\delta_t} = b_{\delta_1} + b_{\delta_2} + \max(0, b - t) \quad (136)$$

$$= b_{\delta_1} + b_{\delta_2} + \max(0, b_1 - t) \quad \text{from (105)} \quad (137)$$

$$= b_{\delta_1} + b_{\delta_2} \quad \text{from (135)} \quad (138)$$

$$b_{\delta_t} + t = b_{\delta_1} + b_{\delta_2} + t \quad (139)$$

Next, from the definition of  $a$

$$a = b_{\delta_1(t)} + t \quad (140)$$

$$= \max(0, b_1 - t) + b_{\delta_1} + t \quad (141)$$

$$= b_{\delta_1} + t \quad \text{from (135)} \quad (142)$$

Now from Case (b)

$$b_1 + \leq t \quad (143)$$

$$b_1 + b_{\delta_1} \leq t + b_{\delta_1} \quad (144)$$

$$b_1 + b_{\delta_1} \leq a \quad \text{from (142)} \quad (145)$$

$$b_2 \leq a \quad \text{from (110), by transitivity} \quad (146)$$

$$b_2 - a \leq 0 \quad (147)$$

$$\max(0, b_2 - a) = 0 \quad (148)$$

$$b_{\delta_2} + \max(0, b_2 - a) = b_{\delta_2} \quad (149)$$

$$b_{\delta_2(a)} = b_{\delta_2} \quad (150)$$

$$b_{\delta_2(a)} + a = b_{\delta_2} + a \quad (151)$$

$$a' = b_{\delta_2} + a \quad \text{from the defintiion of } a' \quad (152)$$

$$= b_{\delta_2} + b_{\delta_1} + t \quad \text{from (142)} \quad (153)$$

$$= b_{\delta_t} + t \quad \text{from (139)} \quad (154)$$

which concludes the proof for Case (2)- (b).  $\square$

The following result states that the representation  $\langle q \rangle_G^{[t,d],b,e}$  is correct:

**PROPOSITION 3.8.** *Let  $G = \langle N, E, \text{conn}, \mathcal{T}_G, \text{val} \rangle$  be a TPG and  $q$  a TRPQ. Then the unfolding of  $\langle q \rangle_G^{[t,d],b,e}$  is  $\llbracket q \rrbracket_G$ .*

**PROOF.** Let  $G = \langle N, E, \text{conn}, \mathcal{T}_G, \text{val} \rangle$  be a TG, and let  $q$  be a TRPQ.

To prove the result, it is sufficient to show that:

- (I) for any  $\langle o_1, o_2, t, d \rangle \in \llbracket q \rrbracket_G$ , there are  $\tau, \delta \in \text{intv}(\mathcal{T})$  and  $b, e \in \mathcal{T}$  such that
  - (a)  $\langle o_1, o_2, \tau, \delta, b, e \rangle \in \langle q \rangle_G^{[t,d],b,e}$ ,
  - (b)  $t \in \tau$ , and
  - (c)  $d \in \delta(t)$  (where  $\delta(t)$  is defined in terms of  $t, \delta, b$  and  $e$ , as explained above).
- (II) for any  $\langle o_1, o_2, \tau, \delta, b, e \rangle \in \langle q \rangle_G^{[t,d],b,e}$  for any  $t \in \tau$  and  $d \in \delta(t)$ ,
  - $\langle o_1, o_2, t, d \rangle$  is in  $\llbracket q \rrbracket_G$ .

Again, the proof is by induction on the structure of  $q$ .

If  $q$  is of the form  $\text{pred}$ , F, B, (test  $\vee$  test), (path + path), path[ $m, n$ ] or path[ $m, \_$ ], then (I) and (II) immediately follow from the definitions of  $\llbracket q \rrbracket_G$  and  $\langle q \rangle_G^{[t,d],b,e}$ .

If  $q$  is of the form test  $\wedge$  test,  $\neg$ test or (?path), then the proof is nearly identical to the one already provided for  $\langle q \rangle_G^{[t]}$ .

And if  $q$  is of the form  $T_\delta$  or path<sub>1</sub>/path<sub>2</sub>, then the proof is nearly identical to the one already provided for  $\langle q \rangle_G^{[t,d]}$ , using Lemma 3.7 instead of 3.5.  $\square$

## 4 FINITENESS

## 5 COMPLEXITY OF QUERY ANSWERING

### 5.1 Problem

We define in this section a decision problem for each representation, similar to the problem COMPACT ANSWER<sup>[t]</sup> defined in the article. First, we define a (possibly partial) order over tuples of each representation, in the expected way, i.e.:

- over  $\mathcal{U}^{[t]}$ :  $\langle o_1, o_2, \tau_1, d_1 \rangle \sqsubseteq_{[t]} \langle o_3, o_4, \tau_2, d_2 \rangle$  iff  $\langle o_1, o_2, d_1 \rangle = \langle o_3, o_4, d_2 \rangle$  and  $\tau_1 \subseteq \tau_2$ ,
- over  $\mathcal{U}^{[d]}$ :  $\langle o_1, o_2, t_1, \delta_1 \rangle \sqsubseteq_{[d]} \langle o_3, o_4, t_2, \delta_2 \rangle$  iff  $\langle o_1, o_2, t_1 \rangle = \langle o_3, o_4, t_2 \rangle$  and  $\delta_1 \subseteq \delta_2$ ,
- over  $\mathcal{U}^{[t,d]}$ :  $\langle o_1, o_2, \tau_1, \delta_1 \rangle \sqsubseteq_{[t,d]} \langle o_3, o_4, \tau_2, \delta_2 \rangle$  iff  $\langle o_1, o_2 \rangle = \langle o_3, o_4 \rangle$ ,  $\tau_1 \subseteq \tau_2$  and  $\delta_1 \subseteq \delta_2$ ,
- over  $\mathcal{U}^{[t,d],b,e}$ :  $\langle o_1, o_2, \tau_1, \delta_1, b_1, b_2 \rangle \sqsubseteq_{[t,d],b,e} \langle o_3, o_4, \tau_2, \delta_2, b_2, e_2 \rangle$  iff  $\langle o_1, o_2 \rangle = \langle o_3, o_4 \rangle$ ,  $\tau_1 \subseteq \tau_2$  and  $\delta_1(t) \subseteq \delta_2(t)$  for all  $t \in \tau_1 \cap \tau_2$  (the notation of  $\delta_i(t)$  is explained above, in Section 2).

Now let  $x$  be one of  $[t]$ ,  $[d]$ ,  $[t, d]$  or  $[t, d], b, e$ .

We decline the notion of compact answer defined in Section XXX in four flavors, as follows:

*Definition 5.1.* Let  $G$  be a TG, let  $q$  be a TRPQ and let  $\mathbf{u} \in \mathcal{U}^x$ .

We say that  $\mathbf{u}$  is a *compact answer* to  $q$  over  $G$  (in  $\mathcal{U}^x$ ) if  $\mathbf{u} \in \max_{\sqsubseteq_x} \{ \mathbf{u}' \in \mathcal{U}^x \mid \text{unfold}(\mathbf{u}') = \llbracket q \rrbracket_G \}$

And similarly, we get four decision problems:

COMPACT ANSWER<sup>x</sup>  
**Input:** TG  $G$ , TRPQ  $q$ , tuple  $\mathbf{u} \in \mathcal{U}^x$   
**Decide:**  $\mathbf{u}$  is a compact answer to  $q$  over  $G$  (in  $\mathcal{U}^x$ )

### 5.2 Hardness

### 5.3 Membership

## 6 MINIMIZATION

## 7 SIZE OF COMPACT ANSWERS