Foundations of Classical Mechanics: The Evolution from Newtonian Dynamics to Universal Conservation Laws

The trajectory of classical mechanics represents one of the most significant intellectual migrations in human history, moving from the descriptive, teleological observations of antiquity to the predictive, deterministic frameworks of the modern era. This evolution is not merely a collection of isolated discoveries but a continuous refinement of how the physical world is conceptualized, transitioning from the localized analysis of individual forces to the global principles of conservation. At its inception, the Newtonian paradigm provided the mathematical tools to describe "how" objects move under specific influences. However, as the field matured, the focus shifted toward the underlying symmetries of nature, leading to the realization that certain quantities—momentum and energy—are fundamental constants of the universe. This report provides an exhaustive exploration of these foundations, examining the transition from the vector-based dynamics of Newton to the scalar-based bookkeeping of conservation laws.

The Laws of Motion: The Formalization of Inertia and Force

The bedrock of classical mechanics is formed by the three laws of motion articulated by Isaac Newton, which synthesized the earlier insights of Copernicus, Kepler, and Galileo into a unified theory of dynamics.[1] These laws established a deterministic framework in which the state of a physical system at any given moment can be used to predict its future evolution with absolute certainty, provided all acting forces are known.[3]

Historical Precursors and the Law of Inertia

Prior to the Newtonian revolution, the prevailing Aristotelian view held that motion was a process that required a continuous cause; an object would only remain in motion as long as a force was actively applied to it. This intuitive but flawed perception was challenged by Galileo Galilei through a series of thought experiments involving inclined planes.[2] Galileo observed that if a ball were rolled down one incline and up another, it would attempt to reach its original height regardless of the slope of the second plane. If the second plane were horizontal, Galileo reasoned, the ball would continue to move forever in an attempt to reach that height, provided friction was absent.[4]

Newton formalized this as his First Law, the Law of Inertia. It states that every object continues in its state of rest, or of uniform motion in a straight line, unless it is compelled to change that state by forces impressed upon it.[3] This law redefines the natural state of matter. "Rest" and

"uniform motion" are no longer seen as distinct physical conditions requiring different explanations; rather, they are both states of zero acceleration, occurring whenever the net force on an object is zero.[6]

Inertia, therefore, is the inherent property of matter to resist changes in its velocity. It is quantitatively measured by mass; a more massive object possesses greater inertia and requires a more significant influence to alter its motion.[7] This resistance to change is not a force itself but a fundamental characteristic of existence in a Galilean-Newtonian space.[1]

The Mathematical Heart of Dynamics: Net Force and Acceleration

Newton's Second Law provides the quantitative link between the external influences acting on a body and the resulting change in its motion. It asserts that the acceleration of an object is directly proportional to the net force acting on it and inversely proportional to its mass.[3] Mathematically, this is expressed as:

$$\sum F = ma$$

The "net force" ($\sum F$) is the vector sum of all individual forces—such as gravity, friction, tension, and normal forces—acting on a single object.[3] This law implies that force is the causal agent of acceleration, not velocity. In a system where the net force is zero, the object is in a state of equilibrium, either remaining stationary or moving at a constant velocity.[5]

The implications of the Second Law are profound for the concept of determinism. If the initial position and velocity of every particle in a system are known, and the forces between them can be calculated, the entire future and past of that system can, in principle, be reconstructed.[1] This Newtonian clockwork universe dominated scientific thought for centuries, providing the tools to calculate planetary orbits and engineer the machinery of the industrial age.[3]

Interaction and Symmetry: Newton's Third Law

While the first two laws describe the behavior of a single mass, the Third Law addresses the nature of forces as mutual interactions. It states that whenever one object exerts a force on a second object, the second object exerts an equal and opposite force on the first.[3] These action-reaction pairs are simultaneous and act on different bodies, meaning they never cancel each other out within the context of a single object's dynamics.[7]

This law introduces a fundamental symmetry into classical mechanics. It suggests that forces do not exist in isolation but are manifestations of the connections between entities.[1] As will be

explored in subsequent sections, the Third Law is the dynamical root of the conservation of momentum; because interaction forces within a system are always equal and opposite, the total internal force of any closed system is zero, ensuring that the system's total momentum cannot be changed by its own internal parts.[8]

Newton's Law	Primary Concept	Physical Implication
First Law	Inertia	Objects resist changes in motion; rest and uniform motion are equivalent.
Second Law	$F = ma$	Net force causes acceleration; mass quantifies resistance to that change.
Third Law	Action-Reaction	Forces are mutual interactions; established the basis for symmetry and conservation.

3

The Non-Inertial Perspective: The Turning Car Thought Experiment

The principles of inertia and net force are often best understood through the lens of non-inertial frames of reference—systems that are themselves accelerating. A classic real-world "thought experiment" involves a passenger in a car making a sharp right-hand turn.[4]

As the car begins to turn to the right, the passenger frequently feels a strong "force" pushing them toward the left, seemingly toward the outside of the curve. In popular terminology, this is often called "centrifugal force".[4] However, from the perspective of an inertial observer (someone standing on the sidewalk), no such outward force exists. Instead, the observer sees the passenger's body attempting to obey Newton’s First Law: it wants to continue traveling in a straight line at a constant speed.[7]

The car, being steered by the driver, experiences a real inward force (centripetal force)

provided by the friction between the tires and the road, which pulls the vehicle to the right.[7] The passenger, possessing inertia, continues straight until they come into contact with the car's door or seat. At that point, the car door exerts a real, inward normal force on the passenger, forcing their body to deviate from its straight-line path and follow the car's curve.[7]

The sensation of being "thrown" to the left is an illusion of perception caused by the passenger's inertia resisting the car's acceleration.[10] This "fictitious force" arises only because the passenger is using the accelerating car as their frame of reference. In a truly inertial frame, all forces have an identifiable physical origin—such as contact or gravity—and the "centrifugal" sensation is revealed to be nothing more than the persistence of motion in the absence of a required inward force.[4]

The Work-Energy Theorem: The Bridge from Force to Energy

As classical mechanics moved beyond simple particle dynamics into the study of complex systems, the limitations of Newton's vector-based approach became apparent. Tracking every individual force vector in a multi-body system is computationally exhausting. This led to the development of energy methods, which utilize scalar quantities to describe the state of a system.[1] The Work-Energy Theorem serves as the vital link between these two worlds, showing how forces acting over a displacement result in a change in the system's energy.[12]

The Definition and Calculus of Work

In the context of physics, "Work" (W) is defined with a specific mathematical rigor that distinguishes it from the colloquial use of the term. Work is done when a force acts upon an object to cause a displacement.[14] For a constant force F acting over a displacement d, work is the dot product of the two vectors:

$$W = F \cdot d = Fd\cos\theta$$

where θ is the angle between the force and the direction of motion.[15] This definition highlights that only the component of force parallel to the motion performs work. Forces perpendicular to the displacement, such as the normal force on a block sliding across a floor, do zero work.[13]

When the force is not constant, such as the force exerted by a stretching spring, work must be calculated by integrating the force function over the path of motion:

$$W = \int_{x_i}^{x_f} F(x)dx$$

This integral represents the area under a force-displacement curve.[14] Work is a scalar quantity, measured in Joules (J), and it represents the amount of energy transferred into or out of a system through mechanical means.[13]

Kinetic Energy: The Energy of Motion

The immediate result of performing work on an object (in the absence of other influences) is a change in its speed. By applying Newton's Second Law to the definition of work, one can derive the expression for Kinetic Energy (K).[12] If a net force $F = ma$ acts over a distance d, and we substitute the kinematic relation $v_f^2 = v_i^2 + 2ad$, we find:

$$W_{net} = m \cdot a \cdot d = m\left(\frac{v_f^2 - v_i^2}{2d}\right)d = \frac{1}{2}mv_f^2 - \frac{1}{2}mv_i^2$$

Thus, Kinetic Energy is defined as $K = \frac{1}{2}mv^2$.[12] The Work-Energy Theorem states that the net work done by all forces acting on a particle is equal to the change in that particle's kinetic energy:

$$W_{net} = \Delta K$$

This theorem is a powerful tool because it allows for the calculation of final speeds without needing to know the details of the acceleration or the time elapsed.[13] It simply requires a "before and after" snapshot of the system's energy state.

Potential Energy: Stored Work

While some work immediately increases kinetic energy, other work is "stored" by changing the configuration of a system against a conservative force. This stored energy is called Potential Energy (U).[13] A force is considered conservative if the work it performs on an object is independent of the path taken and depends only on the starting and ending points.[15]

Common examples include:

1. **Gravitational Potential Energy:** When an object of mass m is lifted to a height h, work is done against gravity. The energy is stored in the gravitational field and is calculated as $U_g = mgh$.[13]
2. **Elastic Potential Energy:** When a spring with stiffness k is compressed or stretched by a distance x, the work done is stored in the spring's deformation, calculated as $U_s = \frac{1}{2}kx^2$.[13]

The relationship between work and potential energy is expressed as $W_{cons} = -\Delta U$. This means that as an object falls, gravity does positive work ($+W$), and the potential energy decreases ($-\Delta U$).[13] The "bridge" here is clear: Work is the process of conversion. When you lift a book, your muscles perform positive work, which is converted into the book's potential energy. When you release the book, the gravitational field performs work, converting that potential energy into kinetic energy.[13]

The Scalar Advantage: Total Mechanical Energy

The synthesis of work, kinetic energy, and potential energy leads to the principle of the Conservation of Mechanical Energy. By separating forces into conservative (W_{cons}) and non-conservative (W_{nc}) types, the Work-Energy Theorem can be rewritten:

$$W_{nc} + W_{cons} = \Delta K$$

$$W_{nc} - \Delta U = \Delta K$$

$$W_{nc} = \Delta K + \Delta U$$

If no non-conservative forces (like friction) are present ($W_{nc} = 0$), then the total mechanical energy $E = K + U$ remains constant.[13] This allows physicists to solve complex motion problems—such as a roller coaster navigating a series of loops—by simply equating the total energy at the top of a hill to the total energy at the bottom.[6]

Energy Concept	Mathematical Definition	Role in the "Bridge"
Work (W)	$\int F \cdot dr$	The active agent of energy transfer/transformation.
Kinetic Energy (K)	$\frac{1}{2}mv^2$	The energy possessed by an object due to its motion.
Potential Energy (U)	mgh or $\frac{1}{2}kx^2$	The energy "waiting" to be converted back into motion.
Mechanical Energy (E)	$K + U$	The "ledger total" that remains constant in ideal systems.

13

Conservation Principles: The Universal Bookkeeping Rules

The transition from the specific laws of Newtonian dynamics to the broader principles of conservation represents a shift toward identifying the fundamental invariants of the universe. Conservation laws—specifically those of momentum and energy—are considered "universal bookkeeping" rules because they describe quantities that must always be accounted for, regardless of how complex the internal interactions of a system may be.[18] These laws are not merely empirical observations; they are deeply rooted in the underlying symmetries of space and time.[1]

Conservation of Momentum: The Symmetry of Space

Momentum ($p = mv$) is a vector quantity that captures both the mass and the velocity of an object. The Principle of Conservation of Momentum states that if the net external force acting on a system is zero, the total momentum of that system remains constant.[3]

This principle is a direct consequence of Newton's Third Law. In any interaction between two objects within a system, the forces they exert on each other are equal in magnitude and opposite in direction. Since these forces act for the same amount of time, the impulses they deliver are also equal and opposite ($F_1 \Delta t = -F_2 \Delta t$). This means any momentum gained by one object is exactly equal to the momentum lost by the other.[9]

In terms of "bookkeeping," momentum behaves like a currency that can be exchanged between participants but never created or destroyed from within. This holds true for all types of collisions—whether they are elastic, inelastic, or explosions.[6]

Conservation of Energy: The Ledger of Reality

While momentum is linked to the symmetry of space (translations), the Conservation of Energy is linked to the symmetry of time.[1] It states that the total energy of an isolated system remains constant; energy may change form—from kinetic to potential, thermal, or chemical—but the total amount in the "ledger" never fluctuates.[5]

The "bookkeeping" analogy is particularly apt here. In any physical process, if energy appears to be "missing," physicists do not assume the law has failed. Instead, they look for "hidden accounts"—new forms of energy or previously unobserved particles that have carried the energy away.[19] This principle has led to major discoveries, such as the realization that heat is a form of energy (thermodynamics) and the prediction of the neutrino in particle physics.[19] In the ledger of the universe, energy is the ultimate unit of account.[18]

Collision Example: The Ballistic Pendulum

A ballistic pendulum is a classic experimental setup used to illustrate the interaction of both conservation laws.[24] It consists of a stationary block (the pendulum) into which a fast-moving projectile is fired.

The event occurs in two distinct stages:

1. **The Inelastic Collision:** The projectile strikes and embeds itself in the pendulum block. During this extremely brief interval, the external forces (gravity and tension) have no time to act, so the total momentum is conserved: $mv_i = (m + M)V_f$. However, because the collision is perfectly inelastic, a significant portion of the initial kinetic energy is "lost" to heat and deformation.[8]
2. **The Swing Phase:** Now moving together with velocity V_f, the projectile-block system swings upward to a maximum height h. During this phase, non-conservative forces like air resistance are negligible, and mechanical energy is conserved: the kinetic energy at the bottom of the swing is converted entirely into gravitational potential energy at the peak (

$\frac{1}{2}(m+M)V_f^2 = (m+M)gh$).[8]

By measuring the final height h, one can use the "bookkeeping" of energy to find the velocity V_f immediately after the impact, and then use the "bookkeeping" of momentum to find the original speed of the bullet v_i.[8]

Collision Example: Billiard Ball Dynamics

Billiards provides a nearly perfect laboratory for observing elastic collisions in two dimensions.[26] When two billiard balls of equal mass collide, the total momentum vector is conserved across both the x and y axes.[9]

Mathematically, if ball A strikes stationary ball B:

- **Momentum Conservation:** $v_A = v'_A + v'_B$ (as a vector sum).[26]
- **Kinetic Energy Conservation:** $v_A^2 = (v'_A)^2 + (v'_B)^2$ (as a scalar sum).[26]

In vector geometry, if the sum of two vectors equals a third vector, and the square of their magnitudes follows the Pythagorean theorem ($a^2 + b^2 = c^2$), the two vectors must be perpendicular.[26] This is why, in a non-head-on collision between two pool balls, they will always move off at a 90-degree angle relative to each other.[26] This "slick" physical rule is a direct manifestation of the universe's rigid bookkeeping requirements for both momentum and energy.

Conservation Law	Conserved Quantity	Symmetry Basis	Primary Application
Momentum	$p = mv$	Space Translation	Analyzing collisions and explosions.
Energy	$E = K + U + Q$	Time Translation	Tracking energy transformations across all states.

Angular Momentum	$L = r \times p$	Rotation	Planetary orbits and rotating rigid bodies.

[1]

The Role of Friction and Air Resistance: The Reality of Dissipation

In an idealized "physics world," mechanical energy is a closed loop, endlessly cycling between kinetic and potential forms. However, the macro-scale world is governed by non-conservative forces, primarily friction and air resistance, which act as "drains" on the mechanical energy ledger.[22] These forces do not destroy energy, but they "degrade" it into forms that are no longer useful for performing mechanical work.[19]

The Molecular Mechanism of Friction

Friction is the resistive force that occurs when two surfaces attempt to slide past one another. At a macroscopic level, we see this as a force $f = \mu N$ that opposes motion.[5] At a microscopic level, friction arises from the electromagnetic interactions between the atoms of the two surfaces and the physical interlocking of microscopic irregularities.[28]

When work is done against friction ($W_f = -f \cdot d$), mechanical energy is converted into Thermal Energy (Heat).[6] This occurs because the kinetic energy of the sliding object is transferred into the random, chaotic vibrations of the molecules in the surfaces.[22] Once energy has been randomized in this way, it is extremely difficult to "re-order" it back into directed mechanical motion. This is the essence of irreversibility in classical systems.[22]

Air Resistance and Fluid Drag

Air resistance (or drag) operates on a similar principle. As an object moves through a fluid like air or water, it must physically push molecules out of its way. This requires the object to perform work on the fluid.[16] The object’s kinetic energy is transferred to the air molecules, resulting in turbulence and an increase in the air's temperature.[28] For high-speed objects, such as a falling skydiver, air resistance eventually equals the force of gravity, leading to a "terminal velocity" where the net force is zero and no further acceleration occurs.[7]

The Impossibility of Perpetual Motion Machines

The reality of non-conservative forces provides the scientific basis for why perpetual motion machines—devices that run forever without an external energy source—are physically impossible.[22] These machines are generally categorized by which fundamental law they attempt to violate:

1. **Perpetual Motion of the First Kind:** These machines claim to produce work without any energy input. They violate the First Law of Thermodynamics (Conservation of Energy).[22] Since energy cannot be created from nothing, a machine can only output as much energy as was put in or stored within it.
2. **Perpetual Motion of the Second Kind:** These machines attempt to spontaneously convert heat back into mechanical work with 100% efficiency. While they may not violate the First Law, they violate the Second Law of Thermodynamics (Entropy).[22] Heat only flows spontaneously from hot to cold; extracting work from a single heat reservoir without a temperature gradient is impossible.[22]
3. **Perpetual Motion of the Third Kind:** These machines aim to eliminate friction and all dissipative forces entirely to maintain motion through inertia.[22] While some low-friction systems (like superconductors or objects in deep space) can move for vast periods, they can never be truly "perpetual." Even in a vacuum, objects eventually lose energy through thermal radiation or collisions with stray atoms.[30]

In the "bookkeeping" of the real world, there is always a "tax" paid to the environment in the form of heat. No mechanical system is perfectly efficient, and the inevitable accumulation of entropy ensures that every machine will eventually run down unless energy is continuously replenished.[22]

Force Type	**Energy Transformation**	**Conservative?**	**Impact on Motion**
Gravity	$U_g \leftrightarrow K$	Yes	Recycles energy; allows for orbital stability.
Spring	$U_s \leftrightarrow K$	Yes	Recycles energy; results in oscillation.
Friction	$K \rightarrow Q$ (Heat)	No	Removes mechanical

			energy; stops motion.
Air Resistance	$K \to Q$ (Heat)	No	Opposes velocity; leads to terminal velocity.

6

Synthesis and Concept Map: The Mechanical Hierarchy

The transition from Newtonian Physics to Conservation Laws is not a replacement but an expansion of perspective. To understand the foundations of classical mechanics, one must see how Force, Work, and Energy are inextricably linked in a hierarchical structure. Force provides the instantaneous mechanism, Work describes the process of change over distance, and Energy/Momentum provide the overarching constraints that define what is physically possible.

The Dynamics of Connection

The following "Concept Map" illustrates the logical flow and the mathematical bridges that connect these foundational principles:

- **FORCE (F)** $\to$ *Acts over a distance (d)* $\to$ **WORK (W)**
  - *Equation:* $W = \int F \cdot dr$ [15]
  - *Bridge:* Work is the "Transformer" that changes energy states.[12]
- **WORK (W)** $\to$ *Results in motion* $\to$ **KINETIC ENERGY (K)**
  - *Equation:* $W_{net} = \Delta K = \frac{1}{2}mv_f^2 - \frac{1}{2}mv_i^2$ [12]
  - *Logic:* The Work-Energy Theorem connects net force to changes in speed.
- **WORK (W)** $\to$ *Results in configuration change* $\to$ **POTENTIAL ENERGY (U)**
  - *Equation:* $W_{cons} = -\Delta U$ [13]
  - *Logic:* Conservative forces store work to be used later.
- **CONSERVATION LAWS** $\to$ *Total Ledger* $\to$ **MOMENTUM & ENERGY**
  - *Momentum (p):* $\sum p_i = \sum p_f$ (Vector conservation) [9]

- *Energy (E):* $K_i + U_i + W_{nc} = K_f + U_f$ (Scalar conservation) [13]
- *Logic:* The "Universal Bookkeeping" rules that govern all interactions.[18]

- **DISSIPATIVE FORCES** $\rightarrow$ *Non-Conservative Work* $\rightarrow$ **THERMAL ENERGY (Q)**
  - *Equation:* $\Delta E_{mech} = W_{friction} = Q$ [6]
  - *Logic:* Why mechanical energy "leaks" and perpetual motion is impossible.[22]

Through this hierarchy, classical mechanics provides a complete and elegant description of the physical world. From the simple resistance of a mass (Inertia) to the complex interplay of momentum in a collision, the principles remain consistent. The shift from Newton's "Lexica" to the universal "Bookkeeping" of energy allows physicists to navigate both the ideal vacuum of theory and the dissipative reality of the world we inhabit.[1]

Works cited

1. Foundations of Classical Mechanics [Hardcover ed.] 110848056X, 9781108480567, accessed March 19, 2026, https://dokumen.pub/foundations-of-classical-mechanics-hardcovernbsped-110848056x-9781108480567.html
2. Foundations of Classical Mechanics, accessed March 19, 2026, https://api.pageplace.de/preview/DT0400.9781108658942_A45555032/preview-9781108658942_A45555032.pdf
3. Foundations of Classical Mechanics | PDF - Scribd, accessed March 19, 2026, https://www.scribd.com/document/794949336/Classical-Mechanics-Detailed
4. Fictitious Forces and Non-inertial Frames: The Coriolis Force | Physics, accessed March 19, 2026, https://courses.lumenlearning.com/suny-physics/chapter/6-4-fictitious-forces-and-non-inertial-frames-the-coriolis-force/
5. Physics Study Guide: Motion, Newton's Laws & Energy | Notes - Pearson, accessed March 19, 2026, https://www.pearson.com/channels/physics/study-guides/physics-i-syllabus-mechanics-motion-and-energy
6. Momentum & Energy Concept Map, accessed March 19, 2026, https://bittingerphysics.weebly.com/uploads/5/2/7/5/52759703/concept_map_-_momentum___energy[1].pdf
7. The Physics Classroom Website, accessed March 19, 2026, https://www.physicsclassroom.com/mmedia/circmot/rht.cfm
8. Physics 215 - Experiment 4 Conservation of Energy and Linear Momentum, accessed March 19, 2026, http://www.phy.olemiss.edu/~thomas/weblab/215_lab_items/10_215_con_ener_spr2006.pdf
9. Conservation Of Momentum In Two Dimensions Study Guide - Inspirit VR, accessed March 19, 2026,

https://inspiritvr.com/conservation-of-momentum-in-two-dimensions-study-guide/
10. Circular Motion | Physics | Research Starters - EBSCO, accessed March 19, 2026, https://www.ebsco.com/research-starters/physics/circular-motion
11. Work–Energy Theorem (Chapter 5) - Rigid Body Dynamics, accessed March 19, 2026, https://www.cambridge.org/core/books/rigid-body-dynamics/workenergy-theorem/6D7839FAE6C145CDBE95D67590BCA658
12. 6.4: Work-Energy Theorem - Physics LibreTexts, accessed March 19, 2026, https://phys.libretexts.org/Bookshelves/University_Physics/Physics_(Boundless)/6%3A_Work_and_Energy/6.4%3A_Work-Energy_Theorem
13. Work, Kinetic Energy, and Potential Energy - Mechanics Map, accessed March 19, 2026, https://mechanicsmap.psu.edu/websites/10_work_energy_particle/10-1_work_kinetic_and_potential_energy/work_kinetic_and_potential_energy.html
14. Work, Energy, Power, and Mechanical Systems - MindMap AI, accessed March 19, 2026, https://mindmapai.app/mind-mapping/%D8%A7%D9%84%D9%81%D8%B5%D9%84-4-%D8%A7%D9%84%D8%B4%D8%BA%D9%84-%D9%88%D8%A7%D9%84%D8%B7%D8%A7%D9%82%D8%A9-%D9%88%D8%A7%D9%84%D8%A2%D9%84%D8%A7%D8%AA
15. Chapter 6 Work, Kinetic Energy and Potential Energy, accessed March 19, 2026, https://www2.tntech.edu/leap/murdock/books/v1chap6.pdf
16. Kinetic Energy and the Work-Energy Theorem | Physics - Lumen Learning, accessed March 19, 2026, https://courses.lumenlearning.com/suny-physics/chapter/7-2-kinetic-energy-and-the-work-energy-theorem/
17. Conservation Laws and The Ballistic Pendulum, accessed March 19, 2026, https://www.cerritos.edu/cmera/Physics101/Lab/Conservation_Laws/Conservation_Laws.pdf
18. A survey of the elementary functions of money and its invariant representations, alongside the complementary topic of ownership. - Mark Burgess, accessed March 19, 2026, http://markburgess.org/MoneyBook.pdf
19. Information: currency of life? - PMC, accessed March 19, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC2801531/
20. The Lugon Framework: Informational Foundations of Physical Law; Part VI — The Möbius Gates: Navigating the Curvature Map, accessed March 19, 2026, https://www.preprints.org/frontend/manuscript/4c16227446d241d2e2369c4b2833a1da/download_pub
21. Conservation of momentum - BoxSand, accessed March 19, 2026, https://boxsand.physics.oregonstate.edu/sites/boxsand.physics.oregonstate.edu/files/CM.L1.sols_.pdf
22. Perpetual motion - Wikipedia, accessed March 19, 2026, https://en.wikipedia.org/wiki/Perpetual_motion
23. The New Quantum Catastrophe: How Decoherence Revived the Infinity Planck

Banished in the Ultraviolet Crisis | by Bill Giannakopoulos | Medium, accessed March 19, 2026, https://medium.com/@bill.giannakopoulos/the-new-quantum-catastrophe-how-decoherence-revived-the-infinity-planck-banished-in-the-ad8e5fc7389f
24. 24.15 -- Ballistic pendulums - UCSB Physics, accessed March 19, 2026, https://web.physics.ucsb.edu/~lecturedemonstrations/Composer/Pages/24.15.html
25. The Ballistic Pendulum: Conservation of Momentum & Conservation of Energy - SUNY Geneseo, accessed March 19, 2026, https://tildesites.geneseo.edu/~pogo/Analyt1Labs/Labphotos/Ballistic.pdf
26. Physics Of Billiards - Real World Physics Problems, accessed March 19, 2026, https://www.real-world-physics-problems.com/physics-of-billiards.html
27. Conservation of Momentum in Two Dimensions | CK-12 Foundation, accessed March 19, 2026, https://www.ck12.org/book/ck-12-physics-concepts-intermediate/r5/section/7.3/
28. Perpetual Motion Machines — Why They Don't Work? | by Devansh Mittal | Intuitive Physics, accessed March 19, 2026, https://medium.com/intuitive-physics/perpetual-motion-machines-why-they-don't-work-1e192d2c0f62
29. How is it possible to know a proposed perpetual motion machine won't work without even looking at it?, accessed March 19, 2026, https://physics.stackexchange.com/questions/826203/how-is-it-possible-to-know-a-proposed-perpetual-motion-machine-wont-work-withou
30. Why are perpetual motion machines impossible? - Physics Stack Exchange, accessed March 19, 2026, https://physics.stackexchange.com/questions/520113/why-are-perpetual-motion-machines-impossible
31. If perpetual motion is impossible, then how do objects in space continue to move forever?, accessed March 19, 2026, https://www.reddit.com/r/AskPhysics/comments/1ctlot4/if_perpetual_motion_is_impossible_then_how_do/