

Computation of neutral K and D meson mass using non-relativistic lattice QCD.

Oscar Simpson
Supervisor: Dr. B. Chakraborty

May 2021

We begin with a recap of QCD[1, 2].

$$\mathcal{L} = \sum_f \psi_f^\dagger (i\not{D} - m_f) \psi_f - \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu}$$

We begin with a recap of QCD[1, 2].

$$\mathcal{L} = \sum_f \psi_f^\dagger (i\not{D} - m_f) \psi_f - \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu}$$

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a - g_s f_{abc} A_\mu^b A_\nu^c$$

Now we move on to a recap of lattice QCD[3, 2].

$$\langle \mathcal{O} \rangle = \frac{1}{\mathcal{Z}} \int \mathcal{D}\psi \mathcal{O} \exp\left(i \int d^4x \mathcal{L}(\psi, \partial\psi)\right)$$
$$\mathcal{Z} = \int \mathcal{D}\psi \exp\left(i \int d^4x \mathcal{L}(\psi, \partial\psi)\right)$$

Lattice QCD Recap




Figure: An example of a discretisation. Red crosses indicate quark fields, and blue dots indicate gauge fields.

- Mesons containing heavy quarks such as b or c require a small lattice size ($a < 0.45$ fm).
- This is computationally expensive if computed fully relativistically.

- Mesons containing heavy quarks such as b or c require a small lattice size ($a < 0.45 \text{ fm}$).
- This is computationally expensive if computed fully relativistically.
- Expand action in $\frac{1}{m_b}$.
- Also expanding in $\frac{v^2}{c^2}$ gives the Schrödinger equation[2],

$$\mathcal{L}_0 = \psi^\dagger(x) \left(iD_t + \frac{\mathbf{D}^2}{2m_b} \right) \psi(x).$$

- This action is only first order in the above expansions. The datasets used for this project are computed using an equivalent third order expansion.

- Data was provided by Dr. B. Chakraborty.
- Data has been produced using isotropic lattice NRQCD[4].

- Data was provided by Dr. B. Chakraborty.
- Data has been produced using isotropic lattice NRQCD[4].

Label	Spacing a (fm)	Lattice size (time)	m_l/m_s
Very Coarse	0.1509	48	0.036
Coarse	0.124 04	64	0.2
Fine	0.090 23	96	0.2

Table: Parameters of the data used in the main part of this project. Corresponding to sets 1, 5, 6 from table I in [4] respectively.

- Data was provided by Dr. B. Chakraborty.
- Data has been produced using isotropic lattice NRQCD[4].

Label	Spacing a (fm)	Lattice size (time)	m_l/m_s
Very Coarse	0.1509	48	0.036
Coarse	0.124 04	64	0.2
Fine	0.090 23	96	0.2

Table: Parameters of the data used in the main part of this project. Corresponding to sets 1, 5, 6 from table I in [4] respectively.

- Finite lattice leads to discretisation errors: must take continuum limit.

- Data was provided by Dr. B. Chakraborty.
- Data has been produced using isotropic lattice NRQCD[4].

Label	Spacing a (fm)	Lattice size (time)	m_l/m_s
Very Coarse	0.1509	48	0.036
Coarse	0.124 04	64	0.2
Fine	0.090 23	96	0.2

Table: Parameters of the data used in the main part of this project. Corresponding to sets 1, 5, 6 from table I in [4] respectively.

- Finite lattice leads to discretisation errors: must take continuum limit.
- Computations requirements mean we must use unphysical light quark masses: must take chiral limit.

Statistical Analysis: Jackknife averaging

- Given a sample of n values $x_i \sim F$, we want to find a way to construct best estimators for F .

Statistical Analysis: Jackknife averaging

- Given a sample of n values $x_i \sim F$, we want to find a way to construct best estimators for F .
- For example, estimates for the mean:
 - Construct $\bar{x}_i = \sum_{j \neq i} \frac{x_j}{n-1}$.
 - $\bar{x} = \sum_i \frac{\bar{x}_i}{n}$ is in fact an unbiased estimator for the mean of F .

Statistical Analysis: Jackknife averaging

- Given a sample of n values $x_i \sim F$, we want to find a way to construct best estimators for F .
- For example, estimates for the mean:
 - Construct $\bar{x}_i = \sum_{j \neq i} \frac{x_j}{n-1}$.
 - $\bar{x} = \sum_i \frac{\bar{x}_i}{n}$ is in fact an unbiased estimator for the mean of F .
- A more complex variant of this procedure can be used to find an unbiased estimator for any parameter of F [5].

Statistical Analysis: Bayesian analysis

- We use the `corrfitter`[6] Python package.

Statistical Analysis: Bayesian analysis

- We use the `corrfitter`[6] Python package.
- Perform least-squares regression on 2-point correlator data.

$$G(t) = \sum_n a_n^2 (e^{-E_n t} + e^{-E_n (T-t)}) + (-1)^t \sum_n a_{o_n}^2 (e^{-E_{o_n} t} + e^{-E_{o_n} (T-t)})$$

Preliminary Study: effective mass

$$G(t) = \sum_n a_n^2 (e^{-E_n t} + e^{-E_n (T-t)}) + (-1)^t \sum_n a_{o_n}^2 (e^{-E_{o_n} t} + e^{-E_{o_n} (T-t)})$$

Assuming only the first term dominates, and there is little contribution from the oscillatory terms, then we have:

$$m_{\text{eff}}(t) = \log \left(\frac{G(t)}{G(t+1)} \right).$$

Preliminary Study: effective mass

$$m_{\text{eff}}(t) = \log\left(\frac{G(t)}{G(t+1)}\right)$$




Figure: Jackknife averaged correlation data for D^0 meson on a fine lattice.

Preliminary Study: effective mass

$$m_{\text{eff}}(t) = \log\left(\frac{G(t)}{G(t+1)}\right)$$




Figure: Estimate for effective mass of D^0 meson.

Data Analysis: fine lattice

- Fine dataset has $a \approx 0.09$ fm and $m_l \approx 0.2m_s$.
- K meson clearly converges for $n \geq 4$.




Figure: Results of fitting K meson dataset on a fine lattice.

Data Analysis: fine lattice

Combined fit quality plot shows that the fitting converged for all t_{\min} and $n \geq 4$.




Figure: Fitting quality for fine lattice.

Data Analysis: coarse lattice

- Coarse dataset has $a \approx 0.12$ fm and $m_l \approx 0.2m_s$.
- K meson converges for $n \geq 3$.




Figure: Results of fitting K meson dataset on a coarse lattice.

Data Analysis: coarse lattice

Combined fit quality plot shows that the fitting converged for all t_{\min} and varying n .




Figure: Fitting quality for coarse lattice.

Data Analysis: very coarse lattice

- Fine dataset has $a \approx 0.15$ fm and $m_l \approx 0.036m_s$.
- K meson appears to converge for $n \geq 2$.




Figure: Results of fitting K meson dataset on a very coarse lattice.

Data Analysis: very coarse lattice

Combined fit quality plot shows that the fitting converged for a very limited selection of t_{\min} and n .




Figure: Fitting quality for very coarse lattice.

Results

For consistency between plots, $t_{\min} = 6$ for D mesons, and $n = 4$ for all results, have been selected.

Label	a (fm)	m_l/m_s	E_0 (MeV)		a_0	
			K	D	K	D
Very Coarse	0.15	0.036	495.92(08)	1887.3(51)	-0.281 01(11)	-0.2286(65)
Coarse	0.12	0.2	528.28(15)	1893.0(15)	+0.608 53(37)	+0.1880(12)
Fine	0.09	0.2	530.16(24)	1889.3(11)	+0.139 78(12)	+0.1242(07)

Table: Results from fitting all datasets. Uncertainties given in parentheses are statistical.

These are reasonable mass values as literature[7] quotes $m_K \approx 498$ MeV and $m_D \approx 1865$ MeV.

- Full extrapolation[8] requires 7 pairs of parameters.
- As we have very few datasets, we restrict to first order in each parameter.

$$m = m_{\text{phys}} \left(1 + c_\delta \frac{m_l}{m_s} \right) \left(1 + c_{a^2} a^2 \right)$$

Chiral/Continuum Extrapolation

The fit has $\chi^2/\text{dof} = 0.23$ so we have a good quality fit, and importantly not an overfit.

Label	E_0 (MeV)		Predicted E_0 (MeV)	
	K	D	K	D
Very Coarse	495.92(08)	1887.3(51)	495.92(08)	1887.3(51)
Coarse	528.28(15)	1893.0(15)	528.31(25)	1892.8(15)
Fine	530.16(24)	1889.3(11)	530.13(24)	1889.4(11)
Chiral/Continuum limit	-	-	494.5(11)	1874.6(88)

Table: Comparison of observed and predicted mass values, including the extrapolated chiral/continuum limit values.

The extrapolated values are consistent with results from the literature[7], $m_{K^0} = 497.611(13)$ MeV and $m_{D^0} = 1864.83(5)$ MeV to within 3σ of our statistical uncertainty.

Remaining Work

- Compute partial decay widths, using amplitudes a_0 .

Bibliography

- [1] G. Shaw F. Mandl. *Quantum Field Theory*. John Wiley & Sons, 2010.
- [2] Brian Colquhoun. "Bottomonium and B Physics with Lattice NRQCD b Quarks". University of Glasgow, 2015.
- [3] R. P. Feynman. "Space-Time Approach to Non-Relativistic Quantum Mechanics". In: *Rev. Mod. Phys.* 20 (2 Apr. 1948), pp. 367–387. DOI: 10.1103/RevModPhys.20.367.
- [4] Bipasha Chakraborty et al. "Improved V_{CS} determination using precise lattice QCD form factors for $D \rightarrow K l \nu$ ". In: *arXiv e-prints* (2021), arXiv-2104.
- [5] Bradley Efron. *The jackknife, the bootstrap and other resampling plans*. SIAM, 1982.
- [6] G Peter Lepage. *corrfitter*. Version 8.1.1. 2020. URL: <https://github.com/gplepage/corrfitter>.
- [7] Particle Data Group et al. "Review of Particle Physics". In: *Progress of Theoretical and Experimental Physics* 2020.8 (Aug. 2020). 083C01. ISSN: 2050-3911. DOI: 10.1093/ptep/ptaa104. eprint: https://academic.oup.com/ptep/article-pdf/2020/8/083C01/34673740/rpp2020-vol2-2015-2092_18.pdf. URL: <https://doi.org/10.1093/ptep/ptaa104>.
- [8] Bipasha Chakraborty et al. "Nonperturbative comparison of clover and HISQ quarks in lattice QCD and the properties of the phi meson". In: *arXiv preprint arXiv:1703.05552* (2017).