

# Obtaining $K$ and $D$ meson properties from lattice QCD

Oscar Simpson  
Supervisor: Dr. B. Chakraborty

May 2021

We begin with a recap of QCD[1, 2].

$$\mathcal{L} = \sum_f \psi_f^\dagger (i\not{D} - m_f) \psi_f - \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu}$$

We begin with a recap of QCD[1, 2].


$$\mathcal{L} = \sum_f \psi_f^\dagger (i\not{D} - m_f) \psi_f - \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu}$$

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a - g_s f_{abc} A_\mu^b A_\nu^c$$

Now we move on to a recap of lattice QCD[3, 2].

$$\langle \mathcal{O} \rangle = \frac{1}{Z} \int \mathcal{D}\psi \mathcal{O} \exp\left(i \int d^4x \mathcal{L}(\psi, \partial\psi)\right)$$
$$Z = \int \mathcal{D}\psi \exp\left(i \int d^4x \mathcal{L}(\psi, \partial\psi)\right)$$

# Lattice QCD Recap



**Figure:** An example of a discretisation. Red crosses indicate quark fields, and blue dots indicate gauge fields.

# Highly Improved Staggered Quark

- Fermions placed on a finite lattice give "doubled" fermions[4].

$$M^{-1} = \frac{-i\gamma_\mu \sin(ap_\mu)/a + m}{\sum_{\mu=0}^3 \sin^2(ap_\mu)/a^2 + m^2} \quad (1)$$

- These are the reflections of the fermion mass within the Brillouin zone in each of the 4 spacetime dimensions, which gives  $16 = 2^4$  extra modes, called "tastes".
- To remove this problem, Wilson[5] used a construction to give these tastes infinite mass, such that they decouple in the continuum limit.
- A procedure called "staggering" transformation[6] allows us to bring the system down to only 4 doublers.
- With some further optimizations this gives the Highly Improved Staggered Quark (HISQ) action, which has very small discretisation errors.

- Data was provided by Dr. B. Chakraborty.
- Data has been produced using isotropic lattice NRQCD[7].

- Data was provided by Dr. B. Chakraborty.
- Data has been produced using isotropic lattice NRQCD[7].

Label	Spacing $a$ (fm)	Lattice size (time)	$m_l/m_s$
Very Coarse	0.1509	48	0.036
Coarse	0.124 04	64	0.2
Fine	0.090 23	96	0.2

**Table:** Parameters of the data used in the main part of this project. Corresponding to sets 1, 5, 6 from table I in [7] respectively.



- Data was provided by Dr. B. Chakraborty.
- Data has been produced using isotropic lattice NRQCD[7].

Label	Spacing $a$ (fm)	Lattice size (time)	$m_l/m_s$
Very Coarse	0.1509	48	0.036
Coarse	0.124 04	64	0.2
Fine	0.090 23	96	0.2

**Table:** Parameters of the data used in the main part of this project. Corresponding to sets 1, 5, 6 from table I in [7] respectively.

- Finite lattice leads to discretisation errors: must take continuum limit.

- Data was provided by Dr. B. Chakraborty.
- Data has been produced using isotropic lattice NRQCD[7].

Label	Spacing $a$ (fm)	Lattice size (time)	$m_l/m_s$
Very Coarse	0.1509	48	0.036
Coarse	0.124 04	64	0.2
Fine	0.090 23	96	0.2

**Table:** Parameters of the data used in the main part of this project. Corresponding to sets 1, 5, 6 from table I in [7] respectively.

- Finite lattice leads to discretisation errors: must take continuum limit.
- Computations requirements mean we must use unphysical light quark masses: must take chiral limit.

# Statistical Analysis: Jackknife averaging

- Given a sample of  $n$  values  $x_i \sim F$ , we want to find a way to construct best estimators for  $F$ .

# Statistical Analysis: Jackknife averaging

- Given a sample of  $n$  values  $x_i \sim F$ , we want to find a way to construct best estimators for  $F$ .
- For example, estimates for the mean:
  - Construct  $\bar{x}_i = \sum_{j \neq i} \frac{x_j}{n-1}$ .
  - $\bar{x} = \sum_i \frac{\bar{x}_i}{n}$  is in fact an unbiased estimator for the mean of  $F$ .

# Statistical Analysis: Jackknife averaging

- Given a sample of  $n$  values  $x_i \sim F$ , we want to find a way to construct best estimators for  $F$ .
- For example, estimates for the mean:
  - Construct  $\bar{x}_i = \sum_{j \neq i} \frac{x_j}{n-1}$ .
  - $\bar{x} = \sum_i \frac{\bar{x}_i}{n}$  is in fact an unbiased estimator for the mean of  $F$ .
- A more complex variant of this procedure can be used to find an unbiased estimator for any parameter of  $F$ [8].

# Statistical Analysis: Bayesian analysis

- We use the `corrfinder`[9] Python package.

# Statistical Analysis: Bayesian analysis

- We use the `corrfitter`[9] Python package.
- Perform least-squares regression on 2-point correlator data.

$$G(t) = \sum_n a_n^2 (e^{-E_n t} + e^{-E_n (T-t)}) + (-1)^t \sum_n a_{o_n}^2 (e^{-E_{o_n} t} + e^{-E_{o_n} (T-t)})$$

# Preliminary Study: effective mass

$$G(t) = \sum_n a_n^2 (e^{-E_n t} + e^{-E_n (T-t)}) + (-1)^t \sum_n a_{o_n}^2 (e^{-E_{o_n} t} + e^{-E_{o_n} (T-t)})$$


Assuming only the first term dominates, and there is little contribution from the oscillatory terms, then we have:

$$m_{\text{eff}}(t) = \log \left( \frac{G(t)}{G(t+1)} \right).$$



# Preliminary Study: effective mass


$$m_{\text{eff}}(t) = \log\left(\frac{G(t)}{G(t+1)}\right)$$



**Figure:** Jackknife averaged correlation data for  $D^0$  meson on a fine lattice.

# Preliminary Study: effective mass


$$m_{\text{eff}}(t) = \log\left(\frac{G(t)}{G(t+1)}\right)$$



**Figure:** Estimate for effective mass of  $D^0$  meson.

# Data Analysis: fine lattice


- Fine dataset has  $a \approx 0.09$  fm and  $m_l \approx 0.2m_s$ .
- K meson clearly converges for  $n \geq 4$ .



**Figure:** Results of fitting K meson dataset on a fine lattice.

# Data Analysis: fine lattice


Combined fit quality plot shows that the fitting converged for all  $t_{\min}$  and  $n \geq 4$ .



**Figure:** Fitting quality for fine lattice.

# Data Analysis: coarse lattice


- Coarse dataset has  $a \approx 0.12$  fm and  $m_l \approx 0.2m_s$ .
- K meson converges for  $n \geq 3$ .



**Figure:** Results of fitting K meson dataset on a coarse lattice.

# Data Analysis: coarse lattice


Combined fit quality plot shows that the fitting converged for all  $t_{\min}$  and varying  $n$ .



**Figure:** Fitting quality for coarse lattice.

# Data Analysis: very coarse lattice


- Fine dataset has  $a \approx 0.15$  fm and  $m_l \approx 0.036m_s$ .
- K meson appears to converge for  $n \geq 2$ .



**Figure:** Results of fitting K meson dataset on a very coarse lattice.

# Data Analysis: very coarse lattice

Combined fit quality plot shows that the fitting converged for a very limited selection of  $t_{\min}$  and  $n$ .



**Figure:** Fitting quality for very coarse lattice.



# Results

For consistency between plots,  $t_{\min} = 6$  for  $D$  mesons, and  $n = 4$  for all results, have been selected.

Label	$a$ (fm)	$m_l/m_s$	$E_0$ (MeV)		$ a_0 $	
			K	D	K	D
Very Coarse	0.15	0.036	495.92(08)	1887.3(51)	0.281 01(11)	0.2286(65)
Coarse	0.12	0.2	528.28(15)	1893.0(15)	0.608 53(37)	0.1880(12)
Fine	0.09	0.2	530.16(24)	1889.3(11)	0.139 78(12)	0.1242(07)

**Table:** Results from fitting all datasets. Uncertainties given in parentheses are statistical.

These are reasonable mass values as literature[10] quotes  $m_K \approx 498$  MeV and  $m_D \approx 1865$  MeV.

- Full extrapolation[11] requires 7 pairs of parameters.
- As we have very few datasets, we restrict to first order in each parameter.

$$m = m_{\text{phys}} \left( 1 + c_\delta \frac{m_l}{m_s} \right) \left( 1 + c_{a^2} a^2 \right)$$

# Chiral/Continuum Extrapolation

The fit has  $\chi^2/\text{dof} = 0.23$  so we have a good quality fit, and importantly not an overfit.

Label	$a$ (fm)	$m_l/m_s$	$E_0$ (MeV)		$ a_0 $	
			K	D	K	D
Very Coarse	0.15	0.036	495.92(08)	1887.3(51)	0.281 01(11)	0.2286(65)
Coarse	0.12	0.2	528.28(15)	1893.0(15)	0.608 53(37)	0.1880(12)
Fine	0.09	0.2	530.16(24)	1889.3(11)	0.139 78(12)	0.1242(07)

**Table:** Comparison of observed and predicted mass values, including the extrapolated chiral/continuum limit values.

The extrapolated values are consistent with results from the literature[10],  $m_{K^0} = 497.611(13)$  MeV and  $m_{D^0} = 1864.83(5)$  MeV to within  $3\sigma$  of our statistical uncertainty.

# Remaining Work

- Compute partial decay widths, using amplitudes  $a_0$ .

# Bibliography

- [1] G. Shaw F. Mandl. *Quantum Field Theory*. John Wiley & Sons, 2010.
- [2] Brian Colquhoun. "Bottomonium and B Physics with Lattice NRQCD b Quarks". University of Glasgow, 2015.
- [3] R. P. Feynman. "Space-Time Approach to Non-Relativistic Quantum Mechanics". In: *Rev. Mod. Phys.* 20 (2 Apr. 1948), pp. 367–387. DOI: [10.1103/RevModPhys.20.367](https://doi.org/10.1103/RevModPhys.20.367).
- [4] Bipasha Chakraborty. "Precision tests of the standard model using lattice QCD". University of Glasgow, 2016.
- [5] Kenneth G. Wilson. "Confinement of quarks". In: *Phys. Rev. D* 10 (8 Oct. 1974), pp. 2445–2459. DOI: [10.1103/PhysRevD.10.2445](https://doi.org/10.1103/PhysRevD.10.2445).
- [6] Quentin Mason et al. "High-precision determination of the light-quark masses from realistic lattice QCD". In: *Physical Review D* 73.11 (2006), p. 114501.
- [7] Bipasha Chakraborty et al. "Improved  $V_{cs}$  determination using precise lattice QCD form factors for  $D \rightarrow K l \nu$ ". In: *arXiv e-prints* (2021), arXiv:2104.00001.
- [8] Bradley Efron. *The jackknife, the bootstrap and other resampling plans*. SIAM, 1982.
- [9] G Peter Lepage. *corrfitter*. Version 8.1.1. 2020. URL: <https://github.com/gplepage/corrfitter>.
- [10] Particle Data Group et al. "Review of Particle Physics". In: *Progress of Theoretical and Experimental Physics* 2020.8 (Aug. 2020). 083C01. ISSN: 2050-3911. DOI: [10.1093/ptep/ptaa104](https://doi.org/10.1093/ptep/ptaa104). eprint: [https://academic.oup.com/ptep/article-pdf/2020/8/083C01/34673740/rpp2020-vol2-2015-2092\\_18.pdf](https://academic.oup.com/ptep/article-pdf/2020/8/083C01/34673740/rpp2020-vol2-2015-2092_18.pdf). URL: <https://doi.org/10.1093/ptep/ptaa104>.
- [11] Bipasha Chakraborty et al. "Nonperturbative comparison of clover and HISQ quarks in lattice QCD and the properties of the phi meson". In: *arXiv preprint arXiv:1703.05552* (2017).