

Obtaining K and D meson properties from lattice QCD

Oscar Simpson
Supervisor: Dr. B. Chakraborty

May 2021

We begin with a recap of QCD[1, 2].

$$\mathcal{L} = \sum_f \psi_f^\dagger (i\not{D} - m_f) \psi_f - \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu}$$

We begin with a recap of QCD[1, 2].

$$\mathcal{L} = \sum_f \psi_f^\dagger (i\not{D} - m_f) \psi_f - \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu}$$

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a - g_s f_{abc} A_\mu^b A_\nu^c$$

Now we move on to a recap of lattice QCD[3, 2].

$$\langle \mathcal{O} \rangle = \frac{1}{\mathcal{Z}} \int \mathcal{D}\psi \, \mathcal{O} \exp\left(i \int d^4x \, \mathcal{L}(\psi, \partial\psi)\right)$$
$$\mathcal{Z} = \int \mathcal{D}\psi \, \exp\left(i \int d^4x \, \mathcal{L}(\psi, \partial\psi)\right)$$

Lattice QCD Recap




Figure: An example of a discretisation. Red crosses indicate quark fields, and blue dots indicate gauge fields.

Highly Improved Staggered Quark

- Fermions placed on a finite lattice give "doubled" fermions[4].

$$M^{-1} = \frac{-i\gamma_\mu \sin(ap_\mu)/a + m}{\sum_{\mu=0}^3 \sin^2(ap_\mu)/a^2 + m^2} \quad (1)$$

- These are the reflections of the fermion mass within the Brillouin zone in each of the 4 spacetime dimensions, which gives $16 = 2^4$ extra modes, called "tastes".
- To remove this problem, Wilson[5] used a construction to give these tastes infinite mass, such that they decouple in the continuum limit.
- A procedure called "staggering" transformation[6] allows us to bring the system down to only 4 doublers.
- With some further optimizations this gives the Highly Improved Staggered Quark (HISQ) action, which has very small discretisation errors.

- Data was provided by Dr. B. Chakraborty.
- This data has been produced by using relativistic HISQ action for both valence and sea quarks on publicly available gauge configurations generated by MILC collaboration. The configurations have sea quark effects from up/down (light), strange, and charm quarks.[7].

- Data was provided by Dr. B. Chakraborty.
- This data has been produced by using relativistic HISQ action for both valence and sea quarks on publicly available gauge configurations generated by MILC collaboration. The configurations have sea quark effects from up/down (light), strange, and charm quarks.[7].

Label	Spacing a (fm)	Lattice size (time)	m_l/m_s
Very Coarse	0.1509	48	0.036
Coarse	0.124 04	64	0.2
Fine	0.090 23	96	0.2

Table: Parameters of the data used in the main part of this project. Corresponding to sets 1, 5, 6 from table I in [7] respectively.

- Data was provided by Dr. B. Chakraborty.
- This data has been produced by using relativistic HISQ action for both valence and sea quarks on publicly available gauge configurations generated by MILC collaboration. The configurations have sea quark effects from up/down (light), strange, and charm quarks.[7].

Label	Spacing a (fm)	Lattice size (time)	m_l/m_s
Very Coarse	0.1509	48	0.036
Coarse	0.124 04	64	0.2
Fine	0.090 23	96	0.2

Table: Parameters of the data used in the main part of this project. Corresponding to sets 1, 5, 6 from table I in [7] respectively.

- Finite lattice leads to discretisation errors: must take continuum limit.

- Data was provided by Dr. B. Chakraborty.
- This data has been produced by using relativistic HISQ action for both valence and sea quarks on publicly available gauge configurations generated by MILC collaboration. The configurations have sea quark effects from up/down (light), strange, and charm quarks.[7].

Label	Spacing a (fm)	Lattice size (time)	m_l/m_s
Very Coarse	0.1509	48	0.036
Coarse	0.124 04	64	0.2
Fine	0.090 23	96	0.2

Table: Parameters of the data used in the main part of this project. Corresponding to sets 1, 5, 6 from table I in [7] respectively.

- Finite lattice leads to discretisation errors: must take continuum limit.
- Computations requirements mean we must use unphysical light quark masses: must take chiral limit.

Statistical Analysis: Jackknife averaging

- Given a sample of n values $x_i \sim F$, we want to find a way to construct best estimators for F .

Statistical Analysis: Jackknife averaging

- Given a sample of n values $x_i \sim F$, we want to find a way to construct best estimators for F .
- For example, estimates for the mean:
 - Construct $\bar{x}_i = \sum_{j \neq i} \frac{x_j}{n-1}$.
 - $\bar{x} = \sum_i \frac{\bar{x}_i}{n}$ is in fact an unbiased estimator for the mean of F .

Statistical Analysis: Jackknife averaging

- Given a sample of n values $x_i \sim F$, we want to find a way to construct best estimators for F .
- For example, estimates for the mean:
 - Construct $\bar{x}_i = \sum_{j \neq i} \frac{x_j}{n-1}$.
 - $\bar{x} = \sum_i \frac{\bar{x}_i}{n}$ is in fact an unbiased estimator for the mean of F .
- A more complex variant of this procedure can be used to find an unbiased estimator for any parameter of F [8].

Statistical Analysis: Bayesian analysis

- We use the `corrfinder`[9] Python package.

Statistical Analysis: Bayesian analysis

- We use the `corrfitter`[9] Python package.
- Perform least-squares regression on 2-point correlator data.

$$G(t) = \sum_n a_n^2 (e^{-E_n t} + e^{-E_n (T-t)}) + (-1)^t \sum_n a_{o_n}^2 (e^{-E_{o_n} t} + e^{-E_{o_n} (T-t)})$$

Preliminary Study: effective mass

$$G(t) = \sum_n a_n^2 (e^{-E_n t} + e^{-E_n (T-t)}) + (-1)^t \sum_n a_{o_n}^2 (e^{-E_{o_n} t} + e^{-E_{o_n} (T-t)})$$

Assuming only the first term dominates, and there is little contribution from the oscillatory terms, then we have:

$$m_{\text{eff}}(t) = \log \left(\frac{G(t)}{G(t+1)} \right).$$

Preliminary Study: effective mass

$$m_{\text{eff}}(t) = \log\left(\frac{G(t)}{G(t+1)}\right)$$




Figure: Jackknife averaged correlation data for D^0 meson on a fine lattice. This is for a slightly different set of parameters than otherwise discussed here, however it is sufficient for this discussion.

Preliminary Study: effective mass

$$m_{\text{eff}}(t) = \log\left(\frac{G(t)}{G(t+1)}\right)$$




Figure: Estimate for effective mass of D^0 meson.

Data Analysis: fine lattice

- Fine dataset has $a \approx 0.09$ fm and $m_l \approx 0.2m_s$.
- K meson clearly converges for $n \geq 4$.




Figure: Results of fitting K meson dataset on a fine lattice.

Data Analysis: fine lattice

Combined fit quality plot shows that the fitting converged for all t_{\min} and $n \geq 4$.




Figure: Fitting quality for fine lattice.

Data Analysis: coarse lattice

- Coarse dataset has $a \approx 0.12$ fm and $m_l \approx 0.2m_s$.
- K meson converges for $n \geq 3$.




Figure: Results of fitting K meson dataset on a coarse lattice.

Data Analysis: coarse lattice

Combined fit quality plot shows that the fitting converged for all t_{\min} and varying n .




Figure: Fitting quality for coarse lattice.

Data Analysis: very coarse lattice

- Fine dataset has $a \approx 0.15$ fm and $m_l \approx 0.036m_s$.
- K meson appears to converge for $n \geq 2$.




Figure: Results of fitting K meson dataset on a very coarse lattice.

Data Analysis: very coarse lattice

Combined fit quality plot shows that the fitting converged for a very limited selection of t_{\min} and n .




Figure: Fitting quality for very coarse lattice.

Results

For consistency between plots, $t_{\min} = 6$ for D mesons, and $n = 4$ for all results, have been selected.

Label	a (fm)	m_l/m_s	E_0 (MeV)		$ a_0 $	
			K	D	K	D
Very Coarse	0.15	0.036	495.92(08)	1887.3(51)	0.281 01(11)	0.2286(65)
Coarse	0.12	0.2	528.28(15)	1893.0(15)	0.608 53(37)	0.1880(12)
Fine	0.09	0.2	530.16(24)	1889.3(11)	0.139 78(12)	0.1242(07)

Table: Results from fitting all datasets. Uncertainties given in parentheses are statistical.

These are reasonable mass values as literature[10] quotes $m_K \approx 498$ MeV and $m_D \approx 1865$ MeV.

- Full extrapolation[11] requires 7 pairs of parameters.
- As we have very few datasets, we restrict to first order in each parameter.

$$m = m_{\text{phys}} \left(1 + c_\delta \frac{m_l}{m_s} \right) \left(1 + c_{a^2} a^2 \right)$$

Chiral/Continuum Extrapolation

The fit has $\chi^2/\text{dof} = 0.23$ so we have a good quality fit, and importantly not an overfit.

Label	a (fm)	m_l/m_s	E_0 (MeV)		$ a_0 $	
			K	D	K	D
Very Coarse	0.15	0.036	495.92(08)	1887.3(51)	0.281 01(11)	0.2286(65)
Coarse	0.12	0.2	528.28(15)	1893.0(15)	0.608 53(37)	0.1880(12)
Fine	0.09	0.2	530.16(24)	1889.3(11)	0.139 78(12)	0.1242(07)

Table: Comparison of observed and predicted mass values, including the extrapolated chiral/continuum limit values.

The extrapolated values are consistent with results from the literature[10], $m_{K^0} = 497.611(13)$ MeV and $m_{D^0} = 1864.83(5)$ MeV to within 3σ of our statistical uncertainty.

Remaining Work

- Compute partial decay widths, using amplitudes a_0 .

Bibliography

- [1] G. Shaw F. Mandl. *Quantum Field Theory*. John Wiley & Sons, 2010.
- [2] Brian Colquhoun. "Bottomonium and B Physics with Lattice NRQCD b Quarks". University of Glasgow, 2015.
- [3] R. P. Feynman. "Space-Time Approach to Non-Relativistic Quantum Mechanics". In: *Rev. Mod. Phys.* 20 (2 Apr. 1948), pp. 367–387. DOI: 10.1103/RevModPhys.20.367.
- [4] Bipasha Chakraborty. "Precision tests of the standard model using lattice QCD". University of Glasgow, 2016.
- [5] Kenneth G. Wilson. "Confinement of quarks". In: *Phys. Rev. D* 10 (8 Oct. 1974), pp. 2445–2459. DOI: 10.1103/PhysRevD.10.2445.
- [6] Quentin Mason et al. "High-precision determination of the light-quark masses from realistic lattice QCD". In: *Physical Review D* 73.11 (2006), p. 114501.
- [7] Bipasha Chakraborty et al. "Improved V_{cs} determination using precise lattice QCD form factors for $D \rightarrow K l \nu$ ". In: *arXiv e-prints* (2021), arXiv:2104.
- [8] Bradley Efron. *The jackknife, the bootstrap and other resampling plans*. SIAM, 1982.
- [9] G Peter Lepage. *corrfitter*. Version 8.1.1. 2020. URL: <https://github.com/gplepage/corrfitter>.
- [10] Particle Data Group et al. "Review of Particle Physics". In: *Progress of Theoretical and Experimental Physics* 2020.8 (Aug. 2020). 083C01. ISSN: 2050-3911. DOI: 10.1093/ptep/ptaa104. eprint: https://academic.oup.com/ptep/article-pdf/2020/8/083C01/34673740/rpp2020-vol2-2015-2092_18.pdf. URL: <https://doi.org/10.1093/ptep/ptaa104>.
- [11] Bipasha Chakraborty et al. "Nonperturbative comparison of clover and HISQ quarks in lattice QCD and the properties of the phi meson". In: *arXiv preprint arXiv:1703.05552* (2017).