

TRIAL-MATCHING–OPTIMIZED MEDICAL CHART – SARCOMA (FICTIONAL)

1. Patient Demographics

- Patient ID: SM-01459233
- Age: 52
- Sex: Female
- DOB: 1973-07-19
- Height: 165 cm
- Weight: 70.2 kg
- BMI: 25.8 kg/m²

2. Performance Status & Functional

- ECOG Performance Status: 1
- Karnofsky (approximate): 80–90%
- ADLs/IADLs: Independent

3. Primary Cancer Diagnosis

- Primary cancer type: Soft tissue sarcoma
- Histologic subtype: Undifferentiated pleomorphic sarcoma (UPS)
- Primary tumor site: Left proximal thigh (deep, intramuscular)
- Disease setting: Metastatic (lung metastases)
- Initial diagnosis date (sarcoma): 2023-11-10
- Clinical stage at initial diagnosis: cT3N0M0 (large, deep soft tissue sarcoma of extremity)
- Pathologic stage at initial resection: pT3N0M0, high grade (FNCLCC grade 3)
- Tumor size (primary, at resection): 9.2 cm greatest dimension
- Margins at primary resection: R0 (negative but close margins, <1 mm in some areas)
- Histologic grade: FNCLCC grade 3 (high grade)
- Lymphovascular invasion: Not definitively identified
- Nodal involvement: None clinically or pathologically documented
- Current disease status: Metastatic recurrence in lungs (multiple bilateral pulmonary nodules)

4. Disease Status / Imaging Summary

- Current disease status: Metastatic soft tissue sarcoma to lungs
- Latest restaging imaging: CT chest/abdomen/pelvis with contrast
- Date of latest imaging: 2025-09-15

Findings:

- Lungs: Multiple bilateral non-calcified pulmonary nodules, largest 1.8 cm in right lower lobe, additional nodules
- Pleura: No effusion
- Mediastinum: No bulky adenopathy
- Abdomen/pelvis: No liver, adrenal, or solid-organ metastases
- Bones: No definite osseous metastases
- Primary site (left thigh): Post-surgical changes only; no definite local recurrence on this scan

RECIST 1.1 Target Lesions (example):

- Target lesion 1: Right lower lobe lung nodule – 1.8 cm
- Target lesion 2: Left upper lobe lung nodule – 1.2 cm
- Target lesion 3: Right middle lobe lung nodule – 1.1 cm
- Sum of diameters (SoD): 4.1 cm

5. Prior Cancer-Directed Therapy (Sarcoma)

5.1 Surgery

- Wide local excision of left proximal thigh mass
- Date: 2024-01-05
- Margin status: R0, very close (<1 mm) in some areas

5.2 Radiation Therapy

- Adjuvant external beam radiotherapy to left thigh
- Dose: 60 Gy in 30 fractions
- Dates: 2024-02-10 to 2024-03-22
- Acute toxicities: Grade 2 skin erythema, resolved

5.3 Systemic Therapy – Prior Lines for Advanced Disease

Line 1 (metastatic setting)

- Regimen: Doxorubicin 75 mg/m² IV q21 days
- Cycles: 6 cycles
- Dates: 2024-05-01 to 2024-09-10
- Best response: Partial response (decrease in pulmonary nodules)
- Key toxicities: Grade 2 fatigue, grade 1 nausea; no clinically significant cardiotoxicity documented
- Reason for discontinuation: Completed planned 6 cycles, response then stable

Line 2 (metastatic setting)

- Regimen: Gemcitabine + docetaxel
- Dosing: Gemcitabine 900 mg/m² days 1 and 8, docetaxel 75 mg/m² day 8 q21 days
- Cycles: 4 cycles
- Dates: 2025-01-05 to 2025-03-25
- Best response: Stable disease, followed by progression in lung lesions
- Reason for discontinuation: Radiographic progression

5.4 Targeted/Other Agents

- Pazopanib, trabectedin, eribulin, or other sarcoma agents: None received
- Checkpoint inhibitor (PD-1/PD-L1): None
- Clinical trial participation for sarcoma: None to date

6. Key Biomarkers / Molecular Data

- Pathology: Undifferentiated pleomorphic sarcoma; no specific lineage-defining markers
- PD-L1 expression: CPS 1 (low)
- NTRK fusion: Not detected
- ALK/ROS1: Negative
- MSI status: Microsatellite stable (MSS)
- Tumor mutational burden (TMB): Intermediate (~7 mut/Mb)
- Genomic panel: No actionable fusions reported; multiple non-actionable mutations

7. Organ Function & Key Labs (Most Recent Pre-Trial Consideration)

- Date of labs: 2025-09-10

7.1 Hematologic

- WBC: 5.9 x10³/μL

- Hemoglobin: 11.8 g/dL
- Hematocrit: 35.1%
- Platelets: $210 \times 10^3/\mu\text{L}$
- ANC: $3.2 \times 10^3/\mu\text{L}$ (adequate for most protocols)
- Absolute lymphocyte count: $1.4 \times 10^3/\mu\text{L}$

7.2 Renal

- Serum creatinine: 0.86 mg/dL
- Estimated eGFR: $\sim 90 \text{ mL/min/1.73 m}^2$
- BUN: 14 mg/dL

7.3 Hepatic

- AST: 23 U/L
- ALT: 26 U/L
- Alkaline phosphatase: 101 U/L
- Total bilirubin: 0.6 mg/dL
- Albumin: 3.8 g/dL

7.4 Metabolic / Electrolytes

- Na: 137 mmol/L
- K: 4.1 mmol/L
- Ca: 9.0 mg/dL
- Glucose (fasting): 104 mg/dL

7.5 Coagulation

- PT/INR: 1.0
- aPTT: Within normal limits

7.6 Urinalysis / Proteinuria

- Dipstick: Negative for protein, negative for blood
- Quantitative protein/creatinine ratio: Not performed (would be required for some VEGF/TKI trials)

7.7 Infectious Disease Screening

- Hepatitis B surface antigen: Negative
- Hepatitis C antibody: Negative
- HIV antibody/antigen: Negative

8. Cardiac & Other Baseline Assessments

- Baseline echocardiogram prior to doxorubicin (2024-04-20): LVEF 60%
- Latest echocardiogram (2025-08-20): LVEF 57%, no heart failure symptoms
- Baseline EKG: Normal sinus rhythm, QTc 430 ms
- No history of symptomatic coronary artery disease, arrhythmias, or heart failure

9. Comorbidities (Trial-Relevant)

- Hypertension: Mild, controlled on amlodipine
- Hyperlipidemia: On statin therapy
- No known diabetes mellitus
- No chronic kidney disease
- No chronic liver disease
- No known autoimmune disease

- No history of thromboembolic disease (DVT/PE)
- No known chronic lung disease
- No other malignancies documented

10. Autoimmune / Infection / Immunosuppression History

- Autoimmune disease: None known
- Chronic systemic steroids: None
- Other immunosuppressive therapy: None
- Organ transplant: None
- Chronic infections: None
- TB history: None known; no symptoms

11. Concomitant Medications (Eligibility-Relevant)

- Amlodipine 5 mg PO daily
- Atorvastatin 20 mg PO nightly
- PRN ondansetron 8 mg PO for nausea (not used regularly)
- PRN acetaminophen for pain

Not on:

- Chronic anticoagulation
- Chronic systemic steroids
- Immunosuppressive biologics
- QT-prolonging antiarrhythmics

12. Reproductive / Consent-Related

- Biologic sex: Female
- Age: 52
- Pregnancy status: Not pregnant (serum pregnancy test negative when last obtained)
- Contraception: Uses barrier method; willing to comply with trial-mandated contraception
- Cognitive status: Intact, fully oriented, able to understand and sign informed consent

13. Social / Behavioral

- Tobacco use: Never smoker
- Alcohol: Occasional (1–2 drinks/month)
- Illicit drugs: Denies
- Occupation: Office administrator; currently on medical leave
- Support system: Lives with spouse and adult daughter; good social support
- Transportation / access: Reliable transportation to trial center

14. Prior/Planned Local Therapies for Metastases

- Prior lung metastasectomy: None
- Prior SBRT to pulmonary lesions: None
- Potential future consideration: SBRT or surgery for oligoprogressive disease, depending on trial protocol

15. Overall Trial Fit Summary

- Diagnosis: Metastatic undifferentiated pleomorphic sarcoma (UPS) of the left thigh with lung metastases
- Disease category: Advanced/metastatic soft tissue sarcoma
- Lines of prior systemic therapy for advanced disease: 2 lines
 - Line 1: Doxorubicin
 - Line 2: Gemcitabine + docetaxel

- Prior exposure to anthracycline: Yes (cumulative dose within typical limits; LVEF preserved)
- Prior exposure to checkpoint inhibitors: No
- Prior exposure to targeted agents (TKIs, ADCs): No
- Organ function: Adequate bone marrow, renal, and hepatic function for most phase I/II trials
- Cardiac function: LVEF preserved, no symptomatic cardiac disease
- Infectious/autoimmune: No major autoimmune disease; viral serologies negative
- Performance status: ECOG 1
- Major exclusion flags (from available data): None obvious