

Quaternions

Oskar Mampe

9th October 2019

Problem

An artist has given you two keyframe rotation matrices of an object. At time step 0, the artist wants the object rotated by 45° CCW around vector $(2, 1, 4)$ and at time step 100, the object should be rotated by 123° CW around vector $(0, 1, 1)$. Use quaternions to determine the axis and angle of rotation required to achieve this rotation in 100 steps. Show all work.

All calculations are rounded to 5 decimal places.

Solution

Taking the vector $\vec{v} = (2, 1, 4)$ and angle $\theta_0 = 22.5$.

$$q_0 = (\cos(\theta_0), \sin(\theta_0) \times |\vec{v}|)$$

$$||\vec{v}|| = \sqrt{2^2 + 1^2 + 4^2} = \sqrt{21} = 4.58258$$

$$|\vec{v}| = \frac{\vec{v}}{||\vec{v}||} = \frac{(2, 1, 4)}{4.58258} = (0.43644, 0.21822, 0.87287)$$

$$\cos(\theta_0) = 0.92388$$

$$\sin(\theta_0) = 0.38268$$

$$q_0 = (0.92388, 0.38268 \times 0.43644, 0.38268 \times 0.21822, 0.38268 \times 0.87287)$$

$$q_0 = 0.92388 + 0.16702i + 0.08351j + 0.33403k$$

Taking the vector $\vec{w} = (0, 1, 1)$ and angle $\theta_{100} = -61.5$.

$$q_{100} = (\cos(\theta_{100}), \sin(\theta_{100}) \times |\vec{w}|)$$

$$||w|| = \sqrt{0^2 + 1^2 + 1^2} = \sqrt{2} = 1.41421$$

$$|w| = \frac{|w|}{||w||} = \frac{(0, 1, 1)}{1.41421} = (0, 0.70711, 0.70711)$$

$$\cos(\theta_{100}) = 0.47716$$

$$\sin(\theta_{100}) = -0.87882$$

$$q_{100} = (0.47716, 0, -0.87882 \times 0.70711, -0.87882 \times 0.70711)$$

$$q_{100} = 0.47716 + 0i - 0.62142j - 0.62142k$$

Next, to find the quaternion required q_δ , the quaternion q_Δ needs to be calculated by multiplying quaternions q_{100} and the inverse of q_0 . Since q_0 is a unit quaternion, its magnitude is 1, so only taking the conjugate q_0^* is required.

$$q_0^* = 0.92388 - 0.16702i - 0.08351j - 0.33403k$$

$$q_{100} = 0.47716 + 0i - 0.62142j - 0.62142k$$

$$q_\Delta = q_0^* \times q_{100}$$

$$\begin{aligned} q_\Delta &= 0.44084 + 0i - 0.57412j - 0.57412k \\ &\quad - 0.07970i - 0i^2 + 0.10379ij + 0.10379ik \\ &\quad - 0.03985j - 0ji + 0.05189j^2 + 0.05189jk \\ &\quad - 0.15939k - 0ki + 0.20757kj + 0.20757k^2 \\ &= 0.44084 - 0.57412j - 0.57412k - 0.07970i \\ &\quad + 0.10379k - 0.10379j - 0.03985j - 0.05189 \\ &\quad + 0.05189i - 0.15939k - 0.20757i - 0.20757 \end{aligned}$$

$$q_\Delta = 0.18138 - 0.23538i - 0.71776j - 0.62972k$$

Now that q_Δ is found, θ_Δ and \vec{v} can be found. Then the quaternion q_δ can be formed by using $\frac{\theta_\Delta}{100}$ and \vec{v} .

$$(q_\delta)^{100} = q_\Delta$$

$$\theta_\delta = \frac{\theta_\Delta}{100}$$

$$\theta_\Delta = \arccos(0.18138) = 79.54985$$

$$\theta_\delta = \frac{79.54985}{100}$$

$$\theta_\delta = 0.79550$$

$$\sin(79.54985) = 0.98341$$

$$\vec{v}_\delta = \left(-\frac{0.23538}{0.98341}, -\frac{0.71776}{0.98341}, -\frac{0.62972}{0.98341}\right)$$

$$\vec{v}_\delta = (-0.23935, -0.72987, -0.64034)$$

Finally, the quaternion q_δ with the required angle and axis can be formed:

$$\cos(0.79550) = 0.99990$$

$$\sin(0.79550) = 0.01388$$

$$q_\delta = (0.99990, 0.01388 \times -0.23935, 0.01388 \times -0.72987, 0.01388 \times -0.64034)$$

$$q_\delta = 0.99990 - 0.00332i - 0.01013j - 0.00889k$$