
Top 5 KEGG pathways for each condition
Type II diabetes mellitus
Th1 and Th2 cell differentiation
Taurine and hypotaurine metabolism
Synaptic vesicle cycle
Salivary secretion
Ribosome
Proximal tubule bicarbonate reclamation
Prion diseases
Phototransduction
Parkinson disease
Oxytocin signaling pathway
Oxidative phosphorylation
Nicotine addiction
Morphine addiction
Mineral absorption
Long−term potentiation
Long−term depression
Insulin secretion
Huntington disease
HIF−1 signaling pathway
GnRH signaling pathway
Glutamatergic synapse
Glioma
Gastric acid secretion
GABAergic synapse
Ferroptosis
Estrogen signaling pathway
Endocrine and other factor−regulated calcium reabsorption
Cortisol synthesis and secretion
Collecting duct acid secretion
Circadian rhythm
Circadian entrainment
Cell adhesion molecules (CAMs)
Cardiac muscle contraction
cAMP signaling pathway
Butanoate metabolism
Antigen processing and presentation
Amyotrophic lateral sclerosis (ALS)
Amphetamine addiction
Alzheimer disease
Aldosterone−regulated sodium reabsorption
Aldosterone synthesis and secretion
Adrenergic signaling in cardiomyocytes
Adherens junction
Term
Astro
L2_3_IT
L4
L5
L6
Lamp5
Non−neuronal
Oligo
Pvalb
Sncg
Sst
Vip
Cell Type
Odds Ratio
25
50
75
Adjusted P−Value
0.04
0.03
0.02
0.01