Generalized Hierarchical Bayesian Segmentation with Irregular Designs, Multi-Sequence Hierarchies, and Grouped/Latent-Group Designs

Omid Shams Solari[1]

[1]sAIm Labs

ABSTRACT

We develop a generalized hierarchical Bayesian method for multiple-changepoint segmentation of ordered data observed on regular or irregular designs. For a regular exponential-family observation model with a proper segmentwise conjugate prior and finite normalizing constants, cumulative sufficient statistics determine the marginal likelihood and posterior moments of every candidate segment. These segment marginal likelihoods enter dynamic-programming recursions for the marginal likelihood conditional on the number of segments, the posterior distribution of the segment count, marginal changepoint probabilities, and posterior moments of the piecewise-constant signal. A separate max-sum recursion with backtracking returns the joint maximum-a-posteriori partition. Thus, the partition calculation is common across observation families once the segment marginal likelihoods are specified, while the statistical meaning of each segment remains family-specific.

The hierarchical formulation includes design-dependent partition priors for irregular coordinates, common-changepoint models for multiple conditionally independent sequences, known-group models, and a finite latent-group template model. The latent-group algorithm uses a Jensen minorization and exact responsibility-weighted max-sum updates for the stated finite objective; because its component scores are not normalized sampling densities, it is not interpreted as a finite-mixture likelihood. When a nonconjugate segment integral is replaced by a deterministic approximation, a uniform error bound on reachable segment log marginal likelihoods implies explicit bounds on partition marginal likelihoods and posterior odds.

Simulation studies report boundary-posterior calibration for Gaussian data (expected calibration error approximately 0.010, Brier score approximately 0.011), applications of common partition recursions to Gaussian, Poisson, Binomial, and Beta-valued responses, and 96% hard allocation accuracy in one two-group template experiment. A separate 400-dataset stress study evaluates template recovery and objective monotonicity. Observed Gaussian computation times range from 0.0455 seconds at $n = 50$ to 1.0585 seconds at $n = 800$ for $k_{\max} = 20$. Analyses of well-log, array-CGH, financial-return, and methylation data illustrate model use, while the lack of independently scored changepoints limits conclusions about external segmentation accuracy. Additional simulation and comparison protocols are specified as planned work and do not contribute numerical findings to this report.

Keywords Bayesian changepoint analysis, multiple changepoints, product partition models, exponential families, conjugate priors, hierarchical Bayesian models, dynamic programming, irregular designs, multiple sequences, latent groups

Correspondence omid.solari@saimai.com

CONTENTS

1 Introduction 2
2 Statistical foundations and related work 3
3 Problem formulation and notation 5
4 Generalized segment models and integrated likelihoods 7
5 Posterior computation over ordered partitions 14
6 Irregular designs and hierarchical extensions 19
7 Approximate segment integration, posterior prediction, and evaluation 25
8 Algorithms, computational complexity, and numerical implementation 33
9 Experimental protocols 36
10 Results 43
11 Limitations and scope 53
12 Conclusion 55
A Stable numerical implementation of posterior summaries 58
B Log-space dynamic programs 59
C Supplementary agreement diagnostics 60
D Reproducibility 61

1 Introduction

Multiple-changepoint analysis represents an ordered series by contiguous segments within which a sampling-model parameter is constant. It is used in statistical signal processing, genomics, environmental science, and econometrics. The statistical task is to estimate the segment count, changepoint locations, segment parameters, and their uncertainty. Bayesian methods assign posterior probabilities to segmentations and can average over these alternatives when estimating a piecewise-constant signal or predicting future observations (Barry & Hartigan, 1992, 1993; Fearnhead, 2006; Hutter, 2007).

For an offline changepoint model with independent segment parameters and a factorizing partition prior, the probability of a partition is a product of integrated segment likelihoods and local prior terms. Dynamic programming then sums these products without enumerating every partition (Barry & Hartigan, 1992; Fearnhead, 2006; Hutter, 2007). This formulation is well established. Its practical use across heterogeneous outcomes depends on calculating segment marginal likelihoods with the correct sampling-model normalization and integrating the partition prior with the same care.

BayesBreak develops a generalized hierarchical formulation of this model. For regular exponential-family observations with proper conjugate priors, segment marginal likelihoods and

posterior moments follow from cumulative sufficient statistics. The resulting partition recursions do not depend on which member of the family supplied the segment scores. The framework includes nonuniform partitions on irregular physical coordinates, several sequences sharing changepoints, known groups, and a finite latent-group template criterion. Nonconjugate segment integrals can also be approximated locally, but exact posterior interpretation of the resulting dynamic program requires control of the segment-level approximation error.

Contributions. The contributions of this paper are as follows.

1. We specify a common conjugate segment calculation for exponential-family sampling models with known exposure, trial-count, and precision descriptors, explicitly distinguishing ordinary marginal likelihoods from powered likelihood integrals. The calculation provides segment marginal likelihoods and the posterior moments required for Bayesian piecewise-constant regression.
2. We derive normalized sum-product recursions for the segment-count posterior, ordered and unordered changepoint marginals, and posterior signal moments. We also derive a separate max-sum recursion for joint MAP segmentation, with immediate proofs of the identities and decision rules.
3. We extend the local-factor model to physical segment lengths and candidate-boundary weights, and to the exact pooling of independently sampled sequences under a common segmentation.
4. We study known-group pooling and a finite latent-template criterion for unknown groups. A Jensen minorization leads to alternating allocation and responsibility-weighted optimal-partition updates, with monotonicity for the specified objective rather than a claim of full Bayesian mixture inference.
5. We analyze the effect of segment marginal-likelihood approximation error on partition probabilities and posterior odds. The analysis is conditional on a uniform bound across admissible segments, not on a numerical method's name or an unverified finite-sample error estimate.

The experiments examine simulation recovery, posterior calibration, numerical approximations, computation time, and applications to well-log measurements, array-CGH, financial returns, and methylation. Existing results are reported separately from the additional experiments specified in Section 9. The latter are planned evaluations and are not used to support conclusions about accuracy or comparative performance.

Section 2 reviews the relevant statistical literature. Section 3 defines the sampling model, partition, and inferential targets. Sections 4 and 5 derive segment integration and exact partition inference. Section 6 covers hierarchical extensions. Sections 7 and 8 address numerical approximation, prediction, and implementation. The experimental protocols and results follow in Sections 9 and 10.

2 Statistical foundations and related work

2.1 Exponential families and conjugate segment models

The exponential-family representation separates a finite-dimensional sufficient statistic from the log normalizer of the sampling distribution. Diaconis and Ylvisaker characterize conjugate priors for natural exponential families and the hyperparameter conditions under which those priors are proper (Diaconis & Ylvisaker, 1979). In a changepoint model, the same calculation is applied separately

to every candidate segment. BayesBreak uses this standard conjugate theory; its contribution is to formulate the resulting segment marginal likelihoods and posterior moment numerators so that they can be reused by the same partition recursions and by the hierarchical extensions developed below. Exactness requires the stated regularity, propriety, and finiteness conditions. The method does not assert closed-form integration for arbitrary nonconjugate parameterizations.

2.2 Product partition models and exact Bayesian recursions

Product partition models assign a partition probability proportional to a product of segment cohesions and combine this prior with segmentwise integrated likelihoods (Barry & Hartigan, 1992, 1993). This is the principal probabilistic lineage of the present work. Fearnhead developed exact recursions and posterior simulation for a broad class of Bayesian multiple-changepoint models with conditionally independent segments (Fearnhead, 2006); Hutter derived exact Bayesian regression for piecewise-constant functions, including posterior break probabilities and model-averaged regression curves (Hutter, 2007). Dependence between adjacent segments can sometimes be accommodated by enlarging the recursion state (Fearnhead & Liu, 2011). Bayesian Blocks provides a related local-fitness and dynamic-programming construction for astronomical time series (Scargle et al., 2013).

BayesBreak does not claim priority for product partition models, conjugate segment integration, exact changepoint recursions, posterior changepoint probabilities, or Bayesian piecewise-constant regression. It develops a generalized hierarchical model in which these established calculations are stated with common notation across exponential-family segment models, irregular-design priors, multiple sequences, observed groups, and a finite latent-group extension. It also keeps posterior marginalization and joint MAP optimization as separate recursions rather than treating one as an approximation to the other.

2.3 Frequentist segmentation and optimization

Dynamic programming also underlies frequentist segmentation with additive segment costs (Auger & Lawrence, 1989; Jackson et al., 2005). Penalized formulations and pruning lead to PELT and related pruned exact algorithms (Killick et al., 2012; Rigaill, 2015). Wild binary segmentation and SMUCE pursue different detection and uncertainty goals (Fryzlewicz, 2014; Frick et al., 2014). Total variation and the fused lasso replace discrete partition search with convex regularization (Rudin et al., 1992; Tibshirani et al., 2005); the group fused lasso encourages aligned changes across several sequences (Bleakley & Vert, 2011). Circular binary segmentation is a standard comparator for array-CGH data (Olshen et al., 2004). These methods are important computational and empirical references, but their estimands are penalized or testing-based rather than the posterior distribution over partitions considered here.

2.4 Irregular designs and design-dependent partition priors

Generalized product partition models and product partition models with covariates show how covariate information can alter partition probabilities through cohesion functions (Park & Dunson, 2010; Müller et al., 2011). BayesBreak specializes this idea to ordered observations. It distinguishes a segment-span cohesion from a candidate-boundary hazard because they encode different prior assumptions. Both factors can be included in the ordered-partition probability when they remain local to a segment or transition. Neither is treated as a likelihood weight.

2.5 Hierarchical and multi-sequence changepoint models

Hierarchical Bayesian changepoint analysis has a long history (Carlin et al., 1992). BASIC models the tendency of changepoints to co-occur across multiple sequences through a hierarchical empirical-Bayes prior and provides posterior sampling and optimization algorithms (Fan & Mackey, 2017). Joint random partition models provide a Bayesian nonparametric construction for dependent partitions across several ordered processes (Quinlan et al., 2024). BayesBreak studies a more restrictive setting for exact finite computation: aligned sequences are conditionally independent given a common partition, so their segment marginal likelihoods multiply. Known groups apply this calculation separately by group. The finite-candidate consistency theorem in Section 6 concerns this common-partition model and does not imply consistency for arbitrary hierarchical dependence.

2.6 Latent groups and common structural changes

Model-based clustering of time-dependent observations according to common structural changes is an active neighboring literature (Corradin et al., 2026). The BayesBreak latent-group construction is narrower than a full probability mixture over all partitions. It optimizes a finite objective of the form $\sum_s \log \sum_g \pi_g S_s(\tau_g)$, where $S_s(\tau_g) > 0$ is the specified subject–template score. A Jensen minorization yields auxiliary weights and responsibility-weighted max-sum updates. Because the S_s need not be normalized densities over the data space, the algorithm is described as minorization–maximization for a finite latent-group objective, not as maximum likelihood for an identifiable finite mixture (Dempster et al., 1977; Teicher, 1963).

2.7 Nonconjugate segment calculations

When a segment integral is unavailable in closed form, deterministic approximations such as Laplace approximation, the Jaakkola–Jordan variational bound, expectation propagation, Pólya–Gamma augmentation, or numerical quadrature may be used (Jaakkola & Jordan, 2000; Minka, 2001; Polson et al., 2013). The ordered-partition recursion is unchanged after an approximate segment marginal-likelihood table is supplied, but its output is then exact only for that table. Section 7 therefore proves a conditional error-propagation result: a uniform bound on reachable segment log marginal likelihoods yields bounds on partition marginal likelihoods and posterior odds. The theorem does not supply the segmentwise bound for a particular approximation routine.

2.8 Position of the present work

The present contribution is a generalized hierarchical Bayesian segmentation model whose observation-family dependence is confined to segment marginal likelihoods and posterior moments. The paper derives the common posterior recursions, the separate MAP recursion, the irregular-design prior factors, exact shared-boundary pooling, the finite latent-group optimization, and the conditional approximation bounds in one notation. Its empirical claims concern the executed simulations and case studies reported here; it does not claim a universal complexity improvement or superiority over every established changepoint method.

3 Problem formulation and notation

3.1 Observations, coordinates, and admissible partitions

Let $\{(x_i, y_i, d_i)\}_{i=1}^n$ be observations with ordered design points $x_1 < \cdots < x_n$. The response y_i may be continuous or discrete; d_i contains known observation-model quantities, such as a Poisson exposure, a Binomial trial count, or a Gaussian precision. A separate indicator $o_i \in \{0, 1\}$ specifies whether the response is observed. Missing outcomes do not contribute likelihood factors. We write y_{obs} for the observed responses. The default analysis uses ordinary sampling likelihoods. When likelihood powers $a_i \geq 0$ are deliberately introduced, they define a *generalized Bayesian* calculation and are not interpreted as exposure, trial count, or physical spacing.

Candidate partition edges are $u_0 < u_1 < \cdots < u_n$, with interior edge u_i separating observations i and $i + 1$. For irregular observations, an admissible choice is $u_i = (x_i + x_{i+1})/2$ for $1 \leq i < n$, together with prespecified endpoints $u_0 < x_1$ and $u_n > x_n$. Where the observation domain has known physical limits, use those limits instead. Choosing these edges is part of the model specification. On an index grid, use $u_i = i$. These physical edges are distinct from the observed positions x_i and from the family descriptor d_i.

A partition into k nonempty contiguous segments has ordered boundary vector

$$t = (t_0, \ldots, t_k), \qquad 0 = t_0 < t_1 < \cdots < t_k = n, \qquad I_q(t) = \{t_{q-1} + 1, \ldots, t_q\}. \tag{1}$$

The candidate block $(i, j]$ denotes observations $i + 1, \ldots, j$ and has physical length $u_j - u_i$. Let $\mathcal{T}_k$ be the set of partitions satisfying the declared candidate-boundary and minimum-length restrictions. These restrictions are set independently of the responses. Write $1 \leq k \leq k_{\max}$ for the admissible segment counts.

3.2 Hierarchical sampling model and prior

Within segment q, observations are conditionally independent given the segment parameter θ_q:

$$p(y_{\text{obs}} \mid t, k, \theta_{1:k}, d) = \prod_{q=1}^{k} \prod_{i \in I_q(t): o_i = 1} p(y_i \mid \theta_q, d_i), \qquad \theta_q \overset{\text{ind}}{\sim} \pi_0. \tag{2}$$

The segment prior π_0 is proper and does not depend on the partition unless explicitly specified. The prior on the count and the conditional prior on the partition are $p(k)$ and $p(t \mid k, u)$, respectively. For the block-local factorization used below, write

$$p(t \mid k, u) = \frac{1}{C_k(u)} \prod_{q=1}^{k} \gamma_u(t_{q-1}, t_q), \qquad C_k(u) = \sum_{t \in \mathcal{T}_k} \prod_{q=1}^{k} \gamma_u(t_{q-1}, t_q). \tag{3}$$

The definition assumes $0 < C_k(u) < \infty$. Segment cohesion and candidate-boundary hazards are two components of γ_u and are defined in Section 4. For the uniform conditional prior, $\gamma_u \equiv 1$ on $\mathcal{T}_k$ and $C_k = |\mathcal{T}_k|$; in the unrestricted case, $C_k = \binom{n-1}{k-1}$.

3.3 Inference and decision targets

Define the integrated likelihood of a candidate block by

$$A_{ij}^{(0)} = \int \pi_0(\theta) \prod_{r=i+1}^{j} p(y_r \mid \theta, d_r)^{o_r} \, \mathrm{d}\theta. \tag{4}$$

Here a missing response contributes the multiplicative identity one; in particular, a block containing no observed responses has $A_{ij}^{(0)} = 1$ under a proper prior. The count-conditional marginal likelihood and posterior are

$$p(y_{\mathrm{obs}} \mid k, u) = \frac{1}{C_k(u)} \sum_{t \in \mathcal{T}_k} \prod_{q=1}^{k} \gamma_u(t_{q-1}, t_q) A_{t_{q-1}, t_q}^{(0)}, \qquad p(k \mid y_{\mathrm{obs}}, u) \propto p(k)\, p(y_{\mathrm{obs}} \mid k, u). \tag{5}$$

The inferential targets are the full count posterior, ordered-boundary marginals $p(t_q = h \mid y, k)$, boundary-event probabilities $p(b_h = 1 \mid y, k)$, posterior moments of a declared segment signal $m_\star(\theta)$, and the posterior predictive distribution. The *marginal-MAP count* maximizes $p(k \mid y)$; a *conditional joint MAP partition* maximizes $p(t \mid y, \widehat{k})$; the *global joint MAP pair* maximizes $p(k, t \mid y)$. These are different posterior summaries and can select different partitions. Exact formulas and algorithms are given in Section 5.

3.4 Extension to related sequences

For S aligned sequences, let $y_i^{(s)}$ and $\theta_q^{(s)}$ denote the subject response and its segment parameter. In the shared-boundary model, t is common to all S sequences while the $\theta_q^{(s)}$ are independent given t. Known groups have distinct group-specific partitions. For unknown group labels, Section 6 defines a separate finite-template score criterion and its minorization–maximization procedure. The latter is not a posterior distribution over a normalized finite mixture unless an additional probabilistic model is supplied.

4 Generalized segment models and integrated likelihoods

4.1 Segment likelihood and support

The observations, candidate boundary edges, admissible partitions, and primary inferential targets are defined in Section 3. For an admissible candidate block $(i, j]$, the segment integral is evaluated under its stated sampling model and proper prior. A structurally disallowed block has zero prior weight; a permitted block with missing observations is integrated over its observed outcomes only. These cases must not be conflated.

> *Assumption* 4.1 (Offline structural support and block independence). The candidate segment-count set $\{1, \ldots, k_{\max}\}$ is finite. Structural admissibility of a segment is fixed by the candidate grid, declared length restrictions, and prior support before the response values are observed. A structurally admissible segment has finite nonnegative local evidence; zero evidence removes its likelihood contribution but does not alter the partition-prior normalizer. Structurally excluded segments have local prior factor zero and log-score $-\infty$. Conditional on a partition, segment parameters are independent across segments. The conditional partition prior factorizes through

Table 1. Notation used throughout the manuscript.

Symbol	Definition	Symbol	Definition
$x_i,\ u_i$	Observed position; candidate partition edge	y_i, d_i, o_i	Response, family descriptor, observation mask
n, S, G	Observations, sequences, latent groups	$k, k_{\max}$	Segment count and imposed upper bound
$t, \mathcal{T}_k$	Partition and its admissible set	$I_q(t)$	Index set of segment q
θ_q, π_0	Segment parameter and proper prior	$m_\star(\theta)$	Reported segment-level signal functional
$A_{ij}^{(0)}$	Integrated block likelihood	$\gamma_u(i,j)$	Local partition-prior factor
$C_k(u)$	Conditional partition-prior normalizer	$L_{k,j}, R_{k,i}$	Forward and backward partition sums
$M_{k,j}$	Max-sum optimal prefix score	$p(b_h = 1 \mid y, k)$	Conditional boundary-event probability

local terms as in Assumption 4.4. For approximate segment routines, the same statements define an exact posterior for the supplied approximate score array; comparison to the intended nonconjugate model additionally requires Assumption 7.1.

4.2 Exponential-family segment likelihoods with exposure and power weights

Let d_r collect scientifically specified family information, such as Poisson exposure, Binomial trial count, or Gaussian precision. Let $a_r \geq 0$ be an optional power or reliability weight that multiplies the complete log contribution. The canonical local likelihood is

$$\begin{aligned} \log p(y_r \mid \theta, d_r)^{a_r} = a_r \log h(y_r, d_r) \\ + a_r S(y_r, d_r)^\top \eta(\theta) \\ - a_r E(d_r) A(\theta). \end{aligned} \tag{6}$$

The corresponding block summaries are

$$\begin{aligned} S_{ij} &= \sum_{r=i+1}^{j} a_r S(y_r, d_r), \\ W_{ij} &= \sum_{r=i+1}^{j} a_r E(d_r), \\ H_{ij} &= \sum_{r=i+1}^{j} a_r \log h(y_r, d_r). \end{aligned} \tag{7}$$

This notation prevents three distinct quantities from being collapsed into a single scalar "weight": design spacing affects the partition prior, d_r changes the observation law, and a_r raises the entire observation contribution to a power.

For conjugate hyperparameters (α, β), use the Diaconis–Ylvisaker normalizer convention

$$p(\theta \mid \alpha, \beta) = \exp\{\eta(\theta)^\top \alpha - \beta A(\theta) - \log Z(\alpha, \beta)\},$$
$$Z(\alpha, \beta) = \int e^{\eta(\theta)^\top \alpha - \beta A(\theta)} \, \mathrm{d}\theta. \tag{8}$$

Fix a scientifically declared reference descriptor $d_\star$ when the conditional observation mean depends on exposure or trial structure, and write $m_\star(\theta) = m(\theta; d_\star)$ for the segment functional reported by the analysis (for example, a Poisson rate at unit exposure rather than an exposure-specific count mean). Define

$$A_{ij}^{(0)} = \int p(\theta \mid \alpha_0, \beta_0) \prod_{r=i+1}^{j} p(y_r \mid \theta, d_r)^{a_r} \, \mathrm{d}\theta,$$
$$A_{ij}^{(q)} = A_{ij}^{(0)} \, \mathbb{E}[m_\star(\theta)^q \mid y_{i+1:j}, d, a], \qquad q = 1, 2.$$

For unit powers $a_r = 1$, $A_{ij}^{(0)}$ is the ordinary block marginal likelihood. For non-unit powers it is a powered/generalized-Bayes integral rather than, in general, a probability-normalized likelihood of the observations. Posterior moments in the second display are taken with respect to the corresponding normalized posterior kernel, which may be a generalized posterior.

Theorem 4.2 (Exposure-aware conjugate block integration)**.** *If $Z(\alpha_0 + S_{ij}, \beta_0 + W_{ij}) < \infty$, then*

$$A_{ij}^{(0)} = e^{H_{ij}} \frac{Z(\alpha_0 + S_{ij}, \beta_0 + W_{ij})}{Z(\alpha_0, \beta_0)}. \tag{9}$$

The block posterior is conjugate with hyperparameters $(\alpha_0 + S_{ij}, \beta_0 + W_{ij})$. Whenever its rth moment exists,

$$A_{ij}^{(r)} = A_{ij}^{(0)} \mathbb{E}[m_\star(\theta)^r \mid \alpha_0 + S_{ij}, \beta_0 + W_{ij}]. \tag{10}$$

Proof. **Step 1 (aggregate the powered sampling contributions).** By conditional independence and Equation (6), multiplication of the local factors yields

$$\prod_{r=i+1}^{j} p(y_r \mid \theta, d_r)^{a_r} = \exp\{H_{ij} + \eta(\theta)^\top S_{ij} - W_{ij} A(\theta)\}.$$

The three sums in Equation (7) collect the base-measure term, the sufficient-statistic coefficient, and the coefficient of the log-partition function, respectively. **Step 2 (combine likelihood and prior).** Multiplication by Equation (8) yields

$$\frac{e^{H_{ij}}}{Z(\alpha_0, \beta_0)} \exp\{\eta(\theta)^\top (\alpha_0 + S_{ij}) - (\beta_0 + W_{ij}) A(\theta)\}.$$

This is a constant times the conjugate kernel with the displayed updated hyperparameters. **Step 3 (integrate).** The assumption of a finite updated normalizer permits integrating the kernel. Its integral is $Z(\alpha_0 + S_{ij}, \beta_0 + W_{ij})$, proving Equation (9). Division of the joint density by this integral gives the posterior conjugate distribution. **Step 4 (moment insertion).** For any integrable $m_\star(\theta)^r$, insert that factor before integration. Dividing and multiplying the resulting integral by $A_{ij}^{(0)}$ identifies the updated posterior expectation, proving Equation (10). For $a_r \neq 1$

the algebra is exact for a powered likelihood, but the result is not in general a normalized ordinary sampling-model marginal likelihood. □

Remark 4.3 (Scope of exactness). Theorem 4.2 is exact for the displayed powered, exposure-aware model. It does not permit a Binomial trial count to be treated as a likelihood power, an irregular coordinate gap to be inserted as exposure, or a numerical integral to be described as conjugate closed form.

4.3 Local prior factors

Assumption 4.4 (Factorized ordered-partition prior). For fixed k and boundary coordinates u, the prior on t is

$$p(t \mid k, u) = C_k(u)^{-1} \prod_{q=1}^{k} c_u(t_{q-1}, t_q) \prod_{q=1}^{k-1} h_u(t_q), \tag{11}$$

where $c_u(i,j) \geq 0$ is a complete-segment cohesion and $h_u(j) \geq 0$ is an interior-boundary hazard. Define $\gamma_u(i,j) = c_u(i,j)h_u(j)^{\mathbb{1}\{j<n\}}$ and $\widetilde{A}_{ij}^{(r)} = A_{ij}^{(r)}\gamma_u(i,j)$.

The normalizer $C_k(u)$ is itself a partition sum and can be computed by the same dynamic program after replacing all likelihood evidences by one. This ensures that fixed-k evidence remains properly normalized for nonuniform local priors.

Table 2. Supported local block constructions. Closed-form rows instantiate Theorem 4.2; the Beta-valued response uses deterministic one-dimensional quadrature and is therefore exact only up to its numerical tolerance.

Family	Descriptor d_r	Segment parameter	Integrated block construction	Evaluation
Gaussian, known variance	precision or inverse variance	mean μ	Gaussian–Gaussian normalizer and posterior moments	closed form
Poisson	exposure e_r	rate λ	Gamma–Poisson normalizer; rate $e_r\lambda$	closed form
Binomial	trial count m_r	probability p	Beta–Binomial normalizer; m_r trials	closed form
Negative Binomial	fixed dispersion	success parameter	Beta-conjugate distributional parameterization	closed form
Beta-valued response	known precision ϕ_r	mean $\mu \in (0,1)$	one-dimensional integral over a transformed mean	quadrature
Nonconjugate GLM	link-specific inputs	regression parameter	Laplace, variational, EP, PG, or quadrature block score	quadrature or local approximation

4.4 Segment marginal-likelihood interface

The partition recursions require only a triangular array of segment log marginal likelihoods and, when posterior signal moments are requested, corresponding segment moment summaries. A family-

specific implementation returns the segment log marginal likelihood, posterior hyperparameters or moments, a validity flag, and any numerical diagnostic needed to interpret an approximate value. Prefix summaries make Gaussian, Poisson, and Binomial segment calculations constant time after $\mathcal{O}(n)$ preprocessing. The recursion over partitions is therefore unchanged when the observation family changes.

Implementation interface. For each admissible block, the local calculation supplies $\log A_{ij}^{(0)}$, the conditional posterior moments required for the scientific target, and convergence information if the integral is evaluated numerically. A structurally excluded block has prior-adjusted log score $-\infty$. The partition dynamic program uses the resulting score table without reference to the observation family's internal parameterization.

4.5 Closed-form segment distributions and posterior moments

The following calculations make the model-specific part of Theorem 4.2 explicit. Throughout this subsection, observations have their ordinary sampling interpretation ($a_r = 1$). An exposure, precision, or trial count is part of the sampling distribution, not an exponent applied to that distribution. The target $m_\star(\theta)$ is specified in each model rather than tacitly identified with an exposure-dependent observation mean.

4.5.1 Gaussian observations with known precisions

Proposition 4.5 (Gaussian evidence and first two signal moments)**.** *Suppose* $y_r \mid \mu, w_r \sim \mathcal{N}(\mu, \sigma^2/w_r)$ *independently, with known* $w_r > 0$*, and* $\mu \sim \mathcal{N}(\nu, \rho^2)$*, where* $\sigma^2, \rho^2 > 0$*. On* $B = (i, j]$ *let*

$$N_B = j - i, \qquad W_B = \sum_{r \in B} w_r, \qquad \bar{y}_B = W_B^{-1} \sum_{r \in B} w_r y_r,$$
$$Q_B = \sum_{r \in B} w_r (y_r - \bar{y}_B)^2, \qquad V_B = (\rho^{-2} + W_B/\sigma^2)^{-1}, \qquad \nu_B = V_B(\nu/\rho^2 + W_B \bar{y}_B/\sigma^2).$$

Then $\mu \mid y_B \sim \mathcal{N}(\nu_B, V_B)$ *and*

$$\log A_B^{(0)} = -\frac{N_B}{2}\log(2\pi\sigma^2) + \frac{1}{2}\sum_{r \in B} \log w_r - \frac{Q_B}{2\sigma^2} + \frac{1}{2}\log\frac{\sigma^2}{\sigma^2 + \rho^2 W_B} - \frac{W_B(\bar{y}_B - \nu)^2}{2(\sigma^2 + \rho^2 W_B)}. \tag{12}$$

With $m_\star(\mu) = \mu$*, the posterior signal moments are* ν_B *and* $V_B + \nu_B^2$*.*

Proof. **Step 1 (separate residual and mean information).** Completing the weighted square gives the exact identity

$$\sum_{r \in B} w_r (y_r - \mu)^2 = Q_B + W_B(\mu - \bar{y}_B)^2,$$

so the block likelihood is

$$(2\pi\sigma^2)^{-N_B/2}\Big(\prod_{r \in B} w_r^{1/2}\Big) \exp\left\{-\frac{Q_B}{2\sigma^2} - \frac{W_B(\mu - \bar{y}_B)^2}{2\sigma^2}\right\}.$$

Step 2 (derive the posterior). Multiplying by the normal prior and collecting coefficients of μ^2 and μ yields precision $\rho^{-2} + W_B/\sigma^2$ and precision-weighted mean $\nu/\rho^2 + W_B\bar{y}_B/\sigma^2$. Hence the posterior is $\mathcal{N}(\nu_B, V_B)$. **Step 3 (integrate the segment mean).** The remaining normal integral is

$$\int \exp\left\{-\frac{W_B(\mu-\bar{y}_B)^2}{2\sigma^2}\right\}\mathcal{N}(\mu \mid \nu,\rho^2)\,\mathrm{d}\mu = \left(\frac{\sigma^2}{\sigma^2+\rho^2 W_B}\right)^{1/2}\exp\left\{-\frac{W_B(\bar{y}_B-\nu)^2}{2(\sigma^2+\rho^2 W_B)}\right\}.$$

Multiplying the constants from Step 1 and taking logarithms gives Equation (12). The first two normal moments follow from the posterior mean and variance. In particular, ν_B may be negative: its contribution must never be stored as the logarithm of a signed moment numerator. □

4.5.2 Poisson counts with genuine exposure

Proposition 4.6 (Poisson–Gamma integration). *If $y_r \mid \lambda, e_r \sim \mathrm{Pois}(e_r\lambda)$ with known $e_r > 0$ and $\lambda \sim \mathrm{Gamma}(a_0, b_0)$ in shape–rate form, define $Y_B = \sum_{r\in B} y_r$, $E_B = \sum_{r\in B} e_r$ and $H_B = \sum_{r\in B}(y_r \log e_r - \log\Gamma(y_r+1))$. Then*

$$A_B^{(0)} = e^{H_B}\frac{b_0^{a_0}\Gamma(a_0+Y_B)}{\Gamma(a_0)(b_0+E_B)^{a_0+Y_B}}, \tag{13}$$

and $\lambda \mid y_B \sim \mathrm{Gamma}(a_0+Y_B, b_0+E_B)$. For unit-exposure rate $m_\star(\lambda)=\lambda$, the first two posterior moments are a_B/b_B and $a_B(a_B+1)/b_B^2$, with $(a_B,b_B)=(a_0+Y_B,b_0+E_B)$.

Proof. **Step 1 (collect the ordinary Poisson likelihood).** The product of probability mass functions is

$$\prod_{r\in B}\frac{e^{-e_r\lambda}(e_r\lambda)^{y_r}}{y_r!} = e^{H_B}\lambda^{Y_B}e^{-E_B\lambda}.$$

The coefficient of $\log\lambda$ is the observed count Y_B, whereas the coefficient of $-\lambda$ is the exposure E_B; replacing both by a common likelihood-power weight would be a different model. **Step 2 (combine with the Gamma prior).** The prior density is $b_0^{a_0}\lambda^{a_0-1}e^{-b_0\lambda}/\Gamma(a_0)$. Multiplication gives a Gamma kernel with shape a_0+Y_B and rate b_0+E_B. **Step 3 (integrate and extract moments).** Apply $\int_0^\infty \lambda^{a_B-1}e^{-b_B\lambda}\,\mathrm{d}\lambda = \Gamma(a_B)/b_B^{a_B}$ to obtain Equation (13). Gamma moment identities give the stated first and second moments. □

4.5.3 Binomial counts with known trial numbers

Proposition 4.7 (Beta–Binomial integration). *If $y_r \mid p, m_r \sim \mathrm{Binom}(m_r, p)$ with $m_r \in \mathbb{N}_0$, $0 \le y_r \le m_r$, and $p \sim \mathrm{Beta}(a_0, b_0)$, set $Y_B = \sum y_r$, $M_B = \sum m_r$, and $H_B = \sum \log\binom{m_r}{y_r}$. Then*

$$A_B^{(0)} = e^{H_B}\frac{B(a_0+Y_B, b_0+M_B-Y_B)}{B(a_0,b_0)}, \tag{14}$$

with posterior $\mathrm{Beta}(a_B,b_B)$, $a_B = a_0+Y_B$, $b_B = b_0+M_B-Y_B$. For $m_\star(p)=p$, the posterior first two moments are $a_B/(a_B+b_B)$ and $a_B(a_B+1)/[(a_B+b_B)(a_B+b_B+1)]$.

Proof. **Step 1 (likelihood factorization).** Independence and the Binomial mass function give

$$\prod_{r\in B}\binom{m_r}{y_r}p^{y_r}(1-p)^{m_r-y_r}=e^{H_B}p^{Y_B}(1-p)^{M_B-Y_B}.$$

Step 2 (integrate with the Beta prior). Multiplication by $p^{a_0-1}(1-p)^{b_0-1}/B(a_0,b_0)$ gives a $\mathrm{Beta}(a_B,b_B)$ kernel. Integration over $p\in(0,1)$ yields the ratio of Beta functions in Equation (14). **Step 3 (posterior signal).** The first two Beta moments give the result. The trial number m_r belongs to the Binomial law, not to an independent power weight; $m_r=0$ and $y_r=0$ is a legitimate no-trial observation with likelihood factor one. □

4.5.4 Negative-Binomial counts with fixed dispersion

Proposition 4.8 (Beta–negative-Binomial integration and signal choice)**.** *Use the convention*

$$p(y_r\mid p,r_r)=\frac{\Gamma(y_r+r_r)}{\Gamma(r_r)\Gamma(y_r+1)}p^{r_r}(1-p)^{y_r},\qquad r_r>0,$$

whose observation mean is $r_r(1-p)/p$. With $p\sim\mathrm{Beta}(a_0,b_0)$, let $R_B=\sum r_r$, $Y_B=\sum y_r$, and $H_B=\sum\log[\Gamma(y_r+r_r)/(\Gamma(r_r)\Gamma(y_r+1))]$. Then

$$A_B^{(0)}=e^{H_B}\frac{B(a_0+R_B,b_0+Y_B)}{B(a_0,b_0)}. \tag{15}$$

The segment posterior is $p\mid y_B\sim\mathrm{Beta}(\alpha_B,\beta_B)$ with $(\alpha_B,\beta_B)=(a_0+R_B,b_0+Y_B)$. The parameter-scale signal $m_\star(p)=p$ has ordinary Beta moments. Alternatively, the per-unit-dispersion count-mean functional $u(p)=(1-p)/p$ has

$$\mathbb{E}[u(p)\mid y_B]=\frac{\beta_B}{\alpha_B-1}\quad(\alpha_B>1),\qquad\mathbb{E}[u(p)^2\mid y_B]=\frac{\beta_B(\beta_B+1)}{(\alpha_B-1)(\alpha_B-2)}\quad(\alpha_B>2).$$

Proof. **Step 1 (sufficient statistics).** Multiplying the negative-Binomial mass functions collects the factor $e^{H_B}p^{R_B}(1-p)^{Y_B}$. **Step 2 (conjugate integration).** Combining with the Beta prior adds R_B to its first shape and Y_B to its second shape, so the posterior is $\mathrm{Beta}(\alpha_B,\beta_B)$. Its integrated normalizing-constant ratio is Equation (15). **Step 3 (distinguish the scientific signals).** For $u(p)=(1-p)/p$, integrate $p^{\alpha_B-2}(1-p)^{\beta_B}$ and $p^{\alpha_B-3}(1-p)^{\beta_B+1}$ against the Beta normalization. The resulting Beta-function ratios simplify to the displayed moments. Their finiteness requires $\alpha_B>1$ and $\alpha_B>2$, respectively. If the reported signal is instead p, no reciprocal-moment conditions arise. A target must be chosen consistently before computing Bayes curves. □

4.5.5 Beta-valued measurements and one-dimensional integration

Proposition 4.9 (Beta-response evidence and posterior curvature)**.** *Let $y_r\in(0,1)$, $y_r\mid\mu,\phi_r\sim\mathrm{Beta}(\phi_r\mu,\phi_r(1-\mu))$ with known $\phi_r>0$, and $\mu\sim\mathrm{Beta}(a_0,b_0)$. Defining*

$$\Psi_B(\mu)=\log\mathrm{BetaPDF}(\mu\mid a_0,b_0)+\sum_{r\in B}\log\mathrm{BetaPDF}(y_r\mid\phi_r\mu,\phi_r(1-\mu)),$$

the segment integrals are $A_B^{(s)} = \int_0^1 \mu^s e^{\Psi_B(\mu)}\,\mathrm{d}\mu$ *for* $s = 0, 1, 2$. *The log likelihood is strictly concave in* $\mu \in (0,1)$ *for every* $\phi_r > 0$*; if also* $a_0, b_0 \geq 1$*, the posterior log density is concave. Nevertheless, fixed-order quadrature has no universal accuracy guarantee for all admissible input data.*

Proof. **Step 1 (derive the integral).** The Beta response is not conjugate to a Beta prior on its mean. Multiplying the response densities by the prior gives $e^{\Psi_B(\mu)}$, and direct integration with factors μ^s gives the required evidence and unnormalized signal moments. **Step 2 (differentiate the likelihood).** Writing ψ_1 for the trigamma function, the terms involving μ through log-Gamma normalizers give

$$\frac{\mathrm{d}^2}{\mathrm{d}\mu^2} \log p(y_B \mid \mu, \phi_B) = -\sum_{r \in B} \phi_r^2 \{\psi_1(\phi_r \mu) + \psi_1(\phi_r(1-\mu))\} < 0.$$

All ϕ_r are strictly positive and ψ_1 is positive on positive arguments. The prior contributes $-(a_0 - 1)/\mu^2 - (b_0 - 1)/(1-\mu)^2$, which is nonpositive when $a_0, b_0 \geq 1$. **Step 3 (numerical consequence).** Log concavity alone does not bound fixed-grid integration error near either endpoint. Evaluating log-integrands after removing their maximum and comparing multiple quadrature orders with an adaptive reference supplies a practical convergence check; failed convergence must be reported rather than silently accepted. □

Missing observations. For any family above, missingness means omitting its likelihood factor by an explicit observation mask, not setting its descriptor to zero. A candidate segment with no observed values for one subject has integrated contribution one under that subject's proper prior; globally excluded segments remain controlled by the partition support instead.

5 Posterior computation over ordered partitions

Once the prior-adjusted block array $\widetilde{A}_{ij}^{(r)} = A_{ij}^{(r)} \gamma_u(i,j)$ has been constructed, the observation family no longer enters the partition recursions. This section gives the two distinct inference routes used by BayesBreak: sum-product for posterior quantities and max-sum for a joint maximizing partition.

5.1 Forward and backward partition sums

For $k \in \{0, \ldots, k_{\max}\}$ and $j \in \{0, \ldots, n\}$, let $L_{k,j}$ be the integrated score of all k-block partitions of the prefix $(0, j]$. Let $R_{k,i}$ be the corresponding score for all k-block partitions of the suffix $(i, n]$. Set

$$\begin{aligned} L_{0,0} &= 1, \qquad L_{0,j} = 0 \quad (j > 0), \\ R_{0,n} &= 1, \qquad R_{0,i} = 0 \quad (i < n). \end{aligned}$$




Figure 1. The same local scores support two algebraically related but inferentially different routes. Sum-product integrates over paths; max-sum selects one globally maximizing path. Coordinatewise modes of marginal boundary distributions need not form that path.

The sum-product recursions are

$$L_{k+1,j} = \sum_{h=k}^{j-1} L_{k,h} \widetilde{A}^{(0)}_{h,j}, \tag{16}$$

$$R_{k+1,i} = \sum_{h=i+1}^{n-k} \widetilde{A}^{(0)}_{i,h} R_{k,h}. \tag{17}$$

In implementation, Equations (16)–(17) are evaluated in log space using `logsumexp`.

Theorem 5.1 (Partition-sum identities and boundary marginals)**.** *Under Assumptions 4.1 and 4.4, for every admissible k,*

$$L_{k,n} = R_{k,0} = \sum_{0=t_0<\cdots<t_k=n} \prod_{q=1}^{k} \widetilde{A}^{(0)}_{t_{q-1},t_q}. \tag{18}$$

Moreover, for $1 \le p \le k-1$ and admissible h,

$$p(t_p = h \mid y, k) = \frac{L_{p,h} R_{k-p,h}}{L_{k,n}},$$

$$p(b_h = 1 \mid y, k) = \sum_{p=1}^{k-1} p(t_p = h \mid y, k). \tag{19}$$

Proof. **Step 1 (enumerate prefixes).** For $k = 0$ the unique empty path has weight one at $j = 0$, and no empty path has a positive endpoint. Assume $L_{k,h}$ sums the local-score products over all k-segment partitions of $(0, h]$. **Step 2 (split by the last segment).** Every $(k+1)$-segment partition of $(0, j]$ has exactly one penultimate boundary $h \in \{k, \dots, j-1\}$. Its weight is the weight of its k-segment prefix times $\widetilde{A}^{(0)}_{h,j}$. Summing over all such h gives Equation (16) and proves the prefix identity by induction. **Step 3 (enumerate suffixes).** The analogous decomposition by the first segment after i gives Equation (17). At $i = 0$, the suffix enumeration covers precisely the same set of full paths as the prefix enumeration at $j = n$. Hence $R_{k,0} = L_{k,n}$, proving

Equation (18). **Step 4 (fix an ordered boundary).** Under the event $t_p = h$, each admissible path is uniquely split into a p-segment prefix ending at h and a $(k-p)$-segment suffix beginning there. Multiplication of their total sums gives the unnormalized numerator $L_{p,h}R_{k-p,h}$. The prior factor C_k^{-1} is common to numerator and denominator and cancels, so division by $L_{k,n}$ proves the first boundary identity. **Step 5 (forget the boundary rank).** Within a strictly ordered partition, at most one of the events $\{t_p = h\}$ can occur at a given interior index h. They are disjoint as p varies. Their probabilities therefore add, yielding the boundary-event marginal in Equation (19). Normalization over h of each ordered-boundary distribution gives one; summation over h of boundary-event probabilities gives $k-1$. □

Let $p(k)$ be a segment-count prior. Because $C_k(u)$ normalizes Equation (11),

$$p(k \mid y) \propto p(k)\frac{L_{k,n}}{C_k(u)}, \qquad k = 1, \ldots, k_{\max}. \tag{20}$$

The posterior may be reported in full; choosing $\widehat{k} = \arg\max_k p(k \mid y)$ is a separate decision rule. Conditional boundary distributions in Equation (19) and boundary-event probabilities can then be computed at fixed k or averaged over $p(k \mid y)$.

5.2 Posterior moments and Bayes curves

For $r \in \{0, 1, 2\}$, define a block-covering contribution at fixed k,

$$F_{ij}^{(r)}(k) = \frac{1}{L_{k,n}} \sum_{m=1}^{k} L_{m-1,i}\widetilde{A}_{ij}^{(r)}R_{k-m,j}. \tag{21}$$

When the target at coordinate s is the segment-level quantity $m_\star(\theta)$, its conditional posterior moments are

$$\mathbb{E}[m_\star(\theta_s)^r \mid y, k] = \sum_{i<s\le j} F_{ij}^{(r)}(k). \tag{22}$$

The posterior mean Bayes curve uses $r = 1$; the $r = 2$ result gives pointwise posterior variance after subtracting the squared mean.

> **Proposition 5.2** (Block-covering decomposition)**.** *If every admissible partition assigns each coordinate to exactly one block and $A_{ij}^{(r)}$ is the integrated rth moment numerator for that block, then Equation (22) is the exact fixed-k posterior moment of the segment-level target at s.*

Proof. **Step 1 (uniqueness of the covering block).** Fix $s \in \{1, \ldots, n\}$. In every admissible partition there is exactly one segment $(i, j]$ for which $i < s \le j$; the corresponding segment position m is also unique. **Step 2 (integrate a fixed covering block).** Conditioning on $(i, j]$ occupying position m, all earlier segments contribute the prefix sum $L_{m-1,i}$. The covering block contributes $\widetilde{A}_{ij}^{(r)}$, because its signal functional is raised to the rth power before integration. All later segments contribute $R_{k-m,j}$. **Step 3 (sum without duplication).** Summing over positions m and blocks $(i, j]$ covering s enumerates each full k-segment path once, multiplied by its integrated target contribution. This gives the numerator of the posterior expectation. **Step 4 (normalize).** The denominator $L_{k,n}$ is the sum of unweighted path contributions by Theorem 5.1.

Division proves Equation (22). When $r = 2$, subtracting the square of the $r = 1$ result yields pointwise posterior variance, with no claim of a simultaneous credible band. □

5.3 Joint MAP segmentation

Define local log scores $\ell_{ij} = \log \widetilde{A}^{(0)}_{ij}$ and max-sum messages

$$M_{k,j} = \max_{h \in \{k-1,\dots,j-1\}} \{M_{k-1,h} + \ell_{h,j}\}, \quad M_{0,0} = 0, \tag{23}$$

with $M_{0,j} = -\infty$ for $j > 0$. Store a deterministic maximizing predecessor $h^\star(k, j)$. Starting at (k, n) and following these predecessors recovers a complete boundary vector.

Theorem 5.3 (Correctness of max-sum backtracking)**.** *For fixed k, the boundary vector returned by Equation (23) and its stored predecessors belongs to*

$$\arg \max_{0=t_0<\cdots<t_k=n} \sum_{q=1}^{k} \ell_{t_{q-1},t_q}. \tag{24}$$

Consequently, it is a joint MAP segmentation conditional on k. Ties are resolved only by the declared predecessor convention.

Proof. **Step 1 (base case).** For a one-segment prefix $(0, j]$, the only admissible path consists of the block $(0, j]$. Hence $M_{1,j} = \ell_{0,j}$ is its maximum score, and its stored predecessor is 0. **Step 2 (induction hypothesis).** Assume that for each feasible h, $M_{k-1,h}$ is the maximum log score among all $(k-1)$-segment partitions ending at h. **Step 3 (optimal substructure).** Every k-segment path ending at j has a unique penultimate endpoint $h \in \{k-1, \dots, j-1\}$. Its score is the prefix path score plus $\ell_{h,j}$. At any fixed h, the best possible prefix is $M_{k-1,h}$ by the hypothesis. Therefore maximizing over h gives exactly Equation (23), the best k-segment score. **Step 4 (recover a feasible maximizing path).** Each predecessor $h^\star(k, j)$ attains the recurrence maximum and satisfies $h^\star(k, j) < j$. Starting from (k, n) and following these predecessors decreases the endpoint and block count by one each step until $(0, 0)$. The resulting ordered boundary vector attains $M_{k,n}$ and maximizes Equation (24). This is conditional on k; joint optimization over k also needs the count-prior offset. □

Marginal modes are not a joint MAP path

The vector formed from $\arg\max_h p(t_p = h \mid y, k)$ for each p optimizes separate marginal decisions. Its entries can be incompatible or can have lower joint posterior probability than another partition. BayesBreak therefore reports marginal modes as diagnostics and reserves "joint MAP" for Theorem 5.3.

5.4 Decision-theoretic segmentation summaries

Posterior summaries answer different decision questions. The distinction is not a formatting convention: the marginal posterior of k sums over segmentations, while joint MAP scoring maximizes over them.

Proposition 5.4 (Three non-equivalent posterior modes)**.** *Define*

$$\widehat{k}_{\mathrm{marg}} \in \arg\max_{k} \{\log p(k) + \log L_{k,n} - \log C_k\},$$

$$\widehat{t}_{\mathrm{cond}} \in \arg\max_{t \in \mathcal{T}_{\widehat{k}_{\mathrm{marg}}}} \sum_{q=1}^{\widehat{k}_{\mathrm{marg}}} \ell_{t_{q-1},t_q},$$

$$(\widehat{k}_{\mathrm{joint}}, \widehat{t}_{\mathrm{joint}}) \in \arg\max_{k, t \in \mathcal{T}_k} \left\{ \log p(k) - \log C_k + \sum_{q=1}^{k} \ell_{t_{q-1},t_q} \right\}.$$

Here $\ell_{ij} = \log \widetilde{A}^{(0)}_{ij}$. The first is a marginal-MAP segment count, the second a conditional joint-MAP partition at that count, and the third a global joint-MAP count–partition pair. In general they are distinct decisions.

Proof. **Step 1 (marginalize partitions).** The evidence for a fixed count is $p(y \mid k) = L_{k,n}/C_k$, so Bayes' rule gives the first maximization after discarding its constant normalizer. **Step 2 (optimize within a selected count).** Conditional on k, the factors $p(k)$ and C_k^{-1} are constant across admissible partitions. Therefore max-sum backtracking solves the second maximization for $k = \widehat{k}_{\mathrm{marg}}$. **Step 3 (optimize jointly).** If k also varies, the complete joint score retains both $\log p(k)$ and $-\log C_k$, giving the third maximization. **Step 4 (show the distinction).** For each k, the marginal score uses a *sum* $L_{k,n}$ over all paths, whereas the best-path score uses their *maximum*. Counts with many moderately probable paths can have greater marginal support than counts with one sharply preferred path, so no identity between the maximizing counts follows. □

Proposition 5.5 (Bayes actions for full-partition and coordinate losses)**.** *For fixed k, 0–1 loss on the entire ordered vector $t = (t_1, \ldots, t_{k-1})$ is minimized by a joint MAP partition. For coordinatewise Hamming loss $\mathcal{L}(a,t) = \sum_{q=1}^{k-1} \mathbf{1}\{a_q \neq t_q\}$, constrained to admissible ordered actions $a_1 < \cdots < a_{k-1}$, the Bayes action instead solves*

$$\max_{1 \le a_1 < \cdots < a_{k-1} \le n-1} \sum_{q=1}^{k-1} p(t_q = a_q \mid y, k). \tag{25}$$

This constrained optimization has its own additive dynamic program; independent coordinatewise marginal modes need not form an admissible action.

Proof. **Step 1 (0–1 loss).** Its posterior risk at action a is $1 - p(t = a \mid y, k)$, minimized exactly by maximizing the joint posterior mass. **Step 2 (Hamming loss).** By linearity of conditional expectation, its posterior risk is $(k-1) - \sum_q p(t_q = a_q \mid y, k)$. Minimizing risk over valid actions is therefore equivalent to Equation (25); dropping the ordering constraint would solve a different optimization problem. **Step 3 (dynamic program).** Set $D_{1,h} = p(t_1 = h \mid y, k)$ and recursively

$$D_{q,h} = p(t_q = h \mid y, k) + \max_{g < h} D_{q-1,g}$$

on valid index ranges, retaining predecessors. Backtracking from the best valid terminal state yields the constrained Bayes action; prefix maxima implement each layer efficiently. **Step 4 (explicit counterexample).** For $n = 7, k = 3$, assign posterior masses $0.17, 0.18, 0.20, 0.20, 0.125, 0.125$ to the admissible pairs $(1,3), (2,3), (4,5), (4,6), (1,5), (2,6)$, respectively, and zero elsewhere.

They sum to one. The marginal mode of t_1 is 4 with probability 0.40, while that of t_2 is 3 with probability 0.35, producing the impossible vector $(4, 3)$. The joint modes are $(4, 5)$ and $(4, 6)$, each of mass 0.20; the constrained Hamming calculation must respect ordering rather than export $(4, 3)$. □

The output interface reports each object by its complete name. Posterior-mean segment levels *conditional on* an exported MAP partition are likewise distinct from posterior-mean Bayes curves averaged over the segmentation distribution.

5.5 Complexity and numerical invariants

If cumulative sufficient statistics make each local conjugate score constant time, there are $\Theta(n^2)$ blocks to precompute. Sum-product and max-sum each inspect $\Theta(k_{\max} n^2)$ predecessor transitions. Values and backpointers require $\Theta(k_{\max} n)$ working storage, while retaining all block-cover terms can require $\Theta(k_{\max} n^2)$ storage.

> **Proposition 5.6** (Complexity)**.** *When cumulative sufficient statistics make each conjugate segment calculation constant time, segment precomputation costs $\Theta(n^2)$ time and storage; either partition recursion costs $\Theta(k_{\max} n^2)$ time. For a nonconjugate block routine with per-block cost T_{block}, local precomputation costs $\Theta(n^2 T_{\text{block}})$ and the global recursions are unchanged.*

Proof. **Step 1 (candidate-segment count).** There are $n(n+1)/2$ nonempty contiguous segments $(i, j]$ with $0 \le i < j \le n$. With constant-time evaluation from prefix statistics, filling the triangular local score array therefore takes $\Theta(n^2)$ time and memory. **Step 2 (count predecessor transitions).** At a feasible state (k, j), sum-product or max-sum examines $j - k + 1$ predecessor endpoints in the unconstrained dense case. For a range of counts $1 \le k \le k_{\max} \le n$, the exact transition total is of order $\sum_{k=1}^{k_{\max}} \sum_{j=k}^{n} (j - k + 1)$, which is $\mathcal{O}(k_{\max} n^2)$ and attains that order for the usual regimes where $k_{\max}$ does not collapse to an empty or singleton state set. **Step 3 (separate local approximation cost).** If each local score costs T_{block} operations rather than a constant, filling the array costs $\Theta(n^2 T_{\text{block}})$. Neither sum-product nor max-sum changes its predecessor traversal as a result. These are worst-case accounting statements; finite-range measured timing slopes need not equal the asymptotic exponents. □


Four numerical identities are checked in every implementation: the normalized segment-count posterior sums to one; forward and backward total evidence agree; conditional boundary-event probabilities sum to $k - 1$; and the backtracked path score equals the terminal max-sum score within numerical tolerance. These checks test algebraic implementation rather than empirical model quality.

6 Irregular designs and hierarchical extensions

The ordered-partition recursions remain valid when prior factors or multiple-sequence likelihoods can be expressed segment by segment. This section gives the corresponding models for irregular coordinates, shared changepoints, observed groups, and latent groups.

6.1 Irregular designs: segment cohesion and boundary hazard

Irregular coordinates can enter the partition prior in two statistically distinct ways. A complete-segment cohesion $c_u(i, j)$ assigns prior mass according to the physical span of $(i, j]$; an interior-boundary hazard $h_u(j)$ assigns prior mass to a candidate split at u_j. They can appear together in Equation (11), but one does not imply the other.



$$p(t \mid k, u) \propto \prod_{q=1}^{k} c_u(t_{q-1}, t_q) \prod_{q=1}^{k-1} h_u(t_q)$$

Figure 2. Design-dependent partition prior. Segment cohesion and interior-boundary hazard represent different prior assumptions. The terminal endpoint receives no boundary-hazard factor.

Theorem 6.1 (Local prior factors preserve exact inference). *Replacing* $A_{ij}^{(r)}$ *by* $\widetilde{A}_{ij}^{(r)} = A_{ij}^{(r)} c_u(i, j) h_u(j)^{\mathbb{1}\{j<n\}}$ *causes the sum-product recursion to enumerate exactly the unnormalized joint weights in Equation* (11)*. Dividing the terminal partition sum by* $C_k(u)$ *yields the proper fixed-*k *marginal likelihood.*

Proof. **Step 1 (factor the intended partition weight).** For fixed k and $t = (t_0, \ldots, t_k)$, multiply its independent segment marginal likelihoods by the conditional prior in Equation (11). **Step 2 (push each factor into a segment).** Every segment ending at $t_q < n$ contributes the prior factor $c_u(t_{q-1}, t_q) h_u(t_q)$, while the terminal segment ending at n contributes only $c_u(t_{k-1}, n)$. Thus $\prod_q \widetilde{A}^{(0)}_{t_{q-1}, t_q}$ is exactly the likelihood product times the *unnormalized* prior factor. **Step 3 (sum over all partitions).** Theorem 5.1 enumerates these path products once each, so $L_{k,n}$ is their unnormalized sum. Running the same recurrence with $A_{ij}^{(0)} = 1$ enumerates exactly the prior weights and returns $C_k(u)$. **Step 4 (normalize once, not once per block).** Multiplication by $C_k(u)^{-1}$ in the fixed-count prior is common across partitions; hence $p(y \mid k) = L_{k,n}/C_k(u)$. Applying the same factor inside each block instead would produce an incorrect power of the prior normalizer. □

A boundary-process interpretation applies only to the hazard factor. Let $\Lambda_j = \int_{u_{j-1}}^{u_j} \lambda(v)\, \mathrm{d}v$ for an inhomogeneous Poisson process. Interval occupancies are independent Bernoulli variables with probability $1 - e^{-\Lambda_j}$. Conditional on exactly m occupied candidate intervals, the subset probability is proportional to the product of Bernoulli odds, so the compatible local factor is

$$h_u(j) \propto \frac{1 - e^{-\Lambda_j}}{e^{-\Lambda_j}} = e^{\Lambda_j} - 1. \tag{26}$$

Constant intensity and equal-width intervals recover an index-uniform fixed-count boundary prior when $c_u \equiv 1$. Without conditioning on the number of occupied intervals, both occupied and

unoccupied Bernoulli factors are part of the prior. Equation (26) does not justify multiplying a segment likelihood by its physical length; a duration-dependent cohesion is a separate prior choice.

6.2 Renewal interpretation and refinement of an irregular design

The local prior factorization admits a renewal interpretation, but its conditioning matters. A conditional prior on partitions given a separately sampled count k does not automatically induce a prior on the count.

Proposition 6.2 (Renewal-length equivalence conditional on k)**.** *On an index grid with $0 = t_0 < t_1 < \cdots < t_k = n$, let the positive segment lengths $d_q = t_q - t_{q-1}$ have an unnormalized renewal mass $g(d_q) \geq 0$. Conditional on $\sum_q d_q = n$ and a fixed count k, the boundary distribution is*

$$p(t \mid k) = \frac{1}{C_k} \prod_{q=1}^{k} g(t_q - t_{q-1}), \qquad C_k = \sum_{t \in \mathcal{T}_k} \prod_{q=1}^{k} g(t_q - t_{q-1}).$$

For geometric $g(d) = \rho(1-\rho)^{d-1}$, $0 < \rho < 1$, the product is constant over $\mathcal{T}_k$, and thus $p(t \mid k)$ is uniform regardless of ρ.

Proof. **Step 1 (one-to-one correspondence).** Each ordered partition uniquely determines a composition $(d_1, \ldots, d_k)$ of n into positive integers, and conversely the partial sums of each such composition recover the boundaries. **Step 2 (condition the renewal mass).** The unnormalized composition weight is $\prod_q g(d_q)$. Summing those weights over every admissible composition defines C_k, so normalizing yields the displayed conditional distribution. **Step 3 (geometric specialization).** For $g(d) = \rho(1-\rho)^{d-1}$, the product equals $\rho^k(1-\rho)^{\sum_q d_q - k} = \rho^k(1-\rho)^{n-k}$, independent of t at fixed k. The factor cancels against C_k and cannot push the separately specified segment-count prior $p(k)$ toward one or toward n. □

Proposition 6.3 (Score preservation under grid refinement)**.** *Suppose a candidate-coordinate grid is refined by adding locations carrying no new observations. Embed an original partition into the refined grid by retaining all of its physical segment endpoints and placing no boundary at a newly inserted location. Each inherited block then has the same likelihood integral; any cohesion or hazard factor depending only on unchanged physical endpoints and retained boundary weights has the same unnormalized prior contribution. The fully normalized partition posterior need not remain unchanged if new boundary locations are made admissible.*

Proof. **Step 1 (likelihood equality).** Each inherited refined block contains exactly the original observations and descriptors, so its likelihood product and prior-integrated segment likelihood agree with those of the original block. **Step 2 (prior equality for inherited paths).** Preserving physical endpoints preserves a cohesion depending only on those endpoints; by hypothesis, each retained interior boundary also retains its hazard factor. Multiplying across blocks thus gives the same unnormalized path weight. **Step 3 (why posterior equality fails).** Refinement that admits an additional cut creates extra partitions in the sum defining C_k and in the data-dependent partition sum. Even if every inherited path has unchanged weight, the two new normalizers generally differ. The equality is for inherited path scores, not for the whole posterior. □

Physical interval conventions. Observation locations $x_1 < \cdots < x_n$ and candidate boundary coordinates $u_0 < \cdots < u_n$ are distinct objects. Interior cell edges may be set to $u_i = (x_i + x_{i+1})/2$ for $1 \le i < n$ and endpoints determined from the scientific measurement domain. The complete-segment span $u_j - u_i$ is not the same object as the observation gap $x_{i+1} - x_i$. An inhomogeneous Poisson boundary process can supply hazard factors through integrated intensities, but an interval-length cohesion does not automatically produce that process.

6.3 Shared changepoints across multiple sequences

Suppose S aligned sequences share a boundary vector while retaining sequence-specific segment parameters and priors. Conditional independence gives a pooled segment marginal likelihood.




Figure 3. Hierarchical common-changepoint model with sequence-specific segment parameters. Conditional independence makes the sequence-specific segment marginal likelihoods multiply.

Assumption 6.4 (Common candidate set). The sequences are aligned to a common ordered candidate set. Every candidate partition considered by the common-changepoint model is admissible for every sequence after the declared missing-data or exposure convention. Conditional on the common partition, sequences and their segment parameters are independent, and all sequence-specific segment marginal likelihoods are finite and positive on the common support.

Theorem 6.5 (Exact pooling under a common partition)**.** *Under Assumption 6.4, the pooled segment marginal likelihood is*

$$A_{ij}^{(0),\text{pool}} = \prod_{s=1}^{S} A_{ij}^{(0,s)}, \qquad \ell_{ij}^{\text{pool}} = \sum_{s=1}^{S} \ell_{ij}^{(s)}. \tag{27}$$

Using this array in Section 5 gives the exact posterior distribution over the common partition. Conditional sequence-specific parameter posteriors are recovered from the original sequence-

specific conjugate updates.

Proof. **Step 1 (integrate independently within a segment).** For sequence s, write $\theta_{B,s}$ for its parameter on a candidate shared segment $B = (i, j]$. By the independent segment priors, integrating its observations over $\theta_{B,s}$ yields $A_{ij}^{(0,s)}$. **Step 2 (factor across sequences).** The model assumes conditional independence across sequences given the shared partition, and the subject-specific segment parameters are independent a priori. Tonelli's theorem for nonnegative integrands permits exchanging product and integration, so

$$\int \prod_{s=1}^{S} p(y_B^{(s)} \mid \theta_{B,s}) p_s(\theta_{B,s}) \, \mathrm{d}\theta_{B,1:S} = \prod_{s=1}^{S} A_{ij}^{(0,s)}.$$

Taking logarithms gives the additive log-score formula in Equation (27). **Step 3 (apply the unchanged partition recursion).** Multiplication over segments produces the exact integrated joint likelihood conditional on every shared partition. Applying the same partition prior and the sum-product recursion therefore yields its exact boundary and count posterior. Given a partition, conjugate posterior calculations for each sequence's segment parameters remain separate; they are not automatically a single joint conjugate density integrated over all partitions. □

Known groups are handled by applying Equation (27) separately within each observed group. The following asymptotic statement is restricted to a finite candidate set and requires information to accumulate across sequences.

Theorem 6.6 (Finite-candidate consistency for a common partition)**.** *Let $\mathcal{T}$ be finite and contain a data-generating candidate $t^\star$. For sequence s, let $M_s(t) > 0$ be the integrated score used by the common-partition model. Assume independent sequences, $p(t^\star) > 0$, and for each $t \neq t^\star$:*

1. $\mathbb{E}|\log M_s(t^\star) - \log M_s(t)| < \infty$*;*
2. *a strong law holds for the centered log-score differences; and*
3. $\liminf_{S\to\infty} S^{-1} \sum_{s=1}^{S} \mathbb{E}[\log M_s(t^\star) - \log M_s(t)] = \delta_t > 0.$

Then $p(t^\star \mid y^{(1:S)}) \to 1$ almost surely, and every joint MAP segmentation is eventually $t^\star$ almost surely.

Proof. **Step 1 (posterior ratio for an incorrect candidate).** Fix $t \neq t^\star$ and put $D_s(t) = \log M_s(t^\star) - \log M_s(t)$. Bayes' rule gives

$$\frac{p(t \mid y^{(1:S)})}{p(t^\star \mid y^{(1:S)})} = \frac{p(t)}{p(t^\star)} \exp\left\{ -\sum_{s=1}^{S} D_s(t) \right\}.$$

If $p(t) = 0$, the ratio is identically zero and needs no further argument. **Step 2 (accumulation of log-score separation).** The assumed strong law makes $S^{-1} \sum_s [D_s(t) - \mathbb{E}D_s(t)] \to 0$ almost surely. The strictly positive limiting lower bound on the mean expected difference then gives $\liminf_{S\to\infty} S^{-1} \sum_s D_s(t) \geq \delta_t > 0$ almost surely. **Step 3 (vanishing posterior odds).** The exponential factor in Step 1 tends to zero at an exponential rate in S, and the fixed prior odds cannot prevent this. Hence every incorrect candidate has posterior mass asymptotically negligible relative to $t^\star$. **Step 4 (finite candidate set).** There are only finitely many $t \in \mathcal{T}$, so the sum of incorrect posterior ratios tends to zero. Normalization yields $p(t^\star \mid y^{(1:S)}) \to 1$, and eventually its

posterior mass exceeds that of every incorrect t, proving the joint MAP conclusion. The theorem requires an informative average across growing numbers of sequences; adding one informative sequence and arbitrarily many uninformative ones does not satisfy the positive-gap condition. □

One fixed informative sequence is insufficient for this theorem because its contribution vanishes in the average as S increases. The assumption requires an accumulating average expected log-score gap.

6.4 Finite latent-group template model

For sequence s and deterministic segmentation template $\tau = (k, t)$, consider the score

$$S_s(\tau) = \frac{p(k)}{C_k} \prod_{q=1}^{k} A^{(0,s)}_{t_{q-1},t_q} \gamma_u(t_{q-1}, t_q). \tag{28}$$

These subject-template scores are not required to integrate to one over the data space. We therefore define the finite criterion

$$\mathcal{F}(\pi, \tau) = \sum_{s=1}^{S} \log \left\{ \sum_{g=1}^{G} \pi_g S_s(\tau_g) \right\},$$
$$\pi_g \geq 0, \qquad \sum_g \pi_g = 1. \tag{29}$$

This is a latent-group clustering criterion. It is not interpreted as a finite-mixture likelihood unless each $S_s(\tau_g)$ is a normalized sampling density.

For row-stochastic auxiliary weights r_{sg}, Jensen's inequality gives

$$\mathcal{F}(\pi, \tau) \geq \mathcal{Q}(r, \pi, \tau),$$
$$\mathcal{Q}(r, \pi, \tau) = \sum_{s,g} r_{sg} \big[\log \pi_g + \log S_s(\tau_g) - \log r_{sg} \big], \tag{30}$$

with equality at

$$r_{sg} = \frac{\pi_g S_s(\tau_g)}{\sum_h \pi_h S_s(\tau_h)}. \tag{31}$$

The mixing-weight update is $\pi_g = S^{-1} \sum_s r_{sg}$. Let $n_g = \sum_s r_{sg}$. For a candidate segment count k, substitution of Equation (28) into the minorizer gives the count offset $n_g\{\log p(k) - \log C_k\}$ and the max-sum local score

$$\ell^{(g)}_{ij} = \sum_{s=1}^{S} r_{sg} \log A^{(0,s)}_{ij} + n_g \log \gamma_u(i, j). \tag{32}$$

The template update solves the fixed-k max-sum problem and then compares the offset-adjusted terminal values across the allowed segment counts.

> **Theorem 6.7** (Monotonicity of exact latent-group updates)**.** *An iteration that sets r by Equation (31), maximizes $\mathcal{Q}$ over π, and exactly maximizes every responsibility-weighted template criterion using Equation (32) and its segment-count offset over the declared finite*

candidate set cannot decrease $\mathcal{F}$.

Proof. **Step 1 (construct a tight Jensen minorizer).** For any row-stochastic r, Jensen's inequality applied to the positive sum over group templates gives $\mathcal{F}(\pi, \tau) \geq \mathcal{Q}(r, \pi, \tau)$. Choosing r as in Equation (31) at the current $(\pi^{\text{old}}, \tau^{\text{old}})$ makes the inequality an equality. **Step 2 (update group weights).** At fixed r and templates, the π-dependent terms are $\sum_g n_g \log \pi_g$. Maximization on the probability simplex gives $\pi_g = n_g/S$ for positive group masses, under the declared boundary convention for empty groups. This cannot decrease $\mathcal{Q}$. **Step 3 (update templates exactly).** For group g, substitute the score definition in Equation (28) into $\sum_s r_{sg} \log S_s(\tau_g)$. Every segment receives $\sum_s r_{sg} \log A_{ij}^{(0,s)} + n_g \log \gamma_u(i,j)$, and the count offset is $n_g[\log p(k) - \log C_k]$. Max-sum dynamic programming and comparison across all admissible k globally maximize this coordinate of the minorizer. **Step 4 (chain the inequalities).** Let $(\pi^{\text{new}}, \tau^{\text{new}})$ denote the updated parameters. Then

$$\mathcal{F}(\pi^{\text{new}}, \tau^{\text{new}}) \geq \mathcal{Q}(r, \pi^{\text{new}}, \tau^{\text{new}}) \geq \mathcal{Q}(r, \pi^{\text{old}}, \tau^{\text{old}}) = \mathcal{F}(\pi^{\text{old}}, \tau^{\text{old}}).$$

The argument establishes monotonicity of the stated finite *score-clustering* criterion, not maximization of a normalized generative mixture likelihood or global convergence across restarts. □

Proposition 6.8 (Permutation symmetry of the finite template criterion)**.** *For every permutation σ of the G group indices, simultaneously replacing (π_g, τ_g) by $(\pi_{\sigma(g)}, \tau_{\sigma(g)})$ leaves $\mathcal{F}$ unchanged. Consequently, the numerical labels of fitted templates are not identified by this criterion; distinct groups with identical templates can also produce non-uniqueness beyond label permutations.*

Proof. **Step 1 (permute inside each sequence score).** For subject s, relabeling the terms of the finite sum gives $\sum_g \pi_{\sigma(g)} S_s(\tau_{\sigma(g)}) = \sum_g \pi_g S_s(\tau_g)$, because σ is a bijection. **Step 2 (preserve the objective).** Taking the logarithm and summing the unchanged subject terms leaves Equation (29) invariant. **Step 3 (non-uniqueness).** If a maximizing set of templates is relabeled, it remains maximizing. If two templates coincide, their contributions can also be aggregated with changed mixture weights without changing their combined term, giving an additional source of non-identifiability. A plotting convention may canonically order labels, but it does not establish parameter identifiability or optimality across restarts. □

Template-label permutations, coincident templates, and an oversized G can make the optimum non-unique. No finite-mixture identifiability or generative-model consistency claim is made. Multiple initializations must be compared using the criterion evaluated at their *final* iterates, rather than a value recorded before the last update.

7 Approximate segment integration, posterior prediction, and evaluation

For a nonconjugate segment model, BayesBreak requires a numerical approximation to the segment marginal likelihood. Posterior prediction requires the posterior predictive distribution associated with the fitted observation model. This section states the error condition under which segmentwise approximation error can be propagated through the partition recursions and then defines the prediction and boundary-evaluation rules used in the empirical analysis.

7.1 Propagation of segment marginal-likelihood error

Let $\ell_{ij} = \log A_{ij}^{(0)}$ be a reference log segment marginal likelihood and let $\widehat{\ell}_{ij}$ be the value returned by a numerical method such as Laplace approximation, the Jaakkola–Jordan bound, expectation propagation, Pólya–Gamma mean-field approximation, or adaptive quadrature.

Assumption 7.1 (Uniform error bound over admissible segments). There exists $\varepsilon \geq 0$ such that

$$|\widehat{\ell}_{ij} - \ell_{ij}| \leq \varepsilon \tag{33}$$

for every segment $(i, j]$ that can occur in a partition with $k \leq k_{\max}$. The numerical analysis must identify the reference value and, when applicable, separate optimization residual, truncation or tail error, quadrature error, and discrepancy from the reference calculation.

Theorem 7.2 (Stability of marginal likelihoods and posterior odds)**.** *Under Assumption 7.1, every k-segment forward or backward quantity satisfies*

$$\begin{aligned} e^{-k\varepsilon} L_{k,j} \leq \widehat{L}_{k,j} \leq e^{k\varepsilon} L_{k,j}, \\ e^{-k\varepsilon} R_{k,i} \leq \widehat{R}_{k,i} \leq e^{k\varepsilon} R_{k,i}. \end{aligned} \tag{34}$$

Consequently, for $k, k' \leq k_{\max}$,

$$\left| \log \frac{\widehat{p}(k \mid y)}{\widehat{p}(k' \mid y)} - \log \frac{p(k \mid y)}{p(k' \mid y)} \right| \leq (k + k')\varepsilon, \tag{35}$$

and, for two locations h, h' of the same ordered boundary t_p under fixed k,

$$\left| \log \frac{\widehat{p}(t_p = h \mid y, k)}{\widehat{p}(t_p = h' \mid y, k)} - \log \frac{p(t_p = h \mid y, k)}{p(t_p = h' \mid y, k)} \right| \leq 2k\varepsilon. \tag{36}$$

Proof. **Step 1 (pathwise perturbation).** For any admissible partition of k blocks, the approximate-to-exact path-weight ratio is $\exp\{\sum_{q=1}^{k} \delta_{t_{q-1}, t_q}\}$, with every summand bounded in absolute value by ε. Consequently that ratio lies between $e^{-k\varepsilon}$ and $e^{k\varepsilon}$. **Step 2 (sum over positive paths).** Multiply the exact weight of each path by the same lower and upper bounds and sum. Positivity preserves both inequalities. The prefix and suffix path sets contain exactly k segments each, proving the two message sandwich bounds in Equation (34). **Step 3 (count posterior odds).** For distinct counts k, k', the ratio $p(k \mid y)/p(k' \mid y)$ equals $[p(k)/C_k]L_{k,n}$ divided by $[p(k')/C_{k'}]L_{k',n}$. The count priors and partition normalizers are unchanged by local numerical approximation. The two summed-path errors therefore add in the log ratio, yielding the bound $(k + k')\varepsilon$. **Step 4 (ordered boundary odds).** The unnormalized weight for $t_p = h$ is $L_{p,h}R_{k-p,h}$; it is a sum of paths each with exactly $p + (k - p) = k$ blocks. Its log perturbation is bounded by $k\varepsilon$. Comparing this weight at h and h' doubles the maximum error, proving the $2k\varepsilon$ statement. **Step 5 (scope).** The argument compares only positive exact and approximate scores on the same admissible support. A bound on typical blocks, a per-method asymptotic formula, or a large observed maximum score discrepancy does not establish the uniform hypothesis. □

Corollary 7.3 (Consequences for probabilities and modal locations)**.** *If finite unnormalized weights satisfy $|\log \widehat{w}_a - \log w_a| \leq \eta$, then their normalized probabilities satisfy $e^{-2\eta} \leq \widehat{p}_a/p_a \leq$*

$e^{2\eta}$ *and* $\|\widehat{p} - p\|_{\mathrm{TV}} \le \min\{1, e^{2\eta} - 1\}$. *Moreover, a fixed-*$k$ *modal boundary separated from every alternative by exact log-odds margin* Δ *is unchanged whenever* $\varepsilon < \Delta/(2k)$.

Proof. **Step 1 (bound the perturbed normalizer).** Write $\widehat{w}_a = w_a e^{\delta_a}$ with $|\delta_a| \le \eta$. Let $p_a = w_a / \sum_b w_b$. Then $\sum_b p_b e^{\delta_b} \in [e^{-\eta}, e^{\eta}]$. **Step 2 (normalized probability ratios and total variation).** Since $\widehat{p}_a / p_a = e^{\delta_a} / \sum_b p_b e^{\delta_b}$, this ratio lies in $[e^{-2\eta}, e^{2\eta}]$. Therefore $|\widehat{p}_a - p_a| \le (e^{2\eta} - 1) p_a$. Summing its positive part bounds total variation by $\min\{1, e^{2\eta} - 1\}$. **Step 3 (preserve the maximizing boundary).** Suppose $h^\star$ exceeds every rival h by exact log-odds margin at least Δ. Theorem 7.2 changes each pairwise log-odds difference by at most $2k\varepsilon$. If $2k\varepsilon < \Delta$, all perturbed differences remain positive, so $h^\star$ remains the unique modal boundary location. □

Proof obligation 7.4 (Error control for individual numerical methods). This paper does not establish a uniform convergence rate for Laplace approximation, the Jaakkola–Jordan bound, expectation propagation, Pólya–Gamma mean-field approximation, or adaptive quadrature over all admissible segments. To apply Theorem 7.2, a nonconjugate analysis must establish Assumption 7.1 under conditions appropriate to the observation family, parameter domain, segment support, initialization, convergence event, and tail behavior. Without that step, the dynamic program is exact for the supplied approximate score array, but the theorem does not compare the resulting posterior with the posterior under the intended segment model.

7.2 Nonconjugate block calculations: what each approximation establishes

For a canonical Binomial–logistic block with a Gaussian prior on its intercept, the exact local evidence is a one-dimensional integral. It provides a numerical reference against which deterministic approximations can be tested before the resulting segment scores enter the same partition recursion. A local approximation formula, its variational lower bound, and a rigorous numerical error bound are three different objects.

Lemma 7.5 (Finite, unique logistic-block mode under a proper normal prior)**.** *Let* $y_r \mid \theta \sim \mathrm{Binom}(m_r, \sigma(\theta))$ *independently, with* $m_r \ge 0$, $\sigma(\theta) = (1 + e^{-\theta})^{-1}$, *and* $\theta \sim \mathcal{N}(\nu, \rho^2)$ *for finite* $\rho^2 > 0$. *Then the block log posterior is strictly concave and has a unique finite maximizer, including blocks containing only successes or only failures.*

Proof. **Step 1 (posterior objective).** Apart from its Binomial coefficients, the log posterior is

$$\Psi_B(\theta) = Y_B \theta - M_B \log(1 + e^{\theta}) - \frac{(\theta - \nu)^2}{2\rho^2} + \text{constant},$$

where $Y_B = \sum y_r$ and $M_B = \sum m_r$. **Step 2 (strict curvature).** Direct differentiation gives

$$\Psi_B''(\theta) = -M_B \sigma(\theta)\{1 - \sigma(\theta)\} - \rho^{-2} < 0$$

for every finite θ, even when $M_B = 0$. **Step 3 (existence, not just uniqueness).** The Binomial log-likelihood is bounded above by the sum of its log combinatorial coefficients. The negative quadratic prior term tends to $-\infty$ as $|\theta| \to \infty$. Hence Ψ_B is coercive, attains a finite maximum by continuity, and strict concavity makes that maximizer unique. This explicit tail condition is why the conclusion does not follow from strict concavity alone under an arbitrary prior. □

7.2.1 Laplace evaluation and its limitations

Let $\widehat{\theta}_B$ be the mode and $H_B = -\Psi_B''(\widehat{\theta}_B) > 0$. The first-order Laplace approximation is

$$\log \widehat{A}_B^{\mathrm{Lap}} = \Psi_B(\widehat{\theta}_B) + \tfrac{1}{2}\log(2\pi) - \tfrac{1}{2}\log H_B. \tag{37}$$

This is a local second-order approximation to the integral, *not* a nonasymptotic upper bound on the absolute log-evidence error. A controlled asymptotic remainder would require derivatives beyond second order, concentration of the integral near the mode, and bounds on the tails outside that neighborhood. Numerical evaluation should use safeguarded Newton steps, check finite curvature and convergence, and fall back to adaptive one-dimensional integration on failure. The software's local curvature diagnostic does not by itself establish Assumption 7.1.

7.2.2 Jaakkola–Jordan lower bound

Proposition 7.6 (Valid logistic tangent bound and block-evidence lower bound)**.** *For $\xi > 0$ put $\lambda(\xi) = \tanh(\xi/2)/(4\xi)$ and $\sigma(x) = (1 + e^{-x})^{-1}$. Then, for every real x,*

$$\log \sigma(x) \geq \log \sigma(\xi) + \frac{x - \xi}{2} - \lambda(\xi)(x^2 - \xi^2). \tag{38}$$

For a $\mathrm{Binom}(m_r, \sigma(\theta))$ observation, this quadratic lower bound has linear coefficient $y_r - m_r/2$ and quadratic coefficient $-m_r\lambda(\xi_r)$. Combining these terms with a Gaussian prior and integrating the resulting Gaussian kernel gives a deterministic lower *bound on $\log A_B^{(0)}$; this bound alone supplies no upper bound on the unknown error gap.*

Proof. **Step 1 (rewrite the log sigmoid).** The identity $\log \sigma(x) = x/2 - \log 2 - \log\cosh(x/2)$ separates a linear term from an even function of x. **Step 2 (bound the even term).** Define $f(v) = \log\cosh(\sqrt{v}/2)$ for $v \geq 0$. Its derivative for $v > 0$ is $f'(v) = \tanh(\sqrt{v}/2)/(4\sqrt{v})$, which decreases with v. Thus f is concave and lies below its tangent at $v = \xi^2$: $f(x^2) \leq f(\xi^2) + \lambda(\xi)(x^2 - \xi^2)$. **Step 3 (recover the inequality).** Substitute this tangent inequality into the expression for $\log \sigma(x)$, then use the identity at $x = \xi$; Equation (38) follows, with equality at $x = \pm\xi$ in the even part. **Step 4 (apply to Binomial data).** Write $\log p(y_r \mid \theta) = \log \binom{m_r}{y_r} + (y_r - m_r/2)\theta - m_r \log[2\cosh(\theta/2)]$. Apply the tangent bound to its even part and sum over observations. Exponentiating and integrating an everywhere-lower-bounded nonnegative integrand preserves the ordering. Thus $\log \widehat{A}_B^{\mathrm{JJ}} \leq \log A_B$, but the value of $\log A_B - \log \widehat{A}_B^{\mathrm{JJ}}$ has not been upper-bounded by this argument. □

7.2.3 Pólya–Gamma augmentation and mean-field computation

Proposition 7.7 (Augmentation identity and Gaussian coordinate updates)**.** *For a Binomial likelihood without its θ-independent coefficient, let $b_r = m_r$, $\kappa_r = y_r - m_r/2$ and denote the Pólya–Gamma density by $p_{\mathrm{PG}}(\omega \mid b, 0)$. Then*

$$\frac{e^{y_r\theta}}{(1 + e^{\theta})^{m_r}} = 2^{-b_r} e^{\kappa_r\theta} \int_0^{\infty} e^{-\omega\theta^2/2} p_{\mathrm{PG}}(\omega \mid b_r, 0)\, \mathrm{d}\omega. \tag{39}$$

With $\theta \sim \mathcal{N}(\nu, \rho^2)$, the mean-field family $q(\theta)\prod_r q(\omega_r)$ admits coordinate updates

$$q(\theta) = \mathcal{N}(\mu_q, V_q), \qquad V_q^{-1} = \rho^{-2} + \sum_r \mathbb{E}_q[\omega_r],$$
$$\mu_q = V_q\left(\nu/\rho^2 + \sum_r \kappa_r\right), \qquad q(\omega_r) = \mathrm{PG}(b_r, c_r), \qquad c_r^2 = \mu_q^2 + V_q, \tag{40}$$

where $\mathbb{E}_q[\omega_r] = b_r \tanh(c_r/2)/(2c_r)$ with its continuous limit at $c_r = 0$. If an additional whole-likelihood *power a_r is used, the identity requires $(b_r, \kappa_r) = (a_r m_r, a_r(y_r - m_r/2))$; multiplying an independent $\mathrm{PG}(m_r, 0)$ variable by a_r is not equivalent.*

Proof. **Step 1 (factor logistic likelihood).** The identity $1 + e^\theta = 2e^{\theta/2}\cosh(\theta/2)$ converts the Binomial likelihood into $2^{-m_r}e^{\kappa_r\theta}\cosh(\theta/2)^{-m_r}$. **Step 2 (introduce the PG mixture).** The Pólya–Gamma Laplace-transform identity is $\cosh(\theta/2)^{-b} = \int e^{-\omega\theta^2/2}p_{\mathrm{PG}}(\omega \mid b, 0)\,\mathrm{d}\omega$ for $b > 0$. Substitution proves Equation (39) (with the $b = 0$ contribution interpreted as one). **Step 3 (update the Gaussian factor).** The augmented log joint is quadratic in θ conditional on the PG variables. Taking the expectation over $q(\omega)$ and adding the Gaussian prior gives precision $\rho^{-2}+\sum_r \mathbb{E}_q[\omega_r]$ and linear coefficient $\nu/\rho^2+\sum_r \kappa_r$, proving the Gaussian update. **Step 4 (update the latent factors).** The coordinate log density for $q(\omega_r)$ is $\log p_{\mathrm{PG}}(\omega_r \mid b_r, 0) - \omega_r\mathbb{E}_q[\theta^2]/2$ plus a constant. Exponential tilting gives $\mathrm{PG}(b_r, c_r)$, $c_r^2 = \mathbb{E}_q[\theta^2]$, and the standard PG mean gives Equation (40). Each exact coordinate maximization cannot lower its *properly evaluated* ELBO; it does not certify the gap to the exact evidence. **Step 5 (external powers).** Raising the complete Binomial factor to a_r changes both the hyperbolic-cosine exponent and linear coefficient, hence the PG shape and κ_r as stated. A scaled PG random variable does not have this tilted law. □

For the intercept-only logistic–Gaussian block, the optimized Jaakkola–Jordan quadratic bound and the corresponding PG mean-field Gaussian updates can be algebraically equivalent. For the present numerical comparison, the `jj` and `pg_vb` labels correspond to the same implementation and hence represent a single computational method. A distinct Pólya–Gamma implementation would require separate numerical validation.

7.2.4 Expectation propagation and numerical reference

Expectation propagation approximates likelihood contributions with Gaussian sites and iterates cavity/tilted-distribution moment matching. A valid approximation to the *segment marginal likelihood* must include the global Gaussian normalizer and the site/tilted normalizing corrections, not merely integrate a normalized approximate posterior (whose integral is one). The EP estimates are not used to establish integrated-likelihood accuracy in the absence of a separately validated normalizer, damping scheme, convergence criterion, and reference comparison. Adaptive one-dimensional quadrature, with a documented integration tolerance and tail or endpoint control, remains the numerical reference for low-dimensional blocks.

7.3 Family-specific posterior prediction

A fitted model records the observation family, the segment posterior state, and the coordinate range used for fitting. Prediction must use the posterior predictive distribution of that observation

family. Coordinates outside the fitted range are rejected by default; an extrapolation rule is permitted only when it is stated as part of the predictive model before evaluation.

Consider a fitted segment B with posterior conjugate hyperparameters (α_B, β_B). Let $(S_B^{\mathrm{new}}, W_B^{\mathrm{new}}, H_B^{\mathrm{new}})$ denote the exposure-aware sufficient summaries of new observations assigned to B.

Proposition 7.8 (Posterior-predictive likelihood for a fixed segment)**.** *Under the exponential-family formulation of Section 4,*

$$p(y_B^{\mathrm{new}} \mid y_B) = e^{H_B^{\mathrm{new}}} \frac{Z(\alpha_B + S_B^{\mathrm{new}}, \beta_B + W_B^{\mathrm{new}})}{Z(\alpha_B, \beta_B)}. \tag{41}$$

Proof. **Step 1 (begin with the fitted segment posterior).** Conditional on the selected fitted segment, the training data update the conjugate hyperparameters from (α_0, β_0) to (α_B, β_B). This posterior, not the original segment prior, is the integrating measure for future observations. **Step 2 (aggregate new sampling information).** Under the same observation family and its declared descriptors, the new data likelihood has the kernel $\exp\{H_B^{\mathrm{new}} + \eta(\theta)^\top S_B^{\mathrm{new}} - W_B^{\mathrm{new}} A(\theta)\}$. **Step 3 (integrate once).** Multiplication by the training posterior produces the conjugate kernel with parameters $(\alpha_B + S_B^{\mathrm{new}}, \beta_B + W_B^{\mathrm{new}})$. Integration yields its normalizer divided by $Z(\alpha_B, \beta_B)$, multiplied by $e^{H_B^{\mathrm{new}}}$, giving Equation (41). **Step 4 (conditioning).** This density is posterior predictive conditional on the fitted segment and its assignment rule. It is not automatically averaged over alternative partitions, nor is it valid for a test family or coordinate support different from the declared predictive model. □

Prediction may condition on an exported joint-MAP partition or average over partitions and segment counts. The first calculation is conditional on a selected partition; the second integrates segmentation uncertainty. In either case, assignment of prediction coordinates to segments is part of the statistical model. Assigning every out-of-range coordinate to the final segment without an explicit extrapolation assumption is not a valid posterior-predictive calculation.

7.4 Joint prediction for sequences, grouped units, and multivariate responses

The fixed-segment predictive formula in Proposition 7.8 supports several input structures, but *joint* scoring must preserve dependence induced by a shared segment parameter. A product of separately integrated single-observation predictive densities is generally not the integrated density of their joint collection.

Proposition 7.9 (Joint predictive integration within a fitted segment)**.** *Condition on a fitted partition t and conjugate posterior $p(\theta_B \mid y_B)$ for each of its segments B. If new observations D_B^{new} assigned to B are conditionally independent given θ_B and have sufficient summaries $(S_B^{\mathrm{new}}, W_B^{\mathrm{new}}, H_B^{\mathrm{new}})$, then*

$$p(D_B^{\mathrm{new}} \mid y_B, t) = e^{H_B^{\mathrm{new}}} \frac{Z(\alpha_B + S_B^{\mathrm{new}}, \beta_B + W_B^{\mathrm{new}})}{Z(\alpha_B, \beta_B)}. \tag{42}$$

The full fixed-partition joint predictive density is the product of these contributions over distinct segments*, not in general over individual observations within one segment.*

Proof. **Step 1 (conditional likelihood).** Given θ_B, independence gives $p(D_B^{\text{new}} \mid \theta_B) = \prod_{r \in D_B^{\text{new}}} p(y_r \mid \theta_B, d_r)$. **Step 2 (aggregate once).** The sufficient-statistic representation collects all new observations into one exponential-family kernel with summaries $S_B^{\text{new}}, W_B^{\text{new}}, H_B^{\text{new}}$. **Step 3 (integrate once).** Multiply this kernel by the training posterior and integrate θ_B, yielding Equation (42). It differs from integrating θ_B separately for each observation, which would spuriously treat new observations as having independent latent parameters. **Step 4 (between-segment factorization).** Under the model assumption of conditionally independent segment parameters, the posterior factorizes across distinct segments given t. Therefore the joint predictive density factors over the distinct segments, completing the claim. □

Corollary 7.10 (Closed-form predictive distributions in three conjugate families)**.** *Conditional on a fitted segment and its conjugate posterior, the following are ordinary posterior-predictive sampling distributions:*

1. *For Gaussian $\mu \mid y_B \sim \mathcal{N}(\nu_B, V_B)$ and new observations with known variances σ^2/w_r, a vector of m new observations in the same segment has joint law*

$$Y^{\text{new}} \mid y_B, t \sim \mathcal{N}_m\Big(\nu_B \mathbf{1}_m,\ V_B \mathbf{1}_m \mathbf{1}_m^\top + \operatorname{diag}(\sigma^2/w_1, \ldots, \sigma^2/w_m)\Big)\,.$$

2. *For Poisson $\lambda \mid y_B \sim \operatorname{Gamma}(a_B, b_B)$ and a new count at exposure $e > 0$, the predictive law is negative Binomial with*

$$p(y^{\text{new}} = z \mid y_B, e) = \frac{\Gamma(a_B + z)}{\Gamma(a_B) z!} \left(\frac{b_B}{b_B + e}\right)^{a_B} \left(\frac{e}{b_B + e}\right)^{z}.$$

3. *For Binomial $p \mid y_B \sim \operatorname{Beta}(a_B, b_B)$ and m future trials, the predictive count law is Beta–Binomial with*

$$p(y^{\text{new}} = z \mid y_B, m) = \binom{m}{z} \frac{B(a_B + z, b_B + m - z)}{B(a_B, b_B)}.$$

Proof. **Step 1 (Gaussian joint law).** Write each new observation as $Y_r = \mu + \epsilon_r$ with independent $\epsilon_r \sim \mathcal{N}(0, \sigma^2/w_r)$ conditional on μ, and with $\mu \sim \mathcal{N}(\nu_B, V_B)$ from the training posterior. Gaussian addition gives mean $\nu_B \mathbf{1}_m$ and covariance $V_B \mathbf{1}_m \mathbf{1}_m^\top + \operatorname{diag}(\sigma^2/w_r)$. The rank-one term shows why multiplying separate Gaussian marginals would be incorrect for joint scoring. **Step 2 (Poisson–Gamma mixture).** Integrate $e^{-e\lambda}(e\lambda)^z/z!$ against the Gamma posterior density. The Gamma integral has shape $a_B + z$ and rate $b_B + e$, which simplifies to the displayed negative-Binomial probability mass function. **Step 3 (Beta–Binomial mixture).** Integrate $\binom{m}{z} p^z (1-p)^{m-z}$ against $\operatorname{Beta}(a_B, b_B)$. Combining exponents produces $B(a_B + z, b_B + m - z)/B(a_B, b_B)$, establishing the third expression. □

Pointwise sequences (Case A). New pointwise observations carry coordinates, outcomes, and the family-specific descriptors. Under an exported partition, coordinates select segments and the data are aggregated *within* those segments before applying Equation (42). An out-of-range coordinate requires a declared extrapolation model; a coordinate gap is not an exposure by default.

Set-valued units (Case B). A unit may contain multiple pairs (x_{ur}, y_{ur}) on a physical support interval. Its segment-specific internal observations can be scored jointly using the same sufficient-statistic aggregation. If several units in a single dataset share segment parameters, the dataset

score aggregates *across all such units before integrating*. Multiplying separately marginalized unit likelihoods is justified only under an additional model in which the units have independent latent segment parameters; it is otherwise a composite score.

Vector-valued responses (Case C). When response coordinates are conditionally independent given distinct component-specific segment parameters, the joint segment evidence and predictive density factor across response dimensions. For a genuinely multivariate exponential family with shared or correlated parameters, the joint sufficient statistics and conjugate normalizer must be used instead. The global ordered-partition recursion depends only on the resulting joint segment score.

Group membership and segmentation averaging. If M_g defines a *normalized* predictive data density and π_g is a prior group probability, then $p(g \mid D^{\text{new}}) \propto \pi_g p(D^{\text{new}} \mid M_g)$ is a Bayesian group posterior. Scores from the finite latent-group MM criterion in Equation (29), which are not asserted to be normalized data densities, do not automatically admit this interpretation. Likewise, true segmentation-averaged prediction sums joint predictive contributions over $p(k, t \mid y)$; prediction conditional on a chosen joint-MAP partition is a different object.

Calibration. For discrete outcomes, randomize the probability integral transform within the probability mass at the observed outcome. For continuous outcomes, PIT values are marginally uniform under a calibrated predictive law, but they are generally dependent within a sequence. Report sequence-level calibration or simulation-based reference distributions rather than applying an independent-uniform goodness-of-fit reference blindly. Held-out predictive scores need not rank candidate k in the same order as training posterior count probabilities.

7.5 Boundary metrics

Definition 7.11 (One-to-one tolerance matching)**.** For estimated boundaries $\widehat{\mathcal{B}}$, reference boundaries $\mathcal{B}^\star$, and tolerance τ, form the bipartite graph containing an edge $(\widehat{b}, b^\star)$ whenever $|\widehat{b} - b^\star| \le \tau$. Choose a matching of maximum cardinality; among maximum-cardinality matchings, choose one of minimum total absolute distance under a deterministic tie rule. Precision, recall, and F_1 use the resulting number of matches. No boundary is used more than once.

Definition 7.12 (Matched location error)**.** For the matching $\mathcal{M}_\tau$,

$$\mathrm{MAE}_\tau = |\mathcal{M}_\tau|^{-1} \sum_{(\widehat{b}, b^\star) \in \mathcal{M}_\tau} |\widehat{b} - b^\star|$$

when $|\mathcal{M}_\tau| > 0$. When no pair is matched, the value is reported as NA, not as zero.

Definition 7.13 (Admissible held-out predictive score)**.** A held-out log score is used as predictive evidence only when the held-out observations were excluded from fitting, the posterior predictive distribution of the fitted observation family was used, and every prediction coordinate lay in the fitted support or followed a pre-specified extrapolation model. A violation of any condition invalidates the number for that predictive interpretation without changing the fact that the computation was executed.

Predictive model checking. A posterior-predictive score is meaningful only for the observation law that was fitted and for new coordinates within its stated support. An earlier methylation calculation used a Gaussian predictive density for a fitted Beta-response model and extrapolated outside the fitted range. That value is not included in the predictive results. The admissible Beta-response held-out analysis is reported in Section 10.

Interpretation of approximate local scores. The forward–backward and max–sum recursions are exact for the block-score array supplied to them. When the array is obtained by numerical approximation, this algorithmic exactness does not establish exactness with respect to the underlying sampling model. A Jaakkola–Jordan tangent gap averaged under a fitted Gaussian variational distribution is not an upper bound on the log-evidence error: the difference in log normalizing constants also depends on the changed posterior measure. Likewise, a truncated Laplace expansion is not a finite-sample error certificate unless its omitted remainder and tails have been bounded. Comparisons with a converged numerical-integration reference are necessary before interpreting either diagnostic as evidence of approximation accuracy.

Interpretation of nonconjugate method labels. In the implementation studied here, the options `jj` and `pg_vb` both invoke the same Jaakkola–Jordan numerical routine. These labels therefore do not constitute two independent approximation methods. A distinct Pólya–Gamma variational implementation, if included later, requires a separately specified augmented model, objective, update equations, and numerical verification.

8 Algorithms, computational complexity, and numerical implementation

The implementation separates calculations specific to an observation model from calculations over ordered partitions. Observation-model routines compute sufficient statistics, segment marginal likelihoods, posterior moments, numerical error information, and posterior-predictive quantities. The partition routines combine prior-adjusted segment scores, evaluate sum-product recursions, recover joint MAP partitions by max-sum recursion and backtracking, pool aligned sequences, and extract posterior summaries.

8.1 Inputs and returned quantities

The fit input separates four objects that have different statistical meanings: ordered coordinate u_r, response y_r, observation descriptor d_r, and optional likelihood power a_r. Descriptors such as exposure or trial count are interpreted by the corresponding observation model. Irregular geometry enters the prior through u and the factors c_u, h_u; it is not treated as exposure in the observation likelihood.

8.2 Log-space evaluation and failure handling

All segment and partition scores are stored in log space. Sums in Equations (16)–(17) are evaluated by a stable `logsumexp`. Cumulative sufficient statistics should be centered or scaled when appropriate, and conjugate hyperparameters must lie in their proper domains. When a numerical segment calculation fails to converge, violates support, or lacks the stated error control,

Algorithm 1: BayesBreak offline fit

Input: Ordered observations $\{(u_r, y_r, d_r, a_r)\}_{r=1}^{n}$; observation family and hyperparameters; $k_{\max}$; segment-count prior $p(k)$; segment-cohesion and boundary-hazard factors c_u, h_u; numerical tolerances.

Output: Posterior distribution over k; boundary-event and ordered-boundary marginals; joint MAP partitions; segment posterior states; Bayes curves; numerical diagnostics and result metadata.

Validate order, observation-family support, descriptors, power weights, and candidate-segment restrictions;
Construct cumulative sufficient statistics when available;
for *each admissible segment* $(i, j]$ **do**
    compute $\log A_{ij}^{(0)}$ and the required moments or posterior state;
    record validity, convergence, and approximation diagnostics;
    assign score $-\infty$ to an inadmissible segment rather than substituting a finite value from another model;
Compute $\log \gamma_u(i, j) = \log c_u(i, j) + \mathbb{1}\{j < n\} \log h_u(j)$ once for each segment;
Run log-space forward and backward recursions and normalize Equation (20);
Compute ordered-boundary marginals, boundary-event probabilities, and Bayes-curve moments;
Run max-sum recursion and deterministic backtracking for every retained k or for the selected $\widehat{k}$;
Check forward–backward equality, posterior normalization, boundary-event mass, and backtrack-score equality;
Export the observation-family identifier, fitted coordinate range, result metadata, and numerical diagnostics;

Algorithm 2: Finite latent-group score-clustering with exact template updates

Input: Per-subject log segment likelihoods $\ell_{ij}^{(s)}$; local partition factors $\gamma_u(i, j)$; count prior $p(k)$ and normalizers C_k; group count G; restarts and stopping tolerance.

Initialize G valid template partitions and mixture weights for each restart;
repeat
    Compute each template score $\log S_s(\tau_g) = \log p(k_g) - \log C_{k_g} + \sum_{(i,j]\in\tau_g}(\ell_{ij}^{(s)} + \log \gamma_u(i, j))$;
    Compute auxiliary weights r_{sg} by stabilized softmax over $\log \pi_g + \log S_s(\tau_g)$;
    Update $\pi_g \leftarrow S^{-1} \sum_s r_{sg}$, handling empty groups under the declared policy;
    For every g, form $\ell_{ij}^{(g)} = \sum_s r_{sg} \ell_{ij}^{(s)} + n_g \log \gamma_u(i, j)$;
    Run max-sum for every feasible k and compare terminal scores with the offset $n_g[\log p(k) - \log C_k]$;
    Backtrack to obtain the updated template τ_g;
    Recompute and record $\mathcal{F}(\pi, \tau)$ *after* all template and weight updates; check nondecrease;
until *the fitted templates are stable and the* recomputed final *objective change is below tolerance*;
Rank restarts by the actually recomputed final $\mathcal{F}$, not by a pre-update trace value;

Output: Fitted templates, group weights, row-stochastic auxiliary weights, and full objective traces.

Table 3. Principal returned quantities and their inferential interpretation.

Quantity	Calculation	Interpretation	Numerical checks
$\log A_{ij}^{(0)}$	observation-model segment calculation	log integrated likelihood or declared numerical approximation	support, finiteness, convergence
$p(k \mid y)$	terminal sum-product quantities	posterior uncertainty in segment count	normalization, prior normalizer
$p(t_p = h \mid y, k)$	forward–backward factorization	marginal distribution of ordered boundary p	normalization over h
$p(b_h = 1 \mid y, k)$	sum over ordered boundaries	probability of any changepoint at h	total mass $k - 1$
$\widehat{t}_k^{\mathrm{MAP}}$	max-sum recursion and backtracking	joint MAP partition for fixed k	terminal-score equality
$\mathbb{E}[m_\star(\theta_s)^r \mid y]$	sum over segments covering s	model-averaged segment functional	common marginal-likelihood denominator
Predictive score	observation-family posterior predictive	out-of-sample log score under the stated support rule	holdout, family dispatch, coordinates

it returns a structured failure state. The analysis then follows a declared rule: exclude the segment by assigning $-\infty$, terminate the fit, or continue while labeling the posterior as based on an unverified approximation. It must not silently substitute a different observation model.

Tie rules are deterministic and recorded. In max-sum recursion, the predecessor rule identifies which member of a set of equally scoring partitions is returned. In one-to-one boundary matching, the minimum-distance refinement and a fixed tie order identify the matching. These conventions make repeated calculations reproducible; they do not alter the statistical estimand.

8.3 Computational complexity

Table 4 separates asymptotic operation counts from finite-range timing measurements.

Enumeration of all admissible segments is the principal cost of the exact calculation. Pruning, screening, or a restricted changepoint grid can reduce computation, but may also remove partitions unless the restriction is proved exact for the specified model. Such accelerations are not included in the exactness claims of this paper unless their preservation of the partition sum is established separately.

8.4 Reproducibility

A reproducible analysis specifies the observation family, hyperparameters, candidate-boundary restrictions, prior on segment count, coordinate construction, and numerical tolerances. For stochastic initialization or simulation, the random seed and the rule for generating replicate seeds must also be specified. The experimental protocols in Section 9 identify the estimands and analysis units; the corresponding executable configurations and data provenance accompany the source code.

Table 4. Computational complexity of the implemented calculations. Empirical timings are reported separately in Section 10.

Component	Time	Working memory	Numerical consideration
Conjugate segment precomputation	$\Theta(n^2)$	$\Theta(n^2)$	cumulative sufficient statistics; invalid segments receive $-\infty$
Nonconjugate segment precomputation	$\Theta(n^2 T_{\text{seg}})$	$\Theta(n^2)$ plus method state	convergence and error diagnostics
Sum-product recursion	$\Theta(k_{\max} n^2)$	$\Theta(k_{\max} n)$ for all message layers	log-sum-exp evaluation
Boundary and moment extraction	$\Theta(k_{\max} n^2)$ for streamed fixed-k summaries	output dependent; $\Theta(k_{\max}^2 n)$ for every ordered-boundary table	common marginal-likelihood normalizer
Max-sum recursion and backtracking	$\Theta(k_{\max} n^2)$	$\Theta(k_{\max} n)$ including backpointers	deterministic predecessor rule
Shared-changepoint pooling	$\Theta(Sn^2 + k_{\max} n^2)$ after sequence-specific segment calculations	sequence arrays or streamed log-score sum	add log marginal likelihoods across sequences
Latent-group template update	$\Theta(GSn^2 + Gk_{\max} n^2)$ per iteration	problem dependent	restart ranking must use the final objective value

9 Experimental protocols

We distinguish completed analyses, reported in Section 10, from additional experiments required to evaluate the statistical properties of the method. The designs below specify the data-generating mechanisms, sample sizes, comparators, estimands, replication, and primary analyses *before* results are generated. Every figure and table labeled **Planned** is an unfilled result location, not an observation or a prediction of performance. The complete configuration, data-source, metric, and reporting specifications are maintained with the implementation in the accompanying reproducibility protocol; the scientific definitions and primary comparisons are given here.

9.1 Common evaluation conventions

All simulations use independent datasets as the unit of replication. A configuration is fixed before the generating seed is drawn; all methods in a configuration receive identical simulated observations and candidate boundaries. The default generation scheme uses NumPy `SeedSequence` with separate components for experiment, design cell, and replicate, so new cells do not change existing seeds. The primary comparison uses 100 independently generated datasets per cell unless otherwise specified below. A five-dataset pilot per cell checks feasibility and memory use; pilot results are not included in reported performance estimates. Tests of the exact recursions use separate non-random exhaustive enumeration and numerical-integration oracles.

For boundary evaluation, use a maximum-cardinality one-to-one matching of estimated to generating boundaries within tolerance τ. Among matchings with maximal cardinality, minimize total displacement. The primary tolerance is $\tau = 3$ indices; report $\tau \in \{0, 1, 5\}$ as sensitivity analyses. Compute precision, recall, and F_1 from the matched pairs, counting unmatched estimated and

true boundaries in the usual manner. Matched-location error is not defined when no boundaries match. The segment-count metrics are $\Pr(\widehat{k} = k^\star)$, $|\widehat{k} - k^\star|$, and the probability of $k = k^\star$ under the count posterior. For signal estimation report mean squared error relative to the *generating segment functional*, not across families with incompatible response scales.

For posterior probabilities $\widehat{p}_{ri} = p(b_i = 1 \mid y^{(r)})$, report Brier score, logarithmic score using a prespecified numerical clipping rule, and reliability curves with counts per bin. Use ten equal-mass bins as the primary diagnostic, separately reporting event prevalence; show equal-width bins as a sensitivity analysis. Confidence intervals for simulation metrics use the independent sequence as the resampling unit (2,000 stratified bootstrap replicates with the resampling seed declared in the protocol). Dependent positions from one sequence are not treated as independent replicates. Intervals are descriptive across generated datasets, not frequentist coverage statements about an individual posterior interval.

For any algorithm fitted to held-out data, hyperparameters and comparator penalties are selected on disjoint training/validation sequences or by a predeclared training-only rule. Methods are compared at matched input, coordinate axis, and observation law. An agreement score against the BayesBreak MAP is never labeled ground-truth accuracy. Unless the models define proper probabilities for the same observed variables and base measure, their marginal-likelihood values are not directly compared.

9.2 E00: exactness checks

For $n \in \{2, \ldots, 10\}$, enumerate every admissible partition at every segment count. Test uniform and nonuniform normalized priors, including zero prior factors and minimum-segment constraints. Compare $L_{k,n}$, $R_{k,0}$, C_k, $p(k \mid y)$, ordered and event boundary marginals, posterior moments, and conditional and global MAP decisions against direct enumeration. Include a Gaussian example with negative segment posterior means to test signed moment aggregation. Independently compare the Gaussian, Poisson, Binomial, and negative-Binomial block integrals to one-dimensional numerical integration for unit and nonunit exposure/trial/precision descriptors. Structural infeasibility is not conflated with a block whose observed-data likelihood equals one.

9.3 E01 and E02: simulated recovery, calibration, and family generality

E01 uses regularly spaced Gaussian sequences of lengths 256 and 512, with $k^\star \in \{1, 3, 6\}$, minimum generating segment length 32, and adjacent mean differences $\Delta/\sigma \in \{0.5, 1, 2\}$. For $k^\star = 1$, the jump-size parameter is immaterial and duplicate null cells are omitted. Use the fixed proper segment prior $\mu \sim N(0, 4)$ and known observation variance $\sigma^2 = 1$; these quantities are held constant for the confirmatory runs. Analyze both the model-averaged boundary-event posterior and the posterior conditional on the true segment count as separately named outputs. Primary metrics are boundary $F_1@3$, count posterior coverage, Brier score, and calibration by boundary probability. Do not pool boundary positions as independent bootstrap units.

E02 fixes three generating segments and $n = 240$ for conjugate families; an initial $n = 120$ setting is used for Beta-valued quadrature because it requires numerical integration for every candidate segment. The exact generating parameters are specified as rates, probabilities, dispersion and observation precisions, with two effect-size settings per family. Report each family on its own natural signal scale and include a standardized estimation metric for display across panels. Use 100 simulated datasets per conjugate cell and 50 per quadrature cell, after feasibility checks. The

Table 5. Planned validation matrix. Entries specify experiments to run; they are not additional results. Exact parameter grids and artifact destinations are included in the companion experimental protocol.

ID	Question	Main design and primary estimand	Intended output
E00	Exactness of partition inference	Exhaustive partitions for $n \leq 10$ and integrated-block quadrature	Oracle agreement table
E01	Posterior calibration and recovery	Gaussian $n \in \{256, 512\}$, $k^\star \in \{1, 3, 6\}$, varying jump size	Reliability curves and recovery table
E02	Observation-model generality	Gaussian, Poisson, Binomial, negative Binomial, and Beta-valued responses	Family recovery table
E03	Irregular-design prior	Uniform, gap-hazard, length-cohesion, combined priors; controlled gaps	Boundary mass and detection plots
E04	Common-changepoint pooling	$S \in \{1, 4, 16, 64\}$ sequences, with and without shared truth	Recovery and calibration table
E05	Latent-template estimation	$G \in \{1, 2, 3\}$; separation/noise sweeps; multiple restarts	ARI and collapse plots
E06	Nonconjugate segment integration	Logistic-normal blocks, adaptive quadrature reference	Error versus block length, posterior discrepancy
E07	Misspecification and missingness	Heavy tails, overdispersion, absent outcomes	Coverage and robustness plots
E08	Posterior prediction	Disjoint held-out contiguous regions; correct family scoring	Test log score and uncertainty table
E09	External comparators and case studies	Matched coordinate/likelihood analyses and annotations when verified	Independent benchmark tables
E10	Scaling and resource cost	$n \in \{128, 256, 512, 1024, 2048\}$ and $k_{\max} \in \{5, 10, 20\}$	Time and memory plots

Table 6. Planned Table P0 (E00). Independent verification of the exact recursions and family-specific integrals. Fill each entry with the maximum observed numerical discrepancy across the enumerated test cases.

Quantity	Cases tested	Maximum absolute error	Acceptance tolerance
Partition sums and normalizers	TBD	TBD	10^{-8} on log scale
Boundary probabilities and moments	TBD	TBD	10^{-9} probability
Family-specific integrated likelihoods	TBD	TBD	Reference integration tolerance

comparison tests the common inference algorithm, not whether one observation family is superior to another.



Figure 4. Planned Figure P1 (E01). Posterior calibration and detection versus signal-to-noise ratio. Show independent-dataset uncertainty bands and per-bin sample counts in the final figure. No numerical result is asserted here.

Table 7. Planned Table P1 (E01–E02). Predeclared reporting columns. One row per experiment, design cell, and method will be generated by the simulation registry. The values below have not been measured.

Design and method	N datasets	F_1@3 [95% CI]	$\Pr(\widehat{k} = k^\star)$	Brier / log score
Gaussian, varying SNR	TBD	TBD	TBD	TBD
Poisson / Binomial / NegBin	TBD	TBD	TBD	TBD
Beta-valued quadrature	TBD	TBD	TBD	TBD

9.4 E03: irregularly spaced observations

Simulate $n = 240$ design points with three piecewise-constant segments and an inserted empty interval between two clusters. Vary the relative gap width over $\{1, 4, 16\}$ times the within-cluster spacing and the changepoint location relative to the large gap. Analyze the same data under (i) index-uniform partitions, (ii) a candidate-boundary hazard proportional to the physical gap, (iii) segment-length cohesion, and (iv) their product. Normalize each conditional partition prior with its own C_k and report posterior boundary mass at and near the gap as well as one-to-one recovery. Report a separate simulation in which the generating boundary is *not* at the large gap so that the prior's false-positive behavior can be measured. This comparison is a prior-sensitivity analysis, not a guarantee that any gap prior is uniformly preferable.




Figure 5. Planned Figure P6 (E03). Comparison of index-uniform, candidate-gap, length-cohesion, and combined priors under both generating boundary placements. Show uncertainty across independent simulated datasets.

9.5 E04 and E05: hierarchical segmentation

For E04, generate $S \in \{1, 4, 16, 64\}$ independent Gaussian sequences conditionally on a common three-segment template, then replace 25% and 50% of the sequences with shifted subject-specific boundaries in separate misspecification cells. Compare the common-boundary posterior, separate fits, and a suitable partial-sharing baseline on boundary $F_1@3$, calibration, and signal error. Sequence-level bootstrap resampling respects the shared generating configuration.

For E05, use $G \in \{1, 2, 3\}$ templates and $S \in \{24, 48\}$ sequences, with predeclared low, medium, and high separation cells. Run five initializations per dataset and select using the *recomputed final* finite-template objective. Report label-invariant clustering accuracy and adjusted Rand index, complete objective traces, the fraction of empty or duplicate templates, and failures under reduced separation. The criterion and its MM algorithm are those of Section 6; do not replace the criterion with a normalized generative mixture during this experiment.




Figure 6. Planned Figure P2 (E04–E05). Effects of replication, assumption violations, and between-template separation. Plot dataset-level uncertainty and template-collapse frequency.

9.6 E06: nonconjugate numerical evidence

Use intercept-only Binomial–logistic segments with normal prior on the logit, block sizes $\{4, 8, 16, 32, 64, 128\}$, and generating success probabilities $\{0.05, 0.5, 0.95\}$. For each cell generate 100 independent blocks. Establish a converged adaptive quadrature reference with absolute and relative tolerances 10^{-11} and 10^{-10}, checking agreement with a second numerical method on a stress subset. Compare the segment log marginal likelihood and its first two posterior signal moments. Include Laplace, the implemented Jaakkola–Jordan bound, and expectation propagation only if its complete normalizing constant and stopping rule pass validation. The current `pg_vb` option calls the JJ routine and is not counted as an independent method. Propagate these same block-score arrays through a segmentation simulation, reporting posterior count discrepancy, boundary probability error, ranking changes, and runtime. The maximum over reachable segments must be used when interpreting the uniform-error theorem; a posterior-weighted error is a distinct diagnostic.

9.7 E07 and E08: robustness and held-out prediction

E07 repeats the $n = 256$, three-segment recovery experiment with Student t_3 noise fitted by a Gaussian block model, negative-Binomial counts fitted by Poisson blocks, and overdispersed success counts fitted by Binomial blocks. Separately remove 10% or 30% of responses completely at random, or a single contiguous 15% region. Missing outcomes are represented by the observation mask, not a zero Gaussian precision or a zero Poisson exposure. Compare boundary calibration, count selection, and frequentist recovery to the correctly specified reference.




Figure 7. Planned Figure P3 (E06). Numerical integration error and its propagation to segmentation posteriors, with failure rates and runtime reported separately. This figure is not a numerical certificate.

Table 8. Planned Table P2 (E06). Quadrature-validated numerical approximation study. Report failed or nonconvergent calls in the denominator.

Method	Median $\lvert\Delta \log A\rvert$	Maximum $\lvert\Delta \log A\rvert$	Failure fraction	Time [95% CI]
Adaptive quadrature (reference)	TBD	TBD	TBD	TBD
Laplace	TBD	TBD	TBD	TBD
Jaakkola–Jordan	TBD	TBD	TBD	TBD
EP, conditional on validation	TBD	TBD	TBD	TBD

E08 holds out contiguous regions from synthetic and real sequences while retaining the fitted coordinate endpoints. The split is made before any hyperparameter estimation. Each fitted observation family uses its own posterior-predictive distribution with the *observed* new-data exposure, trial count, or precision. Report log predictive density per held-out observation and proper prediction intervals or randomized PIT diagnostics as appropriate for the family. Distinguish prediction conditional on the estimated partition from averaging over the segmentation posterior. A pointwise composite score is never called a joint predictive density.

Table 9. Planned Table P3 (E07–E08). Robustness and predictive performance. Comparisons will be paired within identical held-out splits.

Perturbation or model	$F_1@3$ [95% CI]	Held-out log density / obs.	Prediction coverage
Correctly specified reference	TBD	TBD	TBD
Heavy tails / overdispersion	TBD	TBD	TBD
Missingness or blocked holdout	TBD	TBD	TBD

9.8 E09: external comparisons and real-data sensitivity

Use the original well-log record, the probe-aligned array-CGH matrix, the 2015–2023 daily S&P 500 series, and the `methylKit` chromosome-21 example with dataset identity and checksums fixed before fitting. Reproduce the reported preprocessing and include the full well-log record alongside the subsample. Array-CGH must first use a validated Gaussian precision likelihood; report its corrected results separately from the earlier powered-weight fit. For methylation, compare count-

based Binomial or Beta-Binomial modeling when raw methylated and total reads are available to the existing continuous-Beta response model; do not regard a Gaussian predictive fallback as a valid Beta score. Vary $k_{\max}$ until posterior mass near the upper truncation is small, or explicitly report truncation sensitivity if that criterion is not met. Evaluate a grid of segment-count priors and physical-coordinate priors with the normalizer recomputed for every model.

Compare with PELT, optimal partitioning, binary segmentation, wild binary segmentation, and a Bayesian reference implementation when each comparator is appropriate to the response and feasible at the target length. Distinguish (a) reference-annotation accuracy when independent labels and alignment are actually verified, (b) model-to-model agreement when only fitted boundaries are available, and (c) held-out predictive comparisons on the same outcomes. No figure caption or table may relabel model agreement as biological or geological ground truth. Baseline penalties, feature transforms, and stopping criteria are determined by disjoint tuning data or stated a priori.



Figure 8. Planned Figure P4 (E09). Fair matched-axis comparisons and posterior sensitivity to $k_{\max}$, distinguished by whether an independent boundary reference is available.

Table 10. Planned Table P4 (E09). Corrected real-data and comparator study; every row will identify its observed variables, reference, and priors.

Dataset / model	$\widehat{k}$, tail mass	Reference type	F_1@3 [95% CI]	Test log density
Well-log, full and subsampled	TBD	TBD	TBD	TBD
Array-CGH, corrected precision	TBD	TBD	TBD	TBD
S&P 500 and methylation	TBD	TBD	TBD	TBD

9.9 E10: time and memory complexity

Benchmark $n \in \{128, 256, 512, 1024, 2048\}$ and $k_{\max} \in \{5, 10, 20\}$ on a single fixed machine, using three warmup runs and seven measured runs per configuration. Pin Python, NumPy, BLAS thread counts and data types. Record construction of the segment-score array, forward and backward recursions, MAP backtracking, posterior marginal extraction, peak resident memory, and the total wall time separately. Include the ordinary $\Theta(n^2)$ storage path and a documented streaming variant if it is implemented. Report observed finite-range slopes and theoretical worst-case complexity as different quantities.




Figure 9. Planned Figure P5 (E10). Component timing and peak memory with run-to-run dispersion; proposed guide curves represent theoretical complexity and are not fitted performance claims.

9.10 Criteria for replacing planned outputs

A planned slot is populated only when the generating configuration, data or simulation seed, code revision, software dependencies, method configuration, metric definition, and output files are available together. A table that has only some completed rows remains labeled planned until all reported entries can be traced to executed measurements. The statistical results and their limitations must be revised from those measurements, not from expected outcomes. The retained findings in Section 10 are historical experiments with stated conditions; they are not the unfilled cells in the planned validation matrix.

10 Results

The following results describe completed numerical calculations for the model configurations specified in this section. They are analyzed according to their data-generating mechanisms, observation models, and reference sets. Proposed additional evaluations remain unfilled in Section 9; their outcomes are not inferred from the present experiments.

10.1 Single-sequence posterior summaries

Figure 10 shows a Gaussian sequence with two true changepoints. The boundary-event posterior concentrates near the generating changes while retaining a smaller shoulder of ambiguity before the first jump. The joint-MAP signal reflects this local uncertainty by inserting a short intermediate segment. Table 11 shows that the selected and modal segment count is four while the posterior mean is 4.742, illustrating why an exported partition and posterior uncertainty over k should not be collapsed into one object.

Quantity	Value
Selected segment count $\widehat{k}$	4
Posterior mean $\mathbb{E}[k]$	4.742
MAP k	4
$\log p(y)$	-18.031

Table 11. Posterior summary for Figure 10.




Figure 10. Gaussian simulation. Top: marginal boundary-event posterior with generating changepoints shown by dashed lines. Bottom: observations, latent signal, joint-MAP segmentation, and 90% segment-level posterior intervals.

10.2 Portability across local observation models

The same global inference code was applied to Gaussian, Poisson, Binomial, and Beta-valued responses (Figure 11). Table 12 reports the corresponding synthetic metrics. Gaussian, Binomial, and Beta-valued settings selected the generating three segments; Poisson selected four and had the weakest boundary score and largest reconstruction error in this small benchmark. The result supports interface portability and exposes relative difficulty in the bundled settings; it is not a population-level ranking of families.




Figure 11. Observation-family comparison. Gaussian, Poisson, and Binomial use closed-form local conjugate integration; the Beta-valued response uses deterministic one-dimensional quadrature.


Family	n	$k^\star$	$\hat{k}$	F1@τ	MAE	MSE	$-\log p(y)/n$
Gaussian	120	3	3	0.968	0.00	0.0023	0.191
Poisson	120	3	4	0.777	0.20	0.4501	2.162
Binomial	120	3	3	0.960	0.06	0.0004	2.108
Beta-valued	120	3	3	1.000	0.00	0.0001	2.620

Table 12. Synthetic summary across local observation models. The Beta-valued row is quadrature-based rather than closed-form conjugate.

10.3 Calibration and latent-group segmentation

Repeated Gaussian simulations produced a reliability curve close to the diagonal, with ECE approximately 0.010 and Brier score approximately 0.011 (Figure 12a). These values support calibration in the generating design, not robustness to arbitrary misspecification.

The latent-group experiment used 24 sequences of length 80 from two known synthetic groups at noise standard deviation 1.0. It achieved 96% hard assignment accuracy, with one misassignment and visible ambiguity around the class boundary (Figure 12b). Group-specific boundary marginals concentrate near the generating templates. The result is evidence for the finite score-clustering objective in Equation (29); it is not evidence for normalized-mixture identifiability.



(a) Boundary-event reliability.

(b) Latent-group assignments and group-specific boundary marginals.

Figure 12. Calibration and latent-template analysis. Panel (a) reports ECE ≈ 0.010 and Brier ≈ 0.011 in the Gaussian calibration experiment. Panel (b) reports 96% hard recovery for the declared two-template synthetic setting.

A separate stress study evaluated 400 seeded datasets across eight predeclared cells. In the reference simulation cell, mean label-invariant hard accuracy was 0.9742 (95% interval 0.9536–0.9948) and mean adjusted Rand index was 0.9183. All 400 objective traces were monotone, every returned final objective equaled the final trace value, and all 1,200 restarts were valid. Recovery degraded under low separation, duplicate templates, higher noise, and shorter sequences, as summarized in Figure 13. These results support the finite score-clustering criterion in the declared simulations, not normalized-mixture identifiability.




Figure 13. Latent-group stress study. Top: label-invariant hard accuracy with dataset-level 95% intervals. Bottom: collapsed or duplicate-template rate. The dashed 0.5 line is a descriptive reference, not a universal recovery threshold.

10.4 Nonconjugate block approximations

Table 13 reports five method labels in the Bernoulli/logistic example, with quadrature as the local reference. The `jj` and `pg_vb` labels currently share one implementation path and are not independent computational methods. Laplace was the fastest listed routine at 0.049 seconds, whereas quadrature required 0.093 seconds. Maximum absolute block-score discrepancies were 1.235 for Laplace, 1.265 for Jaakkola–Jordan and Pólya–Gamma mean field, and 14.502 for expectation propagation. Boundary F_1 did not vary monotonically with block-score discrepancy: quadrature and Laplace selected six segments with $F_1 = 0.286$, whereas the other listed labels reported $F_1 = 0.500$ under different selected segment counts. The repeated variational row must not be counted twice in method comparisons.

Method	$\max \lvert\Delta \log A^0\rvert$	$q_{95} \lvert\Delta \log A^0\rvert$	median $\lvert\Delta \log A^0\rvert$	time (s)	F1@τ	$\hat{k}$
quadrature	0.000	0.000	0.000	0.093	0.286	6
laplace	1.235	1.188	0.980	0.049	0.286	6
jj	1.265	1.216	1.002	0.071	0.500	3
ep	14.502	14.141	12.680	0.059	0.500	7
pg-vb	1.265	1.216	1.002	0.072	0.500	3

Table 13. Local approximation comparison. The observed maximum errors are too large to treat this example as verification of a small uniform segmentwise error bound.

This result demonstrates why segmentation output alone cannot certify a block approximation. The global dynamic program aggregates many local score differences, and a boundary metric

can improve despite a worse local approximation. Claim-supporting use of Theorem 7.2 therefore requires a uniform error bound over all admissible segments rather than retrospective agreement in the estimated segmentation.

10.5 Finite-range runtime behavior

Figure 14 reports length and segment-budget sweeps. At $k_{\max} = 20$, mean runtime rose from 0.0455 seconds at $n = 50$ to 1.0585 seconds at $n = 800$ (Table 14). Fitted log–log slopes were approximately 1.07 in n for $k_{\max} = 10$, 1.14 for $k_{\max} = 20$, and 0.88 in $k_{\max}$ at fixed $n = 200$. They summarize this implementation over the measured range; they do not replace the $\Theta(k_{\max} n^2)$ worst-case accounting in Proposition 5.6.




Figure 14. Gaussian runtime measurements. Points are means over five repetitions and error bars are one standard deviation. Dashed references show quadratic-in-n and linear-in-$k_{\max}$ guides, not fitted complexity claims.

n	$k_{\max}$	mean (s)	std (s)
50	20	0.0455	0.0008
100	20	0.1033	0.0011
200	20	0.2195	0.0011
400	20	0.5106	0.0307
800	20	1.0585	0.0345

Table 14. Runtime measurements for $k_{\max} = 20$ in the length sweep.

10.6 Applications to four datasets

Table 15 summarizes the executed case studies. Each fit passed the four stated dynamic-programming checks. Their figures provide fitted segmentation, boundary uncertainty, segment-count posterior information, and in-sample diagnostics. The executed run bundles did not include independently scored reference boundaries, so these case studies establish execution and descriptive posterior behavior rather than externally validated accuracy. This does not imply that external annotations are unavailable.

Table 15. Real-data segmentations. The log evidences belong to different data models and are not compared across rows.

Case	n or shape	$\widehat{k}$	Log evidence
Well-log	507	23	-4989.2835
Array-CGH	43×2215	15	76359.7996
S&P 500	566	29	-1296.6537
Methylation	1904	15	-9518.6675

Well-log. The available source contains 4,050 NMR response values and an executed downsampled sequence of $n = 507$. Under the index-uniform prior, the Gaussian fit selected 23 segments with log evidence -4989.2835 (Figure 15). A separately executed length-cohesion configuration selected 25 segments with log evidence -4997.1478. These are distinct prior models; the values do not establish a universal preference between index and physical-length priors. Comparator scores at tolerance three are agreement with the BayesBreak MAP boundary set, not verified geology.




Figure 15. Well-log segmentation. Solid segmentation and red markers are fitted summaries; boundary probabilities and $p(k \mid y)$ are posterior quantities; the cumulative panel is in-sample.

Array-CGH. The shared-boundary model used 43 subjects on 2,215 probes, selected the maximum considered 15 segments, and recorded pooled log evidence 76359.7996 (Figure 16). The sum of independently fitted subject evidences was 109617.6954, but the two totals arise from different factorizations and are not a direct Bayes factor.

The comparator analysis uses the probe index for every method. A previous calculation applied a comparator to a flattened subject-by-probe matrix and cannot be interpreted as boundary




Figure 16. Shared-boundary array-CGH segmentation. The common red markers are fitted joint-MAP boundaries, not external copy-number annotations.

agreement; it is not included in the comparison below.

A probe-aligned comparator analysis used the exact hashed raw $2{,}215 \times 43$ probe-by-subject matrix and scored every method on the common probe-index axis (Figure 17). Agreement $F_1@3$ with the model-derived BayesBreak MAP was 0.8000 for PELT, 0.9286 for dynamic programming, 0.7143 for binary segmentation, and 0.7857 for wild binary segmentation. Dynamic programming, binary segmentation, and wild binary segmentation used the 14-boundary target. The predeclared eight-value PELT grid did not attain 14 boundaries; its closest candidate returned 11 and was not retuned post hoc. These are agreement diagnostics, not external biological accuracy or predictive superiority.

S&P 500 volatility. The source covers January 1, 2015 through December 31, 2023, with approximately 2,263 raw observations and an executed stride-four series of $n = 566$. The Gaussian log-squared-return model selected 29 segments with log evidence -1296.6537 (Figure 18). A secondary Bernoulli threshold-incidence signal selected 50 segments with log evidence -114.4100. It is a different response model, not a Poisson count, and its evidence is not directly comparable with the Gaussian total.

Methylation. The executed data are the `methylKit test1.myCpG` chromosome-21 example with 1,904 CpGs and per-CpG read coverage used as known precision. The Beta-valued quadrature model selected 15 segments and recorded log evidence -9518.6675 (Figure 19). This is not the previously described methylation-atlas analysis. The segmentation fit remains admissible descriptive evidence; the previously reported held-out value is excluded from predictive conclusions under Definition 7.13.




Figure 17. Probe-aligned array-CGH comparison. Top: boundary locations on the shared probe axis. Bottom: $F_1@3$ and exact Jaccard agreement with the model-derived BayesBreak MAP.




Figure 18. S&P 500 volatility fit after stride-four preprocessing. Red markers are fitted boundaries, not independently annotated macroeconomic events.




Figure 19. Beta-valued fit to the methylKit chromosome-21 example. Red markers are fitted MAP boundaries; no independent atlas transition annotation was used.

Scores based on a different predictive observation family or extrapolation outside the fitted domain are excluded from the held-out comparison.

A family-corrected held-out analysis used the Beta-observation predictive distribution, observed held-out coverage as positive ϕ_{new}, and `extrapolation=error`. Ten disjoint stratified interior blocks held out 1,520 CpGs while retaining both fitted-support endpoints. Total log predictive score was -23605.6749 and pooled mean score was -15.5300; the mean split score had a 95% t interval $[-23.1445, -7.9156]$. Mean boundary-stability $F_1@3$ against the model-derived full-data MAP was 0.8786 (Figure 20). The blocks are regions of one chromosome rather than independent biological samples, and no certified Beta-observation PIT or external atlas accuracy is reported. Because both family and split changed, this score is not numerically comparable with the invalid calculation.




Figure 20. Beta-response held-out prediction. Top: mean Beta-observation log predictive score across ten in-support chromosome blocks. Bottom: boundary-stability $F_1@3$ against the model-derived full-data MAP. Shaded bands are split-level 95% t intervals.

10.7 Summary of empirical findings

The empirical results demonstrate that common partition recursions can use closed-form conjugate segment marginal likelihoods, one-dimensional quadrature, other numerical segment calculations, shared-changepoint pooling, and finite latent-group templates. It provides calibrated boundary probabilities in one Gaussian simulation design, a corrected finite latent-group stress grid, finite-range runtime measurements, four traceable real-data fits, a corrected common-axis CGH agreement diagnostic, and a corrected family-specific methylation predictive evaluation. It does not yet establish external boundary accuracy, dominance under independently tuned predictive comparisons, robustness across broad misspecification regimes, routine-wide approximation guarantees, or asymptotic speed better than the quadratic dynamic-programming bound.

11 Limitations and scope

The partition recursions are exact for the stated offline, contiguous, locally factorized segmentation model when each segment marginal likelihood is exact. The following conditions delimit the method and the empirical conclusions.

Computational regime. The exact dynamic program has worst-case cost $\Theta(k_{\max} n^2)$ and is not an online changepoint detector. It is practical for moderate sequence lengths and segment-count bounds. Larger problems require a restricted candidate set, a pruning rule with its own correctness result, or an explicitly approximate calculation. The finite-range timing slopes in Section 10 do not alter the worst-case bound.

Candidate changepoints and dependence across segments. A changepoint absent from the candidate set cannot be recovered. Nonlocal interactions among segments, unrestricted dependence among segment labels, and priors that cannot be represented by local transition factors fall outside the dynamic program derived here. Some forms of dependence can be incorporated by enlarging the state space, but that produces a different recursion and requires a separate analysis.

Observation-model misspecification. Posterior conclusions are conditional on the chosen segment model, descriptor semantics, hyperparameters, and within-segment independence assumptions. Heavy tails under a Gaussian model, zero inflation under a Poisson model, overdispersion under a Binomial model, and unmodeled serial dependence can distort segment marginal likelihoods and hence the posterior over partitions. The method does not by itself identify the scientifically appropriate observation family. The reported precision-weighted array-CGH fit is a historical execution under the existing code path. Its numerical results must be recalculated after aligning Gaussian precision semantics with the implemented local likelihood; that figure does not validate the corrected weighted sampling model.

Prior sensitivity. Posterior distributions for segment count and changepoint locations can be sensitive to $p(k)$, $c_u(i, j)$, and $h_u(j)$, especially with irregularly spaced observations. Sensitivity to segment cohesion and boundary hazard should be reported. Marginal-likelihood values from analyses with different priors describe different models and should not be interpreted as a single Bayes-factor comparison without specifying the compared probability models.

Nonconjugate approximation. Theorem 7.2 is conditional on Assumption 7.1. The effect of a fixed segmentwise log-score error increases with the number of segments. A numerical method that does not establish the uniform error bound yields an exact dynamic-programming calculation for its approximate score matrix, not an error-controlled approximation to the intended posterior. In the reported numerical comparison, expectation propagation had maximum absolute segment-score discrepancy about 14.5, which is not a small-error setting.

Latent-group non-uniqueness. The finite latent-group criterion is invariant to permutation of group labels and may admit duplicate or collapsed templates. Multiple restarts do not guarantee a global maximum. In the reported stress study, the objective was nondecreasing over each recorded trajectory. This does not establish global optimality or identifiability of a generative mixture model.

Shared-changepoint assumptions. Exact pooling assumes aligned candidate coordinates and conditional independence of sequences given a common partition. Dependence among sequences or genuinely sequence-specific changepoints can make a common partition inappropriate. The finite-candidate consistency result additionally requires a positive average expected log-score gap; information from only one sequence is not sufficient as the number of sequences grows.

Posterior prediction. The observation family and coordinate support are part of the predictive model. A fallback to another family or silent assignment of out-of-range coordinates can produce a finite but invalid predictive score, as illustrated by a discarded methylation predictive calculation. Prediction outside the fitted coordinate range requires an explicit extrapolation model.

Evaluation of changepoints. Boundary F_1 depends on the tolerance and matching rule. Agreement with the BayesBreak joint-MAP set is not accuracy against independent truth. Matched

location error is undefined when no pair is matched. Red lines in the case-study plots are fitted changepoints rather than external annotations. Consequently, the reported case studies do not support causal attribution of the detected boundaries to geological, genomic, or financial events.

Incomplete empirical coverage. The current experimental evidence lacks broad repeated uncertainty intervals, a fully fairness-controlled comparator study, comprehensive misspecification grids, larger scaling studies, and method-specific error bounds for the nonconjugate approximations. These omissions limit comparative, robustness, and approximation claims even though the populated numerical values are real executed results.

BayesBreak is therefore a generalized hierarchical Bayesian method for offline segmentation of ordered data under locally factorized segment models. It is not a universal changepoint procedure. Online detection, changes in slope, reversible-jump posterior simulation, and segmentation of nonordered graphs require different models or additional theory (Adams & MacKay, 2007; Ruggieri & Lawrence, 2014; Green, 1995; Fearnhead, 2006).

12 Conclusion

We have developed a generalized hierarchical Bayesian method for multiple-changepoint segmentation with irregular designs, multiple aligned sequences, and known or latent group structure. For regular exponential-family segment models with proper conjugate priors, sufficient statistics give exact segment marginal likelihoods and posterior moments. These quantities enter common dynamic-programming recursions for the posterior distribution over segment counts and changepoint locations. A separate max-sum recursion with backtracking returns the joint MAP partition, so marginal posterior summaries and joint optimization are not conflated.

The same partition formulation accommodates design-dependent segment cohesion and boundary hazards, exact pooling of conditionally independent sequences with common changepoints, and a finite latent-group criterion optimized by alternating minorization and maximization. Nonconjugate segment models can be incorporated through numerical marginal-likelihood calculations; the resulting posterior approximation is related to the intended posterior only when a segmentwise error bound such as Assumption 7.1 is established. Posterior prediction is likewise defined by the fitted observation family and its coordinate support.

The reported numerical studies examine several observation families, Gaussian boundary calibration, a finite latent-group simulation, computational scaling, and four applied datasets. In the reported clustering configuration, hard allocation accuracy is 96%; this quantity characterizes the specified simulation, not a general recovery guarantee. The timing measurements likewise describe the tested implementation and sequence lengths, not an empirical proof of the worst-case complexity bound. For the applied datasets, changepoint agreement without an independently validated reference is not treated as accuracy against ground truth.

The additional protocols in Section 9 address limitations of the present empirical evidence, including comparator tuning, prior sensitivity, model misspecification, numerical integration accuracy, and uncertainty under replicated simulations. Until these studies are completed, no comparative or general robustness claim follows from their planned analyses. The statistical contribution remains the integration of conjugate segment models, exact posterior calculations for locally factorized partitions, irregular-coordinate priors, shared-boundary pooling, and a finite-template clustering extension under explicitly stated assumptions.

Code and data availability

The BayesBreak implementation includes experiment drivers, dataset adapters, and reproducibility specifications for the analyses reported here. Public third-party datasets retain their original distribution and licensing terms. The code and experiment configuration accompanying the published manuscript should be identified by a fixed release revision. No numerical results are imputed for the planned experiments in Section 9.

References

Ryan Prescott Adams and David J. C. MacKay. Bayesian online changepoint detection, 2007. arXiv:0710.3742.

I. E. Auger and C. E. Lawrence. Algorithms for the optimal identification of segment neighborhoods. *Bulletin of Mathematical Biology*, 51(1):39–54, 1989. doi: 10.1016/S0092-8240(89)80047-3.

Daniel Barry and J. A. Hartigan. Product partition models for change point problems. *Annals of Statistics*, 20(1):260–279, 1992. doi: 10.1214/aos/1176348521.

Daniel Barry and J. A. Hartigan. A Bayesian analysis for change point problems. *Journal of the American Statistical Association*, 88(421):309–319, 1993. doi: 10.1080/01621459.1993.10594323.

Kevin Bleakley and Jean-Philippe Vert. The group fused lasso for multiple change-point detection, 2011.

Bradley P. Carlin, Alan E. Gelfand, and Adrian F. M. Smith. Hierarchical Bayesian analysis of changepoint problems. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 41(2):389–405, 1992. doi: 10.2307/2347570.

Riccardo Corradin, Luca Danese, Wasiur R. KhudaBukhsh, and Andrea Ongaro. Model-based clustering of time-dependent observations with common structural changes. *Statistics and Computing*, 36:7, 2026. doi: 10.1007/s11222-025-10756-x. Published online 28 October 2025; volume-year citation 2026.

A. P. Dempster, N. M. Laird, and D. B. Rubin. Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)*, 39(1):1–38, 1977. doi: 10.1111/j.2517-6161.1977.tb01600.x.

Persi Diaconis and Donald Ylvisaker. Conjugate priors for exponential families. *Annals of Statistics*, 7(2): 269–281, 1979. doi: 10.1214/aos/1176344611.

Zhou Fan and Lester Mackey. Empirical bayesian analysis of simultaneous changepoints in multiple data sequences. *The Annals of Applied Statistics*, 11(4):2200–2221, 2017. doi: 10.1214/17-AOAS1075.

Paul Fearnhead. Exact and efficient Bayesian inference for multiple changepoint problems. *Statistics and Computing*, 16(2):203–213, 2006. doi: 10.1007/s11222-006-8450-8.

Paul Fearnhead and Zhen Liu. Efficient bayesian analysis of multiple changepoint models with dependence across segments. *Statistics and Computing*, 21(2):217–229, 2011. doi: 10.1007/s11222-009-9163-6.

Klaus Frick, Axel Munk, and Hannes Sieling. Multiscale change-point inference. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 76(3):495–580, 2014. doi: 10.1111/rssb.12047.

Piotr Fryzlewicz. Wild binary segmentation for multiple change-point detection. *Annals of Statistics*, 42 (6):2243–2281, 2014. doi: 10.1214/14-AOS1245.

Peter J. Green. Reversible jump markov chain monte carlo computation and Bayesian model determination. *Biometrika*, 82(4):711–732, 1995. doi: 10.1093/biomet/82.4.711.

Marcus Hutter. Exact bayesian regression of piecewise constant functions. *Bayesian Analysis*, 2(4):635–664, 2007. doi: 10.1214/07-BA225.

Tommi S. Jaakkola and Michael I. Jordan. Bayesian parameter estimation via variational methods. *Statistics and Computing*, 10(1):25–37, 2000. doi: 10.1023/A:1008932416310.

B. Jackson, J. D. Scargle, D. Barnes, S. Arabhi, A. Alt, P. Gioumousis, E. Gwin, P. Sangtrakulcharoen, L. Tan, and T. T. Tsai. An algorithm for optimal partitioning of data on an interval. *IEEE Signal Processing Letters*, 12(2):105–108, 2005. doi: 10.1109/LSP.2001.838216.

Rebecca Killick, Paul Fearnhead, and Idris A. Eckley. Optimal detection of changepoints with a linear computational cost. *Journal of the American Statistical Association*, 107(500):1590–1598, 2012. doi: 10.1080/01621459.2012.737745.

Thomas P. Minka. Expectation propagation for approximate bayesian inference. In *Proceedings of the 17th Conference on Uncertainty in Artificial Intelligence (UAI)*, pp. 362–369. Morgan Kaufmann, 2001.

Peter Müller, Fernando A. Quintana, and Gary L. Rosner. A product partition model with regression on covariates. *Journal of Computational and Graphical Statistics*, 20(1):260–278, 2011. doi: 10.1198/jcgs.2011.09066.

Adam B. Olshen, E. S. Venkatraman, Robert Lucito, and Michael Wigler. Circular binary segmentation for the analysis of array-based DNA copy number data. *Biostatistics*, 5(4):557–572, 2004. doi: 10.1093/biostatistics/kxh008.

Ju-Hyun Park and David B. Dunson. Bayesian generalized product partition model. *Statistica Sinica*, 20 (3):1203–1226, 2010.

Nicholas G. Polson, James G. Scott, and Jesse Windle. Bayesian inference for logistic models using Pólya–gamma latent variables. *Journal of the American Statistical Association*, 108(504):1339–1349, 2013. doi: 10.1080/01621459.2013.829001.

José J. Quinlan, Garritt L. Page, and Luis M. Castro. Joint random partition models for multivariate change point analysis. *Bayesian Analysis*, 19(1):21–48, 2024. doi: 10.1214/22-BA1344.

Guillem Rigaill. A pruned dynamic programming algorithm to recover the best segmentations with 1 to $k_{\max}$ change-points. *Journal de la Société Française de Statistique*, 156(4):180–205, 2015. URL https://www.numdam.org/item/JSFS_2015__156_4_180_0/. The 2010 arXiv:1004.0887 preprint is the earlier draft of this work; the JSFdS 2015 entry is the verified published version. Author-verification check in CODING_AGENT_HANDOFF.md.

Leonid I. Rudin, Stanley Osher, and Emad Fatemi. Nonlinear total variation based noise removal algorithms. *Physica D: Nonlinear Phenomena*, 60(1-4):259–268, 1992. doi: 10.1016/0167-2789(92)90242-F.

Eric Ruggieri and Charles E. Lawrence. The bayesian change point and variable selection algorithm: Application to the δ^{18}o proxy record of the plio-pleistocene. *Journal of Computational and Graphical Statistics*, 23(1):87–110, 2014. doi: 10.1080/10618600.2012.707852.

Jeffrey D. Scargle, Jay P. Norris, Brad Jackson, and Jed Chiang. Studies in astronomical time series analysis. vi. bayesian block representations. *The Astrophysical Journal*, 764(2):167, 2013. doi: 10.1088/0004-637X/764/2/167. URL https://arxiv.org/abs/1207.5578.

Henry Teicher. Identifiability of finite mixtures. *Annals of Mathematical Statistics*, 34(4):1265–1269, 1963. doi: 10.1214/aoms/1177703862.

Robert Tibshirani, Michael Saunders, Saharon Rosset, Ji Zhu, and Keith Knight. Sparsity and smoothness via the fused lasso. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 67(1): 91–108, 2005. doi: 10.1111/j.1467-9868.2005.00490.x.

Appendix

A Stable numerical implementation of posterior summaries

Family-specific integrated likelihoods, conjugate posterior updates, and complete proofs are provided directly under Propositions 4.5–4.9; the appendix does not repeat those calculations. Here we record two implementation details whose incorrect handling changes posterior inference even when every family-specific segment calculation is correct.

A.1 A separate partition-prior normalizer

For a fixed segment count, the unnormalized prior score of a path is $\prod_q \gamma_u(t_{q-1}, t_q)$. Defining $C_0(0) = 1$ and $C_0(j > 0) = 0$, its normalizer obeys

$$C_q(j) = \sum_{i=q-1}^{j-1} C_{q-1}(i)\gamma_u(i, j), \qquad C_k(u) = C_k(n). \tag{43}$$

Under $\gamma_u \equiv 1$ with all ordered paths allowed, $C_k(u) = \binom{n-1}{k-1}$. That normalizer is never one merely because the local factors equal one, except in the degenerate endpoint counts. Excluded paths receive $\gamma_u(i, j) = 0$; likelihood-zero paths are not removed from the *prior* sum.

Algorithm 3: Log-space normalization of a factorized partition prior

Input: Log prior segment factors $g_{ij} = \log \gamma_u(i, j)$ with $-\infty$ for structurally excluded blocks, and $k_{\max}$.

Initialize $P_{0,0} = 0$ and $P_{0,j>0} = -\infty$;
for $q = 1$ **to** $k_{\max}$ **do**
  for $j = q$ **to** n **do**
    $P_{q,j} \leftarrow \mathrm{LSE}_{i=q-1}^{j-1}(P_{q-1,i} + g_{ij})$;

Output: $\log C_k(u) = P_{k,n}$ for each feasible k; reject k if $C_k = 0$.

A.2 Signed first moments and block inclusion weights

Let $m_{ij}^{(r)}$ denote the posterior block moment $\mathbb{E}[m_\star(\theta)^r \mid y_{i+1:j}]$. In the forward–backward decomposition, the posterior probability that $(i, j]$ is the segment covering a specified coordinate under fixed k is

$$\omega_{ij}(k) = \frac{\widetilde{A}_{ij}^{(0)}}{L_{k,n}} \sum_{q=1}^{k} L_{q-1,i} R_{k-q,j}. \tag{44}$$

For coordinate s the first and second moments are $\sum_{i<s\le j} \omega_{ij}(k) m_{ij}^{(r)}$, $r = 1, 2$. A Gaussian posterior mean can be negative, so $\log A_{ij}^{(1)}$ is not generally defined. The numerically stable interface stores $\log A_{ij}^{(0)}$ and signed $m_{ij}^{(1)}$ separately.

Algorithm 4: Fixed-count Bayes curve with potentially negative segment means

Input: Forward/backward log messages, $\log \widetilde{A}_{ij}^{(0)}$, posterior segment moments $m_{ij}^{(1)}, m_{ij}^{(2)}$, fixed count k.

for *each valid segment* $(i, j]$ **do**
  Compute $\log \omega_{ij}(k)$ with LSE_q using Equation (44);
  Retain the sign of $m_{ij}^{(1)}$ separately from $\log |m_{ij}^{(1)}|$;
for *each coordinate* s **do**
  Sum positive and negative contributions $\omega_{ij}(k) m_{ij}^{(1)}$ for $i < s \le j$ using compensated arithmetic or signed-log-sum-exp;
  Sum nonnegative contributions $\omega_{ij}(k) m_{ij}^{(2)}$ with stabilized arithmetic;
  Set $\widehat{\mu}_s$ to the signed first-moment sum and v_s to the second-moment sum minus $\widehat{\mu}_s^2$;
  Check $v_s \ge -\delta$ for declared numerical tolerance δ, and report any failure;

Output: Fixed-k Bayes posterior-mean curve and pointwise posterior variances.

Masking rule. An unobserved outcome is absent from every likelihood and sufficient statistic. In a shared-boundary analysis, a subject with no observed outcomes in one block contributes log segment evidence zero when it has a proper segment prior. It must not contribute $-\infty$ merely because that subject supplies no information. Minimum-observation requirements are structural admissibility constraints and are represented in the prior support instead.

B Log-space dynamic programs

Algorithm 5: Log-space sum-product messages

Input: Prior-adjusted log scores $\widetilde{\ell}_{ij}$; $k_{\max}$.

Initialize $\Lambda_{0,0} = 0$, $\Lambda_{0,j>0} = -\infty$, $\mathcal{R}_{0,n} = 0$, and $\mathcal{R}_{0,i<n} = -\infty$;
for $k = 0$ **to** $k_{\max} - 1$ **do**
  for $j = k + 1$ **to** n **do**
    $\Lambda_{k+1,j} \leftarrow \mathrm{LSE}_{h=k}^{j-1}\{\Lambda_{k,h} + \widetilde{\ell}_{h,j}\}$;
  for $i \leftarrow n - k - 1, n - k - 2, \ldots, 0$ **do**
    $\mathcal{R}_{k+1,i} \leftarrow \mathrm{LSE}_{h=i+1}^{n-k}\{\widetilde{\ell}_{i,h} + \mathcal{R}_{k,h}\}$;
Normalize $\log p(k) + \Lambda_{k,n} - \log C_k(u)$ over k;

Output: Λ, $\mathcal{R}$, and $p(k \mid y)$.

Algorithm 6: Joint-MAP max-sum and backtracking

Input: Prior-adjusted log scores $\widetilde{\ell}_{ij}$; fixed k.

Set $M_{0,0} = 0$ and $M_{0,j>0} = -\infty$;
for $q = 1$ **to** k **do**
  for $j = q$ **to** n **do**
    $h^\star(q, j) \leftarrow \arg\max_{h=q-1}^{j-1}\{M_{q-1,h} + \widetilde{\ell}_{h,j}\}$ under the declared tie rule;
    $M_{q,j} \leftarrow M_{q-1,h^\star(q,j)} + \widetilde{\ell}_{h^\star(q,j),j}$;
Set $\widehat{t}_k = n$ and recursively $\widehat{t}_{q-1} = h^\star(q, \widehat{t}_q)$;

Output: $\widehat{t}$ and terminal score $M_{k,n}$.

The recommended numerical checks are

$$\sum_{k=1}^{k_{\max}} p(k \mid y) = 1,$$
$$|\Lambda_{k,n} - \mathcal{R}_{k,0}| \le \delta,$$
$$\sum_{h=1}^{n-1} p(b_h = 1 \mid y, k) = k - 1,$$
$$\left| M_{k,n} - \sum_{q=1}^{k} \widetilde{\ell}_{\widehat{t}_{q-1},\widehat{t}_q} \right| \le \delta.$$

with a documented floating-point tolerance δ.

C Supplementary agreement diagnostics

The tables below preserve executed comparator values while making the reference semantics explicit. They are not external-accuracy tables.

Table 16. Well-log boundary agreement with the BayesBreak MAP boundary set at tolerance three. No independently verified geological boundary annotation was used.

Method	Agreement $F_1@3$	Interpretation
Fearnhead-configured reference DP	0.93	agreement with BB MAP
Optimal partitioning	0.91	agreement with BB MAP
Binary segmentation	0.86	agreement with BB MAP
PELT	0.18	agreement with BB MAP
Wild binary segmentation	0.05	agreement with BB MAP

Table 17. S&P 500 and methylation comparator agreement. Segment-count restrictions were matched for the indicated comparisons.

Dataset	Method	$\widehat{k}$	Agreement $F_1@3$	Scope
S&P 500	optimal partitioning, matched k	29	0.82	vs BB Gaussian MAP
S&P 500	PELT	79	0.53	vs BB Gaussian MAP
S&P 500	binary segmentation, matched k	29	0.43	vs BB Gaussian MAP
S&P 500	wild binary segmentation, matched k	29	0.07	vs BB Gaussian MAP
S&P 500	Fearnhead-configured reference DP	13	0.55	vs BB Gaussian MAP
Methylation	optimal partitioning, matched k	15	0.79	vs BB Beta-valued MAP
Methylation	binary segmentation, matched k	15	0.57	vs BB Beta-valued MAP
Methylation	PELT	5	0.44	vs BB Beta-valued MAP
Methylation	wild binary segmentation, matched k	15	0.14	vs BB Beta-valued MAP

The incompatible array-CGH comparator output is intentionally absent from these tables. Its flattened 95,245-value axis and the 2,215-probe boundary axis do not define a common matching problem.

D Reproducibility

Reproduction of an experiment requires specifying the exact data source and coordinate system, the preprocessing and exclusion rules, the observation model and its descriptors, the prior, the numerical method, the random seeds, and the model-selection and evaluation criteria. The prospective simulations and their corresponding figures and tables are specified in Section 9; full machine-readable configurations and run instructions belong with the accompanying code.