ZOE A. HUTCHERSON (e-mail: zoe.hutcherson@enmu.edu), **KEVIN E. GREEN** (e-mail: kevin.e.green@enmu.edu), and **SEBASTIAN D. SUMMO ELIAS** (e-mail: sebastian.summoelias@enmu.edu), Department of Biology, Eastern New Mexico University, ENMU Station 33, 1500 S Avenue K, Portales, New Mexico 88130, USA; **DREW R. DAVIS**, Department of Biology, Eastern New Mexico University, ENMU Station 33, 1500 S Avenue K, Portales, New Mexico 88130, USA and Biodiversity Collections, Department of Integrative Biology, The University of Texas at Austin, 10100 Burnet Road, PRC 176-R400, Austin, Texas 78758, USA (e-mail: drew.davis@enmu.edu).

***RIOPA VOSMAERII* (Vosmaer's Writhing Skink)**. INDIA: MADHYA PRADESH: Ujjain District: Ujjain (23.1765°N, 75.7885°E; WGS 84). 9 September 2017. Ashok Bhilala. Verified by Kaushik Deuti. Zoological Survey of India (ZSI R-28931). Specimen found in a dry scrub forest in semi-arid habitat below boulder. First record from Madhya Pradesh and third specimen after original description by Gray (1839. Ann. Mag. Nat. Hist. 2:331–337). The nearest locality is from Velavadar National Park, Bhabnagar District, Gujarat, India (Vyas 2001. Hamadryad 26:360–361), ca. 400 km to the southwest. The species is also known from the vicinity of Jaggayyapet (16.8669°N, 80.0836°E; WGS 84), Krishna District, Andhra Pradesh, India (Seetharamaraju et al. 2009. J. Threat. Tax. 1:624–626), ca. 810 km to the southeast. We thank staff in the Reptilia section, ZSI for help and support.

PRATYUSH P. MOHAPATRA, Reptilia Section, Zoological Survey of India, Kolkata 700 016, West Bengal, India (e-mail: pratyush.m@zsi.gov.in); **SUMIDH RAY**, Museum & Taxidermy Section, Zoological Survey of India, Kolkata 700 016, West Bengal, India (e-mail: new.sumidh@gmail.com); **ASHOK BHILALA**, Dr. B. R. Ambedkar CM Rise School, Gulana, Shajapur 465 220, Madhya Pradesh, India (e-mail: ashok.pajwal89@gmail.com).

SQUAMATA — SNAKES

***CALLOSELASMA RHODOSTOMA* (Malayan Pitviper)**. MYANMAR: KAYIN: Kawkareik District: Kawkareik Township (ca. 16.85342°N, 97.73549°E; WGS 84), 70 m elev. 1 June 2022. N. Min. Verified by I. Das. Herpetological Collection, Duy Tan University (DTU 655; photo voucher). Individual found at 1800 h while lying motionless on timber forest covered with dry leaves; the habitat was secondary forest. First record for Myanmar as well as the northwest-most record for species; the locality is ca. 90 airline km southwest of the nearest known population in Doi Inthanon National Park, Chiang Mai Province, Thailand (Chan-ard et al. 2015. A Field Guide to the Reptiles of Thailand. Oxford University Press, New York, New York. 352 pp.). The species also occurs in southern Vietnam, central and southern Laos, Cambodia, Thailand, Penisular Malaysia, and Indonesia (Poyarkov et al. 2023. Russian J. Herpetol. 30:255–476).

TAN VAN NGUYEN, College of Medicine and Pharmacy, Duy Tan University, 120 Hoang Minh Thao, Lien Chieu, Da Nang, Vietnam (e-mail: tan.sifasv@gmail.com); **WIN PAING OO** and **NOBLE MIN**, Native Species Conservation and Identification in Myanmar, Mandalay, Myanmar; **NIKOLAY A. POYARKOV**, Department of Vertebrate Zoology, Lomonosov Moscow State University, Leninskiye Gory, GSP–1, Moscow, Russia.

***COLUBER CONSTRICTOR OAXACA* (Mexican Racer)**. USA: TEXAS: Starr Co.: intersection of FM 409 and 2844 (26.51369°N, 98.54205°W; WGS), 114.3 m elev. 29 June 2024. Collected by Mayra Oyervides, David Vela, and John Boyle. Verified by Toby Hibbits. Biodiversity Research and Teaching Collections, Texas A&M University (TCWC PV-041; photo voucher). The specimen is an adult found DOR. *Coluber c. oaxaca* is more commonly known from the sandy soils of the coastal areas of the coast of south Texas and this record represents one of the more inland representatives of the species, where the sandy soils are limited. However, the distribution is continuous with the surrounding counties and is not disjunct. The nearest vouchered specimen is ca. 40.8 km to the southeast from along the Rio Grande in Hildalgo County, Texas (TCWC 52843).

MAYRA OYERVIDES, 1708 Lavender Street, Edinburg, Texas 78539 (e-mail: mayraoyervides@hotmail.com); **DAVID VELA**, U.S. Fish and Wildlife Service, 432 Gonzalez Drive, Donna, Texas 78537 (e-mail: david_vela@usfw.gov); **JOHN BOYLE**, 3801 N McColl Road, Apt 816, McAllen, Texas 78501 (e-mail: johnboyle2020@gmail.com).

***HETERODON PLATIRHINOS* (Eastern Hog-nosed Snake)**. USA: NEW MEXICO: Union Co.: NM Highway 402, 4 km S of Amistad (35.88333°N, 103.16094°W; WGS 84). 15 May 2024. Dylan M. Osterhaus. Verified by Tomasz Giermakowski. Museum of Southwestern Biology, University of New Mexico (MSBObs:Herp:10; photo voucher). A single individual was found crossing the paved road from west to east. A new county record and only the second record from New Mexico (Painter et al. 2017. Western Wildlife 4:29–60). The nearest voucher (University of Kansas Biodiversity Institute [KUH] 72754) is from the same drainage basin ca. 28.5 km to the southeast.

DYLAN M. OSTERHAUS, Department of Fish, Wildlife, and Conservation Ecology, New Mexico State University, 2980 South Espina, Knox Hall 132, Las Cruces, New Mexico 88003, USA; e-mail: dylano@nmsu.edu.

***LAMPROPELTIS PYROMELANA* (Sonoran Mountain Kingsnake)**. USA: ARIZONA: Graham Co.: Galiuro Mountains, High Creek Trail, 0.5 km south of the Galiuro Wilderness boundary (32.58009°N, 110.26275°E; WGS 84), 1760 m elev. 16 July 2024. Charlotte A. Rozanski and Siena Webb. Verified by Andrew Holycross. Arizona State University (ASUHEP000810; photo voucher). One adult was seen crossing High Creek Trail after several days of monsoon storms. Galiuro Mountains, Paddy's River (32.61074°N, 110.28256°E; WGS 84), 1705 m elev. 21 August 2024. Ryan P. O'Donnell and Stephanie M. Cobbold. Verified by Andrew Holycross. ASUHEP000809 (photo voucher). One adult female was found active at 1350 h on the leaf litter of a river canyon, a day after monsoon storms. There are no previous vouchers of *L. pyromelana* (*sensu lato*) from the Galiuro Mountains, but the taxon was previously hypothesized to occur there (Holycross et al. 2020. *In* Holycross and Mitchell [eds.], Snakes of Arizona, pp. 196–204. ECO Publishing, Rodeo, New Mexico; Holycross et al. 2020. *In* Holycross and Mitchell [eds.], Snakes of Arizona, pp. 205–213. ECO Publishing, Rodeo, New Mexico). These two observations provide the first records of *L. pyromelana* (*sensu lato*) from the Galiuro Mountains. Burbrink et al. (2011. Mol. Phylogen. Evol. 60:445–454) proposed a split of the Sonoran Mountain Kingsnake, *L. pyromelana* (*sensu lato*), into the Arizona Mountain Kingsnake, *Lampropeltis pyromelana* (*sensu stricto*) and the Madrean Mountain Kingsnake (*Lampropeltis knoblochi*), but these specimens cannot be assigned to one of these two taxa because those authors did not attempt any morphological diagnosis, and this mountain range is about halfway through a 150–250 km ecological transition zone between the Mogollon Rim of the Colorado Plateau and the Sierra Madre Occidental that they did not sample. Genetic data, if it had been collected, might have been helpful in resolving the taxonomic status of these snakes, but also might not have

Copyright of Herpetological Review is the property of Society for the Study of Amphibians & Reptiles and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.