

Table 1: Overall Demographic and Clinical Characteristics of Long-term Ovarian Cancer Survivors (5+ years) (Grouped by Treatment Type)

Characteristic	Overall Cohort (n = 38,148)	Surgery Only (n = 12,212)	Surgery + Chemotherapy (n = 24,106)	Chemotherapy Alone (n = 750)	P-value
Age at Diagnosis, mean (SD)	54.0y (13.8y)	52.0 (15.0y)	54.6y (12.9y)	62.2y (13.9y)	<0.001
Age Group, n (%)					<0.001
20 - 39 years	4,401 (11.5%)	2,063 (16.9%)	2,157 (8.9%)	39 (5.2%)	
40 - 59 years	18,403 (48.2%)	5,914 (48.4%)	11,826 (49.1%)	218 (29.1%)	
60 - 79 years	13,811 (36.2%)	3,643 (29.8%)	9,397 (39.0%)	396 (52.8%)	
80+ years	1,533 (4.0%)	592 (4.8%)	726 (3.0%)	97 (12.9%)	
Race/Ethnicity, n (%)					<0.001
Non-Hispanic White	26,821 (70.3%)	8,317 (68.1%)	17,299 (71.8%)	512 (68.3%)	
Non-Hispanic Black	2,285 (6.0%)	844 (6.9%)	1,216 (5.0%)	85 (11.3%)	
Hispanic (All Races)	5,048 (13.2%)	1,726 (14.1%)	3,071 (12.7%)	96 (12.8%)	
Asian / Pacific Islander	3,652 (9.6%)	1,198 (9.8%)	2,333 (9.7%)	45 (6.0%)	
American Indian / Alaska Native	237 (0.6%)	72 (0.6%)	148 (0.6%)	9 (1.2%)	
Other/Unknown	105 (0.3%)	55 (0.5%)	39 (0.2%)	3 (0.4%)	
Median Household Income, n (%)					0.001
< \$60,000	3,872 (10.1%)	1,243 (10.2%)	2,348 (9.7%)	98 (13.1%)	
\$60,000 - \$99,999	25,664 (67.3%)	8,212 (67.2%)	16,260 (67.5%)	506 (67.5%)	
\$100,000+	8,606 (22.6%)	2,757 (22.6%)	5,493 (22.8%)	145 (19.3%)	
Unknown/Missing	6 (0.0%)	0 (0.0%)	5 (0.0%)	1 (0.1%)	
Geography (Rural/Urban), n (%)					0.527
Urban (Metropolitan)	34,226 (89.7%)	10,967 (89.8%)	21,671 (89.9%)	673 (89.7%)	
Rural (Nonmetropolitan)	3,889 (10.2%)	1,239 (10.1%)	2,410 (10.0%)	76 (10.1%)	
Unknown/Missing	33 (0.1%)	6 (0.0%)	25 (0.1%)	1 (0.1%)	
Cancer Stage, n (%)					<0.001
Localized	11,086 (29.1%)	7,063 (57.8%)	3,905 (16.2%)	19 (2.5%)	
Regional	11,478 (30.1%)	3,121 (25.6%)	8,191 (34.0%)	87 (11.6%)	
Distant	14,107 (37.0%)	1,639 (13.4%)	11,677 (48.4%)	554 (73.9%)	
Unknown/unstaged	1,477 (3.9%)	389 (3.2%)	333 (1.4%)	90 (12.0%)	

y, years; SD, standard deviation