
a Eleven datasets from ten studies
bar colour = habitat; text = environmental samples the plates came from
Hinsu 2021
2 source samples
Kruger 2019
2 source samples
Li 2023
2 source samples
Zheng 2024 (soil)
6 source samples
Verhoeven 2023
2 source samples
Villanelo 2026
1 source sample
Zheng 2024 (sludge)
5 source samples
Indraningrat 2019
12 source samples
Ordonez 2025
3 source samples
Yu 2020
15 source samples
Zehavi 2018
1 source sample
10^0
10^1
10^2
10^3
Plating conditions analysed (log scale)
b What the analysed plates covered
rich
dilute
defined
habitat extract
selective
not classed
cold (≤20 °C)
moderate
warm (≥33 °C)
not resolved
aerobic
anaerobic
microaerophilic
not stated
V3-V4, 341F/806R
V3-V4, 341F/806R
full length, cut to V4, 515F/806R
V4, 515F/806R
V1-V3, 27F/534R
full length, cut to V4, 515F/806R
V4, 515F/806R
V4, 515FY/806RB
V4, 515F/806R (modified)
V3-V4, 341F/806R (modified)
V4, 515F/806R
Medium class
Temperature
Oxygen
16S region and primers
soil
activated sludge
marine sponge
plant (oak)
food (broiler carcass)
rumen
condition present
present but class not known