

Some subtleties of spherical coordinates

Jonathan Delgado


January 22, 2026

Abstract

The following are some reflections on why the polar angle coordinate ϕ does not globally define a 1-form on S^2 and hence on $\mathbb{R}^3 \setminus \{0\}$.

1 Review of Spherical Coordinates

Recall that any point $(x, y, z) \in \mathbb{R}^3$ can be described by its distance from the origin, $\rho = \sqrt{x^2 + y^2 + z^2}$, its **polar angle** ϕ^1 , and its **azimuthal angle** θ . From this picture we have $\rho \in [0, \infty)$, $\phi \in [0, \pi]$,



and $\theta \in [0, 2\pi)$. Despite the care we take with these possible values, we have still introduced several

¹We are following the mathematics convention: https://en.wikipedia.org/wiki/Spherical_coordinate_system, as opposed to the physics convention which swaps the roles of ϕ and θ .

degeneracies: various choices of “coordinates” that represent the same point. For example, at $\rho = 0$, we identify

$$(0, \phi, \theta) \sim (0, \phi', \theta').$$

Similarly, on the positive z -axis,

$$(\rho, 0, \theta) \sim (\rho, 0, \theta') \quad \text{and} \quad (\rho, \pi, \theta) \sim (\rho, \pi, \theta').$$

As a result, spherical coordinates only really describe a coordinate system in the rigorous sense provided

$$(\rho, \phi, \theta) \in (0, \infty) \times (0, \pi) \times [0, 2\pi).$$

This covers $\mathbb{R}^3 \setminus$ (the z -axis), and we can only guarantee the existence of $d\rho$, $d\phi$, and $d\theta$, on this domain.

2 The problem of the 1-form

From the equation for z we have

$$\phi = \arccos\left(\frac{z}{\rho}\right) = \arccos\left(\frac{z}{\sqrt{x^2 + y^2 + z^2}}\right).$$

The domain for \arccos is $[-1, 1]$ and the inequality $|z| \leq \rho$ ensures that ϕ is a globally defined smooth function on $\mathbb{R}^3 \setminus$ (the z -axis). We have more, however, since this expression for ϕ actually shows that it's well-defined on $\mathbb{R}^3 \setminus \{0\}$. This makes it tempting to conclude that $d\phi$ exists on $\mathbb{R}^3 \setminus \{0\}$. It turns out that this is not the case, $d\phi$ is not even a 1-form on S^2 .

2.1 Direct computation

We can argue that $d\phi$ fails to exist on S^2 by a direct computation.

$$\begin{aligned} d\phi &= d \arccos\left(\frac{z}{\rho}\right) = -\frac{1}{\sqrt{1 - \frac{z^2}{\rho^2}}} d\left(\frac{z}{\rho}\right) = -\frac{1}{\sqrt{\frac{x^2 + y^2}{\rho^2}}} \frac{\rho dz - z d\rho}{\rho^2} \\ &= -\frac{1}{\sqrt{\frac{(\rho \sin \phi)^2}{\rho^2}}} \frac{\rho dz - z d\rho}{\rho^2} = -\frac{1}{\sin \phi} \left(\frac{1}{\rho} dz - \frac{z}{\rho^2} d\rho\right). \end{aligned}$$


Since we're away from the origin, $\rho > 0$ and smooth. So the only possible issue for $d\phi$ happens at $\phi \in \{0, \pi\}$. Since the factor on the right does not vanish for these values of ϕ , it follows that $d\phi$ is not globally defined on S^2 .

This is not an exceptionally insightful calculation, and is also tedious to do. Instead, we will discuss a more geometric way to see why this 1-form does not exist.

2.2 A geometric argument

Focusing on the upper z -axis, let's see what happens as ϕ gets close to 0. The distance any given point has from the z -axis is $\rho \sin \phi$. For small values of ϕ the small angle approximation $\sin \phi \approx \phi$ (justified by Taylor expansion) says that the point is approximately ϕ -distance away from the axis. On the other hand, spherical coordinates also tell us that

$$\rho \sin \phi = r = \sqrt{x^2 + y^2}.$$



Our small angle approximation tells us $\phi \approx r$ and the problem of differentiating ϕ near the z-axis is exactly the problem of differentiating r in polar coordinates. In particular

$$\phi \approx \sqrt{x^2 + y^2},$$

and the derivative of this explodes at 0, as seen by

$$dr = \frac{1}{\sqrt{x^2 + y^2}} d(x^2 + y^2).$$

3 Consequences

The observation $\phi \approx r = \sqrt{x^2 + y^2}$ for small values of ϕ offers a suggested way to smooth it over by instead considering $d\phi^2$. In this case

$$d\phi^2 = 2\phi d\phi$$

and a direct computation shows that $\phi d\phi \rightarrow 0$ as $\phi \rightarrow 0$ for the same reason $r dr \rightarrow 0$ as $r \rightarrow 0$.

Notice that this still fails to fix the issue on the negative z-axis since here $\phi \rightarrow \pi$ which does not remove the singular behavior of its derivative given by the same picture. To justify the small-angle approximation, we need to instead consider the complementary angle $\psi = \pi - \phi$. Now $\phi \rightarrow \pi$ is the same as considering $\psi \rightarrow 0^+$ and our previous argument holds. This means that, away from the positive z-axis,

$$d\psi^2 = (\pi - \phi)d(\pi - \phi) = -(\pi - \phi)d\phi$$

is well-defined.