

UNIVERSITY OF OKLAHOMA

GRADUATE COLLEGE

SEARCHING FOR SUPERSYMMETRIC PARTICLES AT THE LARGE HADRON
COLLIDER USING THE ATLAS DETECTOR

A DISSERTATION

SUBMITTED TO THE GRADUATE FACULTY

in partial fulfillment of the requirements for the

Degree of

DOCTOR OF PHILOSOPHY

By

OTHMANE RIFKI
Norman, Oklahoma
2016

SEARCHING FOR SUPERSYMMETRIC PARTICLES AT THE LARGE HADRON
COLLIDER USING THE ATLAS DETECTOR

A DISSERTATION APPROVED FOR THE
HOMER L. DODGE DEPARTMENT OF PHYSICS AND
ASTRONOMY

BY

Dr. Brad Abbott, Chair

Dr. Peter Parker, Outside Member

Dr. Mike Strauss

Dr. Chung Kao

Dr. Eric Abraham

To my loving parents and sister.

Acknowledgements

I would like to acknowledge...

Table of Contents

List of Tables	vi
List of Figures	vii
Abstract	viii
1 Theoretical Background	4
2 The LHC and the ATLAS Experiment	5
3 The Region of Interest Builder	6
4 Object Reconstruction	7
5 Data-driven techniques to estimate fake lepton backgrounds	8
5.1 The problem of fakes	8
5.2 Common processes for faking leptons	9
5.3 Monte Carlo Template Method	11
5.3.1 Motivation	11
5.3.2 Description of the method	12
5.3.3 Correction factors	13
5.3.4 Control regions	14
5.4 Matrix Method	16
5.4.1 Events with one object	17
5.4.2 Dynamic matrix method	20
5.4.3 Propagation of uncertainties	22
5.5 Real lepton efficiency	23
5.6 Fake lepton rate	23
6 Search for new physics in events with two same sign leptons or three leptons and jets	24
7 Statistical Analysis	25

List of Tables

5.1	The FNP and charge flip multipliers obtained after minimizing the likelihood function using Pythia. The uncertainty in the multipliers takes into account the limited statistics of simulated events.	20
5.2	The FNP and charge flip multipliers obtained after minimizing the likelihood function using Sherpa. The uncertainty in the multipliers takes into account the limited statistics of simulated events.	21

List of Figures

5.1	Lepton classification.	13
5.2	Pre-fit distributions for ee channel (left), for $e\mu$ channel (middle), and for $\mu\mu$ channel (right) from CR0b that were used in the fit to extract the FNP lepton and charge flip multipliers. The generator used in these plots is Powheg. The hashed band represents the sum of systematic uncertainties on the predictions.	16
5.3	Post-fit distributions for ee channel (left), for $e\mu$ channel (middle), and for $\mu\mu$ channel (right) from CR0b that were used in the fit to extract the FNP lepton and charge flip multipliers. The generator used in these plots is Powheg. The hashed band represents the sum of systematic uncertainties on the predictions.	17
5.4	Pre-fit distributions for ee channel (left), for $e\mu$ channel (middle), and for $\mu\mu$ channel (right) from CR1b that were used in the fit to extract the FNP lepton and charge flip multipliers. The generator used in these plots is Powheg. The hashed band represents the sum of systematic uncertainties on the predictions.	18
5.5	Post-fit distributions for ee channel (left), for $e\mu$ channel (middle), and for $\mu\mu$ channel (right) from CR1b that were used in the fit to extract the FNP lepton and charge flip multipliers. The generator used in these plots is Powheg. The hashed band represents the sum of systematic uncertainties on the predictions.	19

Abstract

Well now this is my abstract.

Preface

...

In the spring of 2014, I received an ATLAS Support Center graduate fellowship to move to Argonne National Laboratory to work on hardware and physics analysis. This thesis is based on the work conducted in the period of May 2014 to May 2017. In my research, the supervision of Dr. Brad Abbott has been invaluable.

There is a very long list of people I would like to thank. I would like to thank the Argonne group for the warm welcome to the physics division at Argonne. The scientists at Argonne became colleagues and friends.

The working conditions either in Oklahoma, Argonne, or CERN have been excellent. I had the privilege of interacting with many interesting people at these institutes which I contributed significantly to building my independence as a physicist. I have been particularly fortunate of being part of Argonne National Laboratory, where I am able to work with many talented physicists. I would like to start with Alexander Paramonov and Sergei Chekanov who were the first to introduce me to physics analysis. Robert Blair, Jeremy Love, and Jinlong Zhang with whom I evolved the glorious region of interest builder of ATLAS responsible for processing every event recorded by the ATLAS detector. It was a real joy working with them and bringing the project to completion and taking data in ATLAS. It is fair to say that James Proudfoot became a mentor to me during my stay at Argonne and afterwards. Furthermore, my gratitude goes to Ximo Poveda, Julien Maurer, and Otilia Ducu who made my work within the same sign

SUSY group a real joy.

Finally, I would like to thank my family and frineds for their support and acceptance of my absence from their lives. Last, but not least, I thank ... encouragement and patience during these years.

Othmane Rifki, May 2017

Introduction

Chapter 1

Theoretical Background

Chapter 2

The LHC and the ATLAS Experiment

Chapter 3

The Region of Interest Builder

Chapter 4

Object Reconstruction

Chapter 5

Data-driven techniques to estimate fake lepton backgrounds

5.1 The problem of fakes

The reconstructed objects (leptons, photons, b -jets, etc.) in a collision event are used to perform a wide range of SM measurements or searches for evidence of BSM physics. The assumption is that these objects are ‘real’ representing the desired particles in the final state used in the analysis. In practice, the reconstructed objects might not be always ‘real’. In fact, they may be something completely different that was mistakenly reconstructed as the desired object, called ‘fake’. For the purpose of the analysis presented in this thesis, the focus is on ‘fake’ leptons. To illustrate the problem, a hadronic jet may deposit more energy in the electromagnetic calorimeter than the hadronic calorimeter, or that it leaves a narrow deposit of energy leading the reconstruction algorithms to mistake this jet for an electron. From the analysis point of view, the ‘fake’ electron will pass all the selection criteria and will be indistinguishable from a ‘real’ electron. It is important for the analysis that requires a reconstructed electron to model the fake electron background to get a sound result. This example was given with electrons, but can be generalized to muons as well. In short, any analysis that uses leptons in the final state must account for the ‘fake’ lepton background. This background can be more or less important depending on the detector, the analysis selection,

and the number of leptons required. To estimate this background it is important to first understand what type of processes lead to fake leptons.

5.2 Common processes for faking leptons

The reconstruction of ‘fake’ leptons can be an instrumental effect related to the inability to identify the object based on its measured properties by the detector. In this case, the reconstructed lepton is not a real lepton and the production process will be different for electrons and muons.

The reconstruction of electrons relies on the observation of well aligned particle hits in the layers of the ID that are consistent with an energy deposition in the EM calorimeter. Photons can mimick this signature since they deposit energy in the EM calorimeter that happens to be aligned with a charged track. A jet for example containing charged and neutral pions can lead to such scenarios. It is possible for the jet to have one charged pion leaving a track similar to that of an electron. The decay of π^0 mesons to photons in this jet can deposit energy in the EM calorimeter leading to the required signature. Another mechanism that can lead to fake electrons is the emission of photons via Brehmstrahlung from high energy muons. The muon track can be mistaken for that of an electron and the photons interact with the EM calorimeter leading to a signature similar to that of electrons. An additional process is that of photon conversions into a e^+e^- .

The reconstruction of muons relies on the observation of tracks from the ID matched to tracks from the muon spectrometer. It is possible for charged hadrons

with long lifetime to traverse the calorimeter layers and leave hits in the muon spectrometer. These hits may coincide with other hits from the ID due to the random activity in the event. As a result, a muon can get reconstructed. Another instance may occur when pions or kaons decay in-flight to muons in the muon spectrometer and happen to align with the primary vertex.

The leptons that are used in the physics analyses must be coming from the hard scatter, generally referred to as prompt leptons. There is another case where the reconstructed lepton is a real lepton but is not a lepton coming from the hard interaction, referred to as non-prompt leptons. Non-prompt leptons can be produced from heavy flavor meson decays with a low energy activity around the lepton which allows it to pass isolation requirements. A good example of this type of process is the semi-leptonic decay of top quark pair which contribute to final states with two leptons.

For the rest of the thesis, the fake leptons will be referred to as fake/non-prompt (FNP) leptons. There are several methods used to perform the estimation of FNP lepton backgrounds. A method that the author developed will be described next along with a standard method for estimating this type of backgrounds. The benefit of having two methods for estimating the FNP lepton background is to have enough confidence in the final estimate. The two methods use different assumptions which naturally leads to a more robust estimation of this difficult background. Moreover, the final estimate of the FNP lepton background is taken as a statistical combination of the estimates from the two methods leading to a reduction of the systematic uncertainties on the estimate.

5.3 Monte Carlo Template Method

5.3.1 Motivation

The processes leading to FNP leptons depend on the selection applied in the analysis. For instance, a selection with same-sign leptons will have contributions from top quark pair production ($t\bar{t}$) or the associated production of a vector boson and jets (W +jets or Z +jets). These processes cannot give two leptons of the same electric charge unless there is a charge mis-measurement (mainly affecting electrons) or that a FNP lepton was produced. It is possible to generate the processes that can contribute to a FNP lepton, such as $t\bar{t}$ or V +jets, with Monte Carlo event generators processed through Geant4 detector simulation of the ATLAS detector. This approach will yield an estimate however it might not be reliable. For instance, the detector simulation itself might not reproduce the true behavior of the interaction of the physics objects with the detector, particularly when looking at rare processes such as the production of FNP leptons. The second limitation is in the generation of enough MC events to probe the region of the phase space targetted by the analysis which affects the statistical uncertainties in the estimates. The latter concern is addressed by ensuring that the simulations for the major backgrounds ($t\bar{t}$ and V +jets) have much higher event count than the corresponding number of events observed in the data sample. In fact, these backgrounds have a large number of simulated events because they are important for many analyses (including SM measurements and BSM searches). The rest of the section will concentrate on addressing the former limitation.

5.3.2 Description of the method

The MC template method relies on the correct modelling of FNP leptons kinematics in MC simulation to extrapolate background predictions from control regions to the signal regions. The method assumes that the kinematic shapes for each source of FNP lepton is correctly modeled in the simulations, and the normalization for each source is extracted in a combined fit to data control regions. The number of normalization factors depend on the number of identified origins of the FNP lepton in the signal regions and the control regions are designed to constrain these factors in regions enriched with FNP leptons from the same origin.

To illustrate the approach, we describe the application of the method in SS/3L analysis later described in this thesis. The processes of interest that may lead to a FNP lepton or a charge flip are $t\bar{t}$ and V +jets. FNP leptons are classified using an algorithm that navigates the generator particle record to determine where the FNP lepton is originating from. The lepton is classified as either an electron or a muon that is prompt from decays of on-shell W and Z bosons, non-prompt from a heavy flavor b decay (HF), or fake from mis-identification of a light flavor jet or a photon (LF). In the case of an electron, we further classify the prompt electrons to prompt electrons with the correct charge or with a charge mis-measurement, commonly named charge flip. In total, five categories referred to as MC templates are constructed following the classification illustrated in figure 5.1.




Figure 5.1: Lepton classification.

5.3.3 Correction factors

The FNP estimate relies on kinematic extrapolation using processes expected to contribute via FNP leptons from control regions with low jet multiplicity and E_T^{miss} , to the signal regions that require high jet multiplicity and E_T^{miss} . The control regions are chosen to separate FNP leptons from HF origins and FNP leptons from LF origins. For instance, a control sample characterized by the presence of a b -jet will be enriched in processes with one FNP lepton that is coming from a HF decay, while a sample characterized by the absence of a b -jet will have one FNP lepton from LF decay. The presence of one FNP lepton in the control sample allows the correction of the production rate of these FNP leptons by performing a fit to data.

For example, if a $Z \rightarrow \mu\mu + \text{LF jet}$ event is reconstructed as a $\mu^+\mu^-e^+$ event, then the electron is fake. Therefore, a correction of $\text{LF jet} \rightarrow e$ ($\text{Fr}(\text{LF} \rightarrow e)$) is applied to the rate of $\mu\mu e$ events. The correction $\text{Fr}(\text{LF} \rightarrow e)$ is constrained by a fit to data in control regions dominated by $\text{LF jet} \rightarrow e$ type fakes. Similarly, three other corrections are defined as $\text{LF jet} \rightarrow \mu$ ($\text{Fr}(\text{LF} \rightarrow \mu)$), $\text{HF jet} \rightarrow e$

($\text{Fr}(\text{HF} \rightarrow e)$), $\text{HF jet} \rightarrow \mu$ ($\text{Fr}(\text{HF} \rightarrow \mu)$). An additional correction is applied to correct the charge flip rate predicted by simulation. For example, a $Z \rightarrow e^+e^-$ event is reconstructed as e^+e^+ or e^-e^- . The simulation takes into account the charge flip rate but it might be off. The charge flip ($\text{Cf}(e)$) correction derived from a data fit is expected to recover this mis-modeling. The charge flip rate only concern electrons as the muon charge flip rate is negligible.

A likelihood fit is defined as the product of the Poisson probabilities describing the observed events in the binned distributions from the expected number of events rescaled by the five multipliers which are left free to float in the fit. These multipliers are applied to the MC predictions in the signal regions to obtain an estimation of the charge flip and FNP backgrounds.

5.3.4 Control regions

The corrections depend on the simulated sample, the reconstructed final state, and the flavor of the leptons. As a result, care must be taken when designing the control regions used to perform the fit of the FNP leptons and electron charge flip templates. For instance, each template needs to be constrained in a selection that is representative of the processes leading to FNP leptons and charge flip electrons present in the kinematic region targetted by the search for BSM physics.

In the SS/3L analysis discussed in this thesis the control regions are defined with at least two same-sign leptons, $E_{\text{T}}^{\text{miss}} > 40$ GeV, two or more jets. This preselection ensures that the FNP leptons are not from fakes originating from QCD like event topologies. They are further split in regions with or without

b -jets to constrain the HF and LF leptons respectively. In addition, they are also split with different flavours of the same-sign lepton pair ee , $e\mu$, and $\mu\mu$, giving a total of six control regions. Any event entering the signal region is vetoed. The ee channel will constrain the charge flip correction factor, fake leptons from LF decays in the selection without b -jets, and non-prompt decay from HF in the selection with b -jets. The $\mu\mu$ channel will constrain the muon fake rates in the LF and HF decays for the selection without or with b -jets, respectively. The $e\mu$ channel will constrain both the electron and muon fakes for events containing both lepton flavors.

The six distributions are chosen for variables that provide the best separation between processes with prompt leptons and processes with FNP leptons and charge flip and are shown before and after the fit in Figures 5.2-5.4 and Figures 5.3-5.5, respectively.

The minimization of the negative log likelihood using the MINUIT package leads to the multipliers shown in Tables 5.1 and 5.2. The tables represent the multipliers obtained from the fit upon using two different parton showers, POWHEG-BOX and SHERPA for the processes that lead to FNP leptons and charge flips. The systematic uncertainty is obtained by varying the generator from POWHEG-BOX to SHERPA and evaluating the impact on the expected background from FNP and charge flip leptons. It is found to be the dominant contribution to the systematic uncertainty of the method (up to 80%). The uncertainties in the multipliers themselves correspond to how much the parameter needs to be varied for a one standard deviation change in the likelihood function. This uncertainty takes into




Figure 5.2: Pre-fit distributions for ee channel (left), for $e\mu$ channel (middle), and for $\mu\mu$ channel (right) from CR0b that were used in the fit to extract the FNP lepton and charge flip multipliers. The generator used in these plots is Powheg. The hashed band represents the sum of systematic uncertainties on the predictions.

account the limited number of simulated events and is included as a systematic uncertainty on the expected number of background events.

5.4 Matrix Method

The FNP leptons do not often pass one of the lepton selection criteria but have non-zero impact parameter, and are often not well-isolated. These selection requirements are key ingredients to control the FNP leptons. The number of events with at least one FNP lepton is estimated using two classes of leptons: a real-enriched class of “tight” leptons corresponding to signal leptons and a fake-enriched class of “loose” leptons corresponding to candidate leptons with




Figure 5.4: Pre-fit distributions for ee channel (left), for $e\mu$ channel (middle), and for $\mu\mu$ channel (right) from CR1b that were used in the fit to extract the FNP lepton and charge flip multipliers. The generator used in these plots is Powheg. The hashed band represents the sum of systematic uncertainties on the predictions.

$$\begin{pmatrix} n_{\text{pass}} \\ n_{\text{fail}} \end{pmatrix} = \begin{pmatrix} \varepsilon & \zeta \\ 1 - \varepsilon & 1 - \zeta \end{pmatrix} \begin{pmatrix} n_{\text{real}} \\ n_{\text{FNP}} \end{pmatrix}; \quad (5.1)$$

allowing to determine the unknown number of events n_{FNP} from the observed n_{pass} and n_{fail} given measurements of the probabilities ε/ζ .

The predictive power of the matrix method comes from the fact that the real and FNP leptons have different composition in the two collections of tight and loose objects leading to $\varepsilon \neq \zeta$. In fact, the tight lepton collection will be dominated by real objects while the loose region will be dominated by fake objects. As a result, the inequality $\varepsilon \gg \zeta$ will always hold true which guarantees that the matrix in Eq. 5.1 is invertible and gives positive estimates.

The next step is to invert the relation in Eq. 5.1 to obtain




Figure 5.5: Post-fit distributions for ee channel (left), for $e\mu$ channel (middle), and for $\mu\mu$ channel (right) from CR1b that were used in the fit to extract the FNP lepton and charge flip multipliers. The generator used in these plots is Powheg. The hashed band represents the sum of systematic uncertainties on the predictions.

$$\begin{pmatrix} n_{\text{real}} \\ n_{\text{FNP}} \end{pmatrix} = \frac{1}{\varepsilon - \zeta} \begin{pmatrix} \bar{\zeta} & -\zeta \\ -\bar{\varepsilon} & \varepsilon \end{pmatrix} \begin{pmatrix} n_{\text{pass}} \\ n_{\text{fail}} \end{pmatrix}; \quad (5.2)$$

where $\bar{\varepsilon} = 1 - \varepsilon$ and $\bar{\zeta} = 1 - \zeta$. The FNP lepton component is:

$$n_{\text{FNP}} = \frac{1}{\varepsilon - \zeta} (\varepsilon (n_{\text{pass}} + n_{\text{fail}}) - n_{\text{pass}}). \quad (5.3)$$

However, the quantity of interest is the expected FNP lepton background that passes the tight selection criteria: $n_{\text{pass} \cap \text{FNP}} = \zeta n_{\text{FNP}}$. To obtain this quantity, the identity from Eq. 5.1 is used to get:

$$n_{\text{pass} \cap \text{FNP}} = \frac{\zeta}{\varepsilon - \zeta} (\varepsilon (n_{\text{pass}} + n_{\text{fail}}) - n_{\text{pass}}). \quad (5.4)$$

Table 5.1: The FNP and charge flip multipliers obtained after minimizing the likelihood function using Pythia. The uncertainty in the multipliers takes into account the limited statistics of simulated events.

Category	Multiplier	Uncertainty
chFlip	1.49	0.58
HF EL	2.80	0.98
LF EL	2.89	0.88
HF MU	1.59	0.31
LF MU	1.00	1.34

5.4.2 Dynamic matrix method

This readily generalizes to events with more than one lepton, with a formalism described in details. An implementation of this formalism (known also as the dynamic or generalized matrix method) can handle an arbitrary number of leptons in the event. The method should be applied event-by-event, effectively resulting into a weight being assigned to each event. The predicted yield of events with FNP leptons is simply the sum of weights.

The application of the matrix method to multilepton final states comes with two specificities. Firstly, contributions of events with charge-flip electrons would bias a straightforward matrix method estimate (in particular for a final state formed by two leptons with same electric charge). This happens because the candidate-to-signal efficiency for such electrons is typically lower than for real

Table 5.2: The FNP and charge flip multipliers obtained after minimizing the likelihood function using Sherpa. The uncertainty in the multipliers takes into account the limited statistics of simulated events.

Category	Multiplier	Uncertainty
chFlip	1.34	0.58
HF EL	2.40	0.85
LF EL	1.83	1.04
HF MU	1.17	0.16
LF MU	2.40	0.81

electrons having a correctly-assigned charge. One therefore needs to subtract from n_{pass} and n_{fail} the estimated contributions from charge-flip. This can be performed by including as well events with pairs of opposite-sign candidate leptons in the matrix method estimate, but assigning them an extra weight corresponding to the charge-flip weight. Thanks (again) to the linearity of the matrix method with respect to n_{pass} and n_{fail} , this weight-based procedure is completely equivalent (but more practical) to the aforementioned subtraction.

Secondly, the analytic expression of the matrix method event weight depends on the lepton multiplicity of the final state. This concerns events with three or more candidate leptons: one such event takes part both in the evaluation of the FNP lepton background for a selection with two signal leptons or a selection with

three signal leptons, but with different weights². Therefore, for a given event used as input to the matrix method, one should consider all possible leptons combinations, each with its own weight and its own set of kinematic variables. For example, a $e^+e^-\mu^+$ event is comptabilized in the background estimate both as an $e^+\mu^+$ event (with a weight w_1) and as an $e^+e^-\mu^+$ event (with a weight $w_2 \neq w_1$).

5.4.3 Propagation of uncertainties

The two parameters (ε and ζ respectively) can be measured in data, and depend on the flavor and kinematics of the involved leptons. Systematic uncertainties resulting from the measurement of these two parameters, and their extrapolation to the signal regions, can be propagated to uncertainties on the event weight through standard first-order approximations. The different sources of uncertainties should be tracked separately so that correlations of uncertainties across different events can be accounted for correctly. The resulting set of uncertainties on the cumulated event weights can be then added in quadrature to form the systematic uncertainty on the predicted FNP lepton background yield. The corresponding statistical uncertainty can be taken as the RMS of the event weights.

²This can appear for inclusive selections: for example an event with two signal leptons may or not contain additional candidate leptons, in a transparent way

5.5 Real lepton efficiency

5.6 Fake lepton rate

The FNP leptons efficiency parameter should be measured in a CR dominated by same FNP lepton sources as the signal regions. The definition of this control region is analysis dependent, and dedicated truth composition studies should be performed to understand the different sources of FNP leptons in the signal and FNP leptons enriched CR regions.

Chapter 6

Search for new physics in events with two same sign leptons or three leptons and jets

Chapter 7

Statistical Analysis

Conclusion