

i, j, k, l, m, n

$ident, x$

$constant ::=$
 $| \quad ()$

$expr, e ::=$
 $| \quad constant$

$subst_x ::=$
 $| \quad x_1 < -e_1, \dots, x_n < -e_n$

$valu, v ::=$
 $| \quad constant$

$formula ::=$
 $| \quad judgement$
 $| \quad formula_1 \dots formula_n$

$Jred ::=$
 $| \quad v << subst_x >>$
 $| \quad (v_1 \& \dots \& v_n) << subst_x >>$

$judgement ::=$
 $| \quad Jred$

$user_syntax ::=$
 $| \quad i$
 $| \quad ident$
 $| \quad constant$
 $| \quad expr$
 $| \quad subst_x$
 $| \quad valu$
 $| \quad formula$

$v << subst_x >>$

$$\frac{v << x_1 < -v_1, \dots, x_n < -v_n >>}{v << x_1 < -v_1, \dots, x_n < -v_n, x < -v >>}$$
 JR_MATCH_ALIAS

$(v_1 \& \dots \& v_n) << subst_x >>$

$$\frac{v_1 << subst_x >> \dots v_n << subst_x >>}{(v_1 \& \dots \& v_n) << subst_x >>}$$
 JR_MATCHES_AND

Definition rules: 2 good 0 bad

Definition rule clauses: 4 good 0 bad