

SEDAR 36: South Atlantic Snowy Grouper

Research recommendations from the Standard assessment report.

- Increased fishery independent information, particularly for developing reliable indices of abundance, would greatly improve the assessments of deepwater species.
- More age samples should be collected from the general recreational sector and with more complete spatial coverage.
- Snowy grouper were modeled in this assessment as a unit stock off the southeastern U.S. For any stock, variation in exploitation and life-history characteristics might be expected at finer geographic scales. Modeling such sub-stock structure would require more data, such as information on the movements and migrations of adults and juveniles, as well as spatial patterns of larval dispersal and recruitment. Even when fine-scale spatial structure exists, incorporating it into a model may or may not lead to better assessment results (e.g., greater precision, less bias). Spatial structure in a snowy grouper assessment model might range from the very broad (e.g., a single Atlantic stock) to the very narrow (e.g., a connected network of meta-populations living on individual reefs). What is the optimal level of spatial structure to model in an assessment of snapper-grouper species such as snowy grouper? Are there well defined zoogeographic breaks (e.g., Cape Hatteras) that should define stock structure? Research into these questions could help inform future stock assessments.
- Protogynous life history: 1) Investigate possible effects of hermaphroditism on the steepness parameter; 2) Investigate the sexual transition for temporal patterns, considering possible mechanistic explanations if any patterns are identified; 3) Investigate methods for incorporating the dynamics of sexual transition in assessment models.
- In this assessment, the number of spawning events per mature female per year was implicitly assumed to be constant. The underlying assumptions are that spawning frequency and spawning season duration do not change with age or size. Research is needed to address whether these assumptions for snowy grouper are valid. Age or size dependence in spawning frequency and/or spawning season duration would have implications for estimating spawning potential as it relates to age structure in the stock assessment (Fitzhugh et al. 2012).