

Section 2.1

Exercise 5

$$\forall x (x \in A \leftrightarrow x \in B) \leftrightarrow A = B$$

- a) $A = B$ since A contains all elements of B and vice versa
- b) $((1)) \in A \wedge ((1)) \in B$ but $(1) \notin A \rightarrow A \neq B$
- c) \emptyset is an empty set, and is not equal to a set containing an empty set.

Exercise 24

- a) No. The power set of a set always include a set of an empty set: $P(\emptyset) = (\emptyset)$ meaning a power set of a set is never empty. $\forall A ((\emptyset) \in P(A)) \rightarrow P(A) \neq \emptyset$
- b) Yes. $P((a)) = (\emptyset, (a))$
- c) No. $P((\emptyset, a)) = (\emptyset, (a), (\emptyset), (a, \emptyset))$ The set in question is missing a (\emptyset)
- d) Yes. $P((a, b)) = (\emptyset, (a), (b), (a, b))$

Section 2.2

Exercise 18

- c) $(A - B) \equiv \forall x \mid x \in A \wedge x \notin B$
 $(A - B) - C \equiv \forall x \mid x \in (A - B) \wedge x \notin C$
 $(A - B) - C \equiv \forall x \mid x \in A \wedge x \notin B \wedge x \notin C$
 $(A - C) \equiv \forall x \mid x \in A \wedge x \notin C$
 Since all elements of $(A - B) - C$ are in $(A - C)$, $(A - B) - C \subseteq (A - C)$

- d) $(A - C) \equiv \forall x \mid x \in A \wedge x \notin C$
 $(C - B) \equiv \forall x \mid x \in C \wedge x \notin B$
 $(A - C) \cap (C - B) \equiv \forall x \mid x \in (A - C) \wedge x \in (C - B)$
 $(A - C) \cap (C - B) \equiv \forall x \mid (x \in A \wedge x \notin C) \wedge (x \in C \wedge x \notin B)$

Rewriting the paranthesis we get

$$(A - C) \cap (C - B) \equiv \forall x \mid x \in A \wedge \underbrace{(x \notin C \wedge x \in C)}_{x \in \emptyset} \wedge x \notin B$$

Clearly, $(A - C) \cap (C - B) = \emptyset$

Exercise 46

The set $A \cup B$ consists of all elements in A that are also in B.

$$A \cup B = \{x \mid x \in A \wedge x \in B\}$$

Another way of describing $A \cup B$ is to look at all elements in A plus all elements in B minus the duplicates, which are elements in both A and B.

$$|A \cup B| = |A| + |B| - |A \cap B|$$

This also holds for $A \cup B \cup C$ but this time we have more duplicates. Elements in $A \cap B$, $A \cap C$ and $B \cap C$ are "counted" twice, and elements in $A \cap B \cap C$ are "counted" an extra time.

$$|A \cup B \cup C| = |A| + |B| + |C| - \underbrace{|A \cap B| - |A \cap C| - |B \cap C|}_{\text{elements covered twice by } |A|+|B|+|C|} - \underbrace{|A \cap B \cap C|}_{\text{elements covered three times by } |A|+|B|+|C|}$$

Section 2.3**Exercise 12**

- a) $f(n) = n - 1$ is one-to-one since the function is strictly growing. Since every Δn increase on its domain will produce a Δy increase on its codomain, all values of n maps to a unique value
- b) $f(n) = n^2 + 1$ is one-to-one since $f : \mathbb{Z} \mapsto \mathbb{Z}$ and is strictly growing (even in $n=0$ because of \mathbb{Z}).
- c) $f(n) = n^3$ is one to one for the same reason as in b)
- d) $f(n) = \lceil n/2 \rceil$ is not one to one since every value of n maps to two values (except for $n = 0$).

Exercise 38

$$f(x) = ax + b$$

$$g(x) = cx + d$$

$$f \circ g = f(g(x)) = a(cx + d) + b = acx + ad + b$$

$$g \circ f = g(f(x)) = c(ax + b) + d = acx + cb + d$$

$$f \circ g = g \circ f \iff ad + b = cb + d$$

Exercise 42

$f : \mathbb{R} \mapsto \mathbb{R}$ given by $f(x) = x^2$ has an inverse f^{-1} given by $f^{-1}(x) = \pm\sqrt{x}$

a) $f^{-1}(\{1\}) = \pm 1$

b) $f^{-1}(\{x \mid 0 < x < 1\}) = \pm\{x \mid 0 < x < \sqrt{1}\} = \{x \mid -1 < x < 1 \underbrace{\wedge x \neq 0}_{\text{input not defined for } x=0}}\}$

c) $f^{-1}(\{x \mid x > 4\}) = \pm\{x \mid x > \sqrt{4}\} = \{x \mid -2 > x \wedge x > 2\}$

Section 2.4**Exercise 12**

- a) c
- b)

Exercise 33

- a) d
- b)

Section 2.5**Exercise 16**

- a)