

Single Variable Calculus

Otto Martinwall

February 28, 2022

Contents

1	Calculus A Complete Course	2
1.1	Limits and Continuity	3
1.1.1	Exercises 1.1	5
1.1.2	Exercises 1.2	5
2	Lectures	6

1 Calculus A Complete Course

1.1 Limits and Continuity

Theorem 1 (The Squeeze Theorem). *Suppose that $f(x) \leq g(x) \leq h(x)$ holds for all x in some open interval containing c , except possibly at $x = c$. Suppose also that*

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} h(x) = L$$

Then $\lim_{x \rightarrow c} g(x) = L$.

Proof. For this proof, the (ϵ, δ) -definition of the limit will be used.

The goal is to prove that $\lim_{x \rightarrow c} g(x) = L$, which is true if

$$\forall \epsilon > 0, \exists \delta > 0 : \forall x, (|x - a| < \delta \Rightarrow |g(x) - L| < \epsilon).$$

Since $\lim_{x \rightarrow c} f(x) = L$,

$$\forall \epsilon > 0, \exists \delta_1 > 0 : \forall x, (|x - a| < \delta_1 \Rightarrow |f(x) - L| < \epsilon) \quad (1)$$

And since $\lim_{x \rightarrow c} h(x) = L$,

$$\forall \epsilon > 0, \exists \delta_2 > 0 : \forall x, (|x - a| < \delta_2 \Rightarrow |h(x) - L| < \epsilon). \quad (2)$$

Then we have

$$f(x) \leq g(x) \leq h(x)$$

$$f(x) - L \leq g(x) - L \leq h(x) - L$$

We can choose $\delta = \min\{\delta_1, \delta_2\}$, then if $|x - a| < \delta$, and combining (1) and (2), we have

$$-\epsilon < f(x) - L \leq g(x) - L \leq h(x) - L < \epsilon$$

$$-\epsilon < g(x) - L < \epsilon$$

$$|g(x) - L| < \epsilon$$

So $\lim_{x \rightarrow c} g(x) = L$, which completes the proof.

□

1.1.1 Exercises 1.1

1.1.2 Exercises 1.2

2 Lectures