# Multi-agent learning

No-regret learning

Gerard Vreeswijk, Intelligent Software Systems, Computer Science Department, Faculty of Sciences, Utrecht University, The Netherlands.

Monday 11<sup>th</sup> May, 2020



Author: Gerard Vreeswijk. Slides last modified on May  $11^{\mathrm{th}}$ , 2020 at 17:21

■ Reinforcement
Learning. Play those
actions that were
successful in the past.

- Reinforcement
  Learning. Play those
  actions that were
  successful in the past.
- **No-regret learning**: might be considered as an extension of reinforcement learning. *Play those actions that would have been successful in the past*.

- Reinforcement
  Learning. Play those
  actions that were
  successful in the past.
- Similarities:

- Reinforcement
  Learning. Play those
  actions that were
  successful in the past.
- Similarities:
  - 1. Driven by past payoffs.

- Reinforcement
  Learning. Play those
  actions that were
  successful in the past.
- Similarities:
  - 1. Driven by past payoffs.
  - 2. Not interested in (the behaviour of) the opponent.

- Reinforcement
  Learning. Play those
  actions that were
  successful in the past.
- Similarities:
  - 1. Driven by past payoffs.
  - 2. Not interested in (the behaviour of) the opponent.
  - 3. Probabilistic.

■ Reinforcement
Learning. Play those
actions that were
successful in the past.

# Similarities:

- 1. Driven by past payoffs.
- 2. Not interested in (the behaviour of) the opponent.
- 3. Probabilistic.
- 4. Smooth adaptation.

- Reinforcement
  Learning. Play those
  actions that were
  successful in the past.
- No-regret learning: might be considered as an extension of reinforcement learning. Play those actions that would have been successful in the past.

- Similarities:
  - 1. Driven by past payoffs.
  - 2. Not interested in (the behaviour of) the opponent.
  - 3. Probabilistic.
  - 4. Smooth adaptation.
  - 5. Myopic.

- Reinforcement
  Learning. Play those
  actions that were
  successful in the past.
- Similarities:
  - 1. Driven by past payoffs.
  - 2. Not interested in (the behaviour of) the opponent.
  - 3. Probabilistic.
  - 4. Smooth adaptation.
  - 5. Myopic.

- **No-regret learning**: might be considered as an extension of reinforcement learning. *Play those actions that would have been successful in the past*.
- *Differences*:

■ Reinforcement
Learning. Play those
actions that were
successful in the past.

#### ■ Similarities:

- 1. Driven by past payoffs.
- 2. Not interested in (the behaviour of) the opponent.
- 3. Probabilistic.
- 4. Smooth adaptation.
- 5. Myopic.

■ **No-regret learning**: might be considered as an extension of reinforcement learning. *Play those actions that would have been successful in the past*.

#### ■ *Differences*:

1. Keeping counts of hypothetical actions rests on the assumption that a player is able to estimate payoffs of actions that were actually not played.

■ Reinforcement
Learning. Play those
actions that were
successful in the past.

#### ■ Similarities:

- 1. Driven by past payoffs.
- 2. Not interested in (the behaviour of) the opponent.
- 3. Probabilistic.
- 4. Smooth adaptation.
- 5. Myopic.

■ **No-regret learning**: might be considered as an extension of reinforcement learning. *Play those actions that would have been successful in the past*.

#### ■ *Differences*:

1. Keeping counts of hypothetical actions rests on the assumption that a player is able to estimate payoffs of actions that were actually not played.

(Knowledge of the payoff matrix helps, but is a stronger assumption.)

■ Reinforcement
Learning. Play those
actions that were
successful in the past.

#### ■ Similarities:

- 1. Driven by past payoffs.
- 2. Not interested in (the behaviour of) the opponent.
- 3. Probabilistic.
- 4. Smooth adaptation.
- 5. Myopic.

■ **No-regret learning**: might be considered as an extension of reinforcement learning. *Play those actions that would have been successful in the past*.

#### Differences:

- Keeping counts of hypothetical actions rests on the assumption that a player is able to estimate payoffs of actions that were actually not played.
   (Knowledge of the payoff matrix helps, but is a stronger assumption.)
- 2. It is more easy to obtain results regarding performance.

■ Reinforcement
Learning. Play those
actions that were
successful in the past.

#### ■ Similarities:

- 1. Driven by past payoffs.
- 2. Not interested in (the behaviour of) the opponent.
- 3. Probabilistic.
- 4. Smooth adaptation.
- 5. Myopic.

■ **No-regret learning**: might be considered as an extension of reinforcement learning. *Play those actions that would have been successful in the past*.

#### ■ *Differences*:

- Keeping counts of hypothetical actions rests on the assumption that a player is able to estimate payoffs of actions that were actually not played.
   (Knowledge of the payoff matrix helps, but is a stronger assumption.)
- 2. It is more easy to obtain results regarding performance. (*Correlated equilibrium*.)



1. **Probabilistic choice**. A choice of action is never completely determined by history but has a random component.

- 1. **Probabilistic choice**. A choice of action is never completely determined by history but has a random component.
  - The randomness ensures exploration.

- 1. **Probabilistic choice**. A choice of action is never completely determined by history but has a random component.
  - The randomness ensures exploration.
  - The different magnitudes of the probabilities (arisen through experience) ensures exploitation of past experience.

- 1. **Probabilistic choice**. A choice of action is never completely determined by history but has a random component.
  - The randomness ensures exploration.
  - The different magnitudes of the probabilities (arisen through experience) ensures exploitation of past experience.
- 2. **Smooth adaptation**. The strategy of play changes gradually.

- 1. **Probabilistic choice**. A choice of action is never completely determined by history but has a random component.
  - The randomness ensures exploration.
  - The different magnitudes of the probabilities (arisen through experience) ensures exploitation of past experience.
- 2. **Smooth adaptation**. The strategy of play changes gradually.
  - No-regret learning. Select a pure strategy that would have been most successful, given past play.

- 1. **Probabilistic choice**. A choice of action is never completely determined by history but has a random component.
  - The randomness ensures exploration.
  - The different magnitudes of the probabilities (arisen through experience) ensures exploitation of past experience.
- 2. **Smooth adaptation**. The strategy of play changes gradually.
  - No-regret learning. Select a pure strategy that would have been most successful, given past play.
  - Smoothed fictitious play. Give a soft-max response to the (recent) empirical frequency of opponents' actions.

- 1. **Probabilistic choice**. A choice of action is never completely determined by history but has a random component.
  - The randomness ensures exploration.
  - The different magnitudes of the probabilities (arisen through experience) ensures exploitation of past experience.
- 2. **Smooth adaptation**. The strategy of play changes gradually.
  - No-regret learning. Select a pure strategy that would have been most successful, given past play.
  - Smoothed fictitious play. Give a soft-max response to the (recent) empirical frequency of opponents' actions.
  - Hypothesis testing with smoothed best responses. Give a best response to maintained beliefs about *patterns of play*.

Three parts.

Three parts.

1. Basic concepts.

Three parts.

- 1. Basic concepts.
- 2. **Proportional regret matching**. Hart and Mas-Colell (2000).

#### Three parts.

- 1. Basic concepts.
- 2. **Proportional regret matching**. Hart and Mas-Colell (2000).
- 3.  $\epsilon$ -Greedy off-policy regret matching. Foster and Vohra (1999).

#### Three parts.

- 1. Basic concepts.
- 2. **Proportional regret matching**. Hart and Mas-Colell (2000).
- 3.  $\epsilon$ -Greedy off-policy regret matching. Foster and Vohra (1999).

This presentation almost exclusively follows the second half of Ch. 2 of (Peyton Young, 2004).

#### Three parts.

- 1. Basic concepts.
- 2. **Proportional regret matching**. Hart and Mas-Colell (2000).
- 3.  $\epsilon$ -Greedy off-policy regret matching. Foster and Vohra (1999).

This presentation almost exclusively follows the second half of Ch. 2 of (Peyton Young, 2004).

Peyton Young, H. (2004): *Strategic Learning and it Limits*, Oxford UP. Ch. 2: "Reinforcement and Regret"

#### Three parts.

- 1. Basic concepts.
- 2. **Proportional regret matching**. Hart and Mas-Colell (2000).
- 3.  $\epsilon$ -Greedy off-policy regret matching. Foster and Vohra (1999).

This presentation almost exclusively follows the second half of Ch. 2 of (Peyton Young, 2004).

Peyton Young, H. (2004): Strategic Learning and it Limits, Oxford UP. Ch. 2: "Reinforcement and Regret"

Foster, D., and Vohra, R. (1999). "Regret in the on-line decision problem". *Games and Economic Behavior*, **29**, pp. 7-36.

#### Three parts.

- 1. Basic concepts.
- 2. **Proportional regret matching**. Hart and Mas-Colell (2000).
- 3.  $\epsilon$ -Greedy off-policy regret matching. Foster and Vohra (1999).

This presentation almost exclusively follows the second half of Ch. 2 of (Peyton Young, 2004).

Peyton Young, H. (2004): Strategic Learning and it Limits, Oxford UP. Ch. 2: "Reinforcement and Regret"

Foster, D., and Vohra, R. (1999). "Regret in the on-line decision problem". *Games and Economic Behavior*, **29**, pp. 7-36.

Hart, S., and Mas-Colell, A. (2000). "A simple adaptive procedure leading to correlated equilibrium". *Econometrica*, **68**, pp. 1127-1150.

# Part I: Basic concepts

■ Suppose *A* is offered to replay the first 11 periods, under the proviso that he must play one pure strategy (i.e., action) throughout.

Suppose A is offered to replay the first 11 periods, under the proviso that he must play one pure strategy (i.e., action) throughout.

Rounds 1-11: 3
Had L played: 6 6-3 (6-3)/11Had R played: 5 5-3 (5-3)/11

Suppose A is offered to replay the first 11 periods, under the proviso that he must play one pure strategy (i.e., action) throughout.

Rounds 1-11: 3
Had L played: 6 6-3 (6-3)/11Had R played: 5 5-3 (5-3)/11

■ It is ignored that *B* likely would have played different if he knew *A* would have played different.

## No-regret: example

Suppose A is offered to replay the first 11 periods, under the proviso that he must play one pure strategy (i.e., action) throughout.

Rounds 1-11: 3
Had 
$$L$$
 played:  $6 6-3 (6-3)/11$ 
Had  $R$  played:  $5 5-3 (5-3)/11$ 

■ It is ignored that *B* likely would have played different if he knew *A* would have played different.

So no-regret does not take the interactive nature of play into account.

Author: Gerard Vreeswijk. Slides last modified on May  $11^{\mathrm{th}}$ , 2020 at 17:21

■ The average payoff up to and including round t is

$$\bar{u}^t =_{Def} \frac{1}{t} \sum_{s=1}^t u(x^s, y^s).$$

The average payoff up to and including round t is

$$\bar{u}^t =_{Def} \frac{1}{t} \sum_{s=1}^t u(x^s, y^s).$$

For each action x, the hypothetical average payoff for playing x is

$$\bar{h}_{x}^{t} =_{Def} \frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}).$$

The average payoff up to and including round t is

$$\bar{u}^t =_{Def} \frac{1}{t} \sum_{s=1}^t u(x^s, y^s).$$

For each action x, the hypothetical average payoff for playing x is

$$\bar{h}_x^t =_{Def} \frac{1}{t} \sum_{s=1}^t u(x, y^s).$$

For each action x, the average regret from not having played x is

$$\bar{r}_x^t =_{Def} \bar{h}_x^t - \bar{u}^t.$$

The average payoff up to and including round t is

$$\bar{u}^t =_{Def} \frac{1}{t} \sum_{s=1}^t u(x^s, y^s).$$

For each action x, the hypothetical average payoff for playing x is

$$\bar{h}_{x}^{t} =_{Def} \frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}).$$

For each action x, the average regret from not having played x is

$$\bar{r}_{x}^{t} =_{Def} \bar{h}_{x}^{t} - \bar{u}^{t}.$$

Average regret may be represented as a vector

$$\bar{r}^t =_{Def} (\bar{r}_1^t, \dots, \bar{r}_k^t)^T.$$

The average payoff up to and including round t is

$$\bar{u}^t =_{Def} \frac{1}{t} \sum_{s=1}^t u(x^s, y^s).$$

For each action x, the hypothetical average payoff for playing x is

$$\bar{h}_{x}^{t} =_{Def} \frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}).$$

For each action x, the average regret from not having played x is

$$\bar{r}_{x}^{t} =_{Def} \bar{h}_{x}^{t} - \bar{u}^{t}.$$

 Average regret may be represented as a vector

$$\bar{r}^t =_{Def} (\bar{r}_1^t, \dots, \bar{r}_k^t)^T.$$

■ A given realisation of play

$$\omega = (x_1, y_1), \ldots, (x_t, y_t), \ldots$$

is said to have no regret if, for all actions x,

The average payoff up to and including round t is

$$\bar{u}^t =_{Def} \frac{1}{t} \sum_{s=1}^t u(x^s, y^s).$$

■ For each action *x*, the hypothetical average payoff for playing *x* is

$$\bar{h}_{x}^{t} =_{Def} \frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}).$$

For each action x, the average regret from not having played x is

$$\bar{r}_{x}^{t} =_{Def} \bar{h}_{x}^{t} - \bar{u}^{t}.$$

 Average regret may be represented as a vector

$$\bar{r}^t =_{Def} (\bar{r}_1^t, \dots, \bar{r}_k^t)^T.$$

■ A given realisation of play

$$\omega = (x_1, y_1), \ldots, (x_t, y_t), \ldots$$

is said to have no regret if, for all actions x,

$$\limsup_{t\to\infty} \bar{r}_x^t(\omega) \le 0$$

The average payoff up to and including round t is

$$\bar{u}^t =_{Def} \frac{1}{t} \sum_{s=1}^t u(x^s, y^s).$$

For each action x, the hypothetical average payoff for playing x is

$$\bar{h}_{x}^{t} =_{Def} \frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}).$$

For each action x, the average regret from not having played x is

$$\bar{r}_{x}^{t} =_{Def} \bar{h}_{x}^{t} - \bar{u}^{t}.$$

 Average regret may be represented as a vector

$$\bar{r}^t =_{Def} (\bar{r}_1^t, \dots, \bar{r}_k^t)^T.$$

■ A given realisation of play

$$\omega = (x_1, y_1), \ldots, (x_t, y_t), \ldots$$

is said to have no regret if, for all actions x,

$$\limsup_{t\to\infty} \bar{r}_{x}^{t}(\omega) \leq 0$$

i.e.  $\lim_{T\to\infty} \sup\{ \bar{r}_x^t(\omega) \mid T \le t \} \le 0$ 

The average payoff up to and including round t is

$$\bar{u}^t =_{Def} \frac{1}{t} \sum_{s=1}^t u(x^s, y^s).$$

For each action x, the hypothetical average payoff for playing x is

$$\bar{h}_{x}^{t} =_{Def} \frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}).$$

For each action x, the average regret from not having played x is

$$\bar{r}_{x}^{t} =_{Def} \bar{h}_{x}^{t} - \bar{u}^{t}.$$

 Average regret may be represented as a vector

$$\bar{r}^t =_{Def} (\bar{r}_1^t, \dots, \bar{r}_k^t)^T.$$

■ A given realisation of play

$$\omega = (x_1, y_1), \ldots, (x_t, y_t), \ldots$$

is said to have no regret if, for all actions x,

$$\limsup_{t\to\infty} \bar{r}_{x}^{t}(\omega) \leq 0$$

i.e.  $\lim_{T\to\infty} \sup\{\bar{r}_x^t(\omega) \mid T \leq t\} \leq 0$ 

$$\Leftrightarrow \lim_{t\to\infty} [\bar{r}_x^t(\omega)]_+ = 0.$$

# Part II: proportional regret matching

A strategy  $g: H \to \Delta(X)$  is said to have no regret if almost all of its realisations of play have no regret.

A strategy  $g: H \to \Delta(X)$  is said to have no regret if almost all of its realisations of play have no regret. The objective is to formulate a strategy without regret.

A strategy  $g: H \to \Delta(X)$  is said to have no regret if almost all of its realisations of play have no regret. The objective is to formulate a strategy without regret.

One candidate strategy is proposed by Hart and Mas-Colell (2000):

$$q_x^{t+1} =_{Def} \frac{[\bar{r}_x^t]_+}{\sum_{x' \in X} [\bar{r}_{x'}^t]_+}$$

where  $[z]_{+} =_{Def} z \ge 0 ? z : 0$ .

A strategy  $g: H \to \Delta(X)$  is said to have no regret if almost all of its realisations of play have no regret. The objective is to formulate a strategy without regret.

One candidate strategy is proposed by Hart and Mas-Colell (2000):

$$q_x^{t+1} =_{Def} \frac{[\bar{r}_x^t]_+}{\sum_{x' \in X} [\bar{r}_{x'}^t]_+}$$

where  $[z]_+ =_{Def} z \ge 0$  ? z : 0. This rule is called proportional regret matching, or regret matching (RM for short).

A strategy  $g: H \to \Delta(X)$  is said to have no regret if almost all of its realisations of play have no regret. The objective is to formulate a strategy without regret.

One candidate strategy is proposed by Hart and Mas-Colell (2000):

$$q_x^{t+1} =_{Def} \frac{[\bar{r}_x^t]_+}{\sum_{x' \in X} [\bar{r}_{x'}^t]_+}$$

where  $[z]_+ =_{Def} z \ge 0$  ? z : 0. This rule is called proportional regret matching, or regret matching (RM for short). Indeed:

**Theorem** (Hart & Mas-Colell, 2000). *In a finite game, regret matching yields no regret a.s.* 

A strategy  $g: H \to \Delta(X)$  is said to have no regret if almost all of its realisations of play have no regret. The objective is to formulate a strategy without regret.

One candidate strategy is proposed by Hart and Mas-Colell (2000):

$$q_x^{t+1} =_{Def} \frac{[\bar{r}_x^t]_+}{\sum_{x' \in X} [\bar{r}_{x'}^t]_+}$$

where  $[z]_+ =_{Def} z \ge 0$  ? z : 0. This rule is called proportional regret matching, or regret matching (RM for short). Indeed:

**Theorem** (Hart & Mas-Colell, 2000). *In a finite game, regret matching yields no regret a.s.* 

Hart & Mas-Colell (2000). "A simple adaptive procedure leading to correlated equilibrium". *Econometrica*, **68**, pp. 1127-1150.

# Regret matching differs from reinforcement learning

0 0 0 1 1 0 0 0 1 0 0

A L R L L R R L R R R R ?

B R L R L R L R L R L R .

## Regret matching differs from reinforcement learning

	0	0	0	1	1	0	0	0	1	0	0	
$\boldsymbol{A}$	L	R	L	L	R	R	L	R	R	R	R	?
В	R	L	R	L	R	L	R	L	R	L	L	?

## Proportional regret matching:

	Payoff	Average regret	Regret matching
Rounds 1-11:	3		
Had <i>L</i> been played:	6	(6-3)/11	3/5
Had <i>R</i> been played:	5	(5-3)/11	2/5

# Regret matching differs from reinforcement learning

	0	0	0	1	1	0	0	0	1	0	0	
A	L	R	L	L	R	R	L	R	R	R	R	?
B	R	L	R	L	R	L	R	L	R	L	L	?

## Proportional regret matching:


	Payoff	Average regret	Regret matching
Rounds 1-11:	3		
Had $L$ been played:	6	(6-3)/11	3/5
Had <i>R</i> been played:	5	(5-3)/11	2/5

### Cumulative payoff matching:

	Accumulated payoff	Mixed strategy
Action <i>L</i> :	1	1/3
Action $R$ :	2	2/3

## Regret matching in a 5-person 5-action game

Payoff matrix uninformative. Omitted ...




Netlogo simulation of regret matching in a 5-person 5-action game.

## Regret matching in Shapley's game


	R	Y	В
R	(1,0)	(0,0)	(0,1)
Y	(0,1)	(1,0)	(0,0)
В	(0,0)	(0,1)	(1,0)

Column is "fashion leader", row is "fashion follower". Column wants to wear a different color than row.


## Regret matching in Shapley's game



Column is "fashion leader", row is "fashion follower". Column wants to wear a different color than row.



Netlogo simulation of regret matching in Shapley's game.



Quantities:


Author: Gerard Vreeswijk. Slides last modified on May 11<sup>th</sup>, 2020 at 17:21

Quantities:

 $r_x^t =_{Def}$  total regret for not playing x, up to and including t

### Quantities:

 $r_x^t =_{Def}$  total regret for not playing x, up to and including t  $\bar{r}_x^t =_{Def}$  average regret for not playing x, up to and including t



### Quantities:

 $r_x^t =_{Def}$  total regret for not playing x, up to and including t  $\bar{r}_x^t =_{Def}$  average regret for not playing x, up to and including t  $[\bar{r}_x^t]_+ =_{Def}$  positive average regret for not playing x

### Quantities:

 $r_x^t =_{Def}$  total regret for not playing x, up to and including t  $\bar{r}_x^t =_{Def}$  average regret for not playing x, up to and including t  $[\bar{r}_x^t]_+ =_{Def}$  positive average regret for not playing x  $\Delta r_x^t =_{Def}$  incremental regret for not playing  $x : r_x^t - r_x^{t-1}$ 

### Quantities:

```
r_x^t =_{Def} total regret for not playing x, up to and including t
\bar{r}_x^t =_{Def} average regret for not playing x, up to and including t
[\bar{r}_x^t]_+ =_{Def} positive average regret for not playing x
\Delta r_x^t =_{Def} \text{ incremental regret for not playing } x : r_x^t - r_x^{t-1}
E[\Delta r_x^t] = \text{ expected incremental regret for not playing } x
```

### Quantities:

```
r_x^t =_{Def} total regret for not playing x, up to and including t
\bar{r}_x^t =_{Def} average regret for not playing x, up to and including t
[\bar{r}_x^t]_+ =_{Def} positive average regret for not playing x
\Delta r_x^t =_{Def} \text{ incremental regret for not playing } x : r_x^t - r_x^{t-1}
E[\Delta r_x^t] = \text{ expected incremental regret for not playing } x
```

Vector versions:  $r^t$ ,  $\bar{r}^t$ ,  $[\bar{r}^t]_+$ , ...,  $E[r^t]$ ,  $E[\Delta r^t]$ .



### Quantities:

```
r_x^t =_{Def} total regret for not playing x, up to and including t
\bar{r}_x^t =_{Def} average regret for not playing x, up to and including t
[\bar{r}_x^t]_+ =_{Def} positive average regret for not playing x
\Delta r_x^t =_{Def} \text{ incremental regret for not playing } x : r_x^t - r_x^{t-1}
E[\Delta r_x^t] = \text{ expected incremental regret for not playing } x
```

Vector versions:  $r^t$ ,  $\bar{r}^t$ ,  $[\bar{r}^t]_+$ , ...,  $E[r^t]$ ,  $E[\Delta r^t]$ .



### Quantities:

 $r_x^t =_{Def}$  total regret for not playing x, up to and including t  $\bar{r}_x^t =_{Def}$  average regret for not playing x, up to and including t  $[\bar{r}_x^t]_+ =_{Def}$  positive average regret for not playing x  $\Delta r_x^t =_{Def} \text{ incremental regret for not playing } x : r_x^t - r_x^{t-1}$   $E[\Delta r_x^t] = \text{ expected incremental regret for not playing } x$ 

Vector versions:  $r^t$ ,  $\bar{r}^t$ ,  $[\bar{r}^t]_+$ , ...,  $E[r^t]$ ,  $E[\Delta r^t]$ .

### Objective:

$$\lim_{t\to\infty} \left[\bar{r}^t\right]_+ = 0 \text{ a.s.}$$

### Quantities:

 $r_x^t =_{Def}$  total regret for not playing x, up to and including t  $\bar{r}_x^t =_{Def}$  average regret for not playing x, up to and including t  $[\bar{r}_x^t]_+ =_{Def}$  positive average regret for not playing x  $\Delta r_x^t =_{Def} \text{ incremental regret for not playing } x : r_x^t - r_x^{t-1}$   $E[\Delta r_x^t] = \text{ expected incremental regret for not playing } x$ 

Vector versions:  $r^t$ ,  $\bar{r}^t$ ,  $[\bar{r}^t]_+$ , ...,  $E[r^t]$ ,  $E[\Delta r^t]$ .

## Objective:

$$\lim_{t\to\infty} \left[\bar{r}^t\right]_+ = 0 \text{ a.s.}$$

i.e., the regret vector must approach the negative orthant with probability one.

## Incremental regret and expected incremental regret

Suppose there are only two actions, "1" and "2," say.

## Incremental regret and expected incremental regret

Suppose there are only two actions, "1" and "2," say.

1. If 1 is executed at t + 1 then

- 1. If 1 is executed at t + 1 then
  - $ightharpoonup r_1^{t+1}$  will not change.

- 1. If 1 is executed at t + 1 then
  - $ightharpoonup r_1^{t+1}$  will not change.
  - $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1})$ .

- 1. If 1 is executed at t + 1 then
- 2. If 2 is executed at t + 1 then

- $\blacksquare$   $r_1^{t+1}$  will not change.
- $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1})$ .

- 1. If 1 is executed at t + 1 then
  - $\blacksquare$   $r_1^{t+1}$  will not change.
  - $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1})$ .

- 2. If 2 is executed at t + 1 then
  - $r_1^{t+1}$  changes with  $u(1, y^{t+1}) u(2, y^{t+1})$ .

- 1. If 1 is executed at t + 1 then
  - $\blacksquare$   $r_1^{t+1}$  will not change.
  - $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1})$ .

- 2. If 2 is executed at t + 1 then
  - $r_1^{t+1}$  changes with  $u(1, y^{t+1}) u(2, y^{t+1})$ .
  - $\blacksquare$   $r_2^{t+1}$  will not change.

Suppose there are only two actions, "1" and "2," say.

- 1. If 1 is executed at t + 1 then
  - $\blacksquare$   $r_1^{t+1}$  will not change.
  - $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1})$ .

- 2. If 2 is executed at t + 1 then
  - $r_1^{t+1}$  changes with  $u(1, y^{t+1}) u(2, y^{t+1})$ .
  - $\blacksquare$   $r_2^{t+1}$  will not change.

If  $\alpha^{t+1} =_{Def} u(1, y^{t+1}) - u(2, y^{t+1})$  then incremental regret will be either  $(0, -\alpha^{t+1})$ 

Suppose there are only two actions, "1" and "2," say.

- 1. If 1 is executed at t + 1 then
  - $\blacksquare$   $r_1^{t+1}$  will not change.
  - $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1})$ .

- 2. If 2 is executed at t + 1 then
  - $r_1^{t+1}$  changes with  $u(1, y^{t+1}) u(2, y^{t+1})$ .
  - $\blacksquare$   $r_2^{t+1}$  will not change.

If  $\alpha^{t+1} =_{Def} u(1, y^{t+1}) - u(2, y^{t+1})$  then incremental regret will be either  $(0, -\alpha^{t+1})$  or  $(\alpha^{t+1}, 0)$ .

Suppose there are only two actions, "1" and "2," say.

- 1. If 1 is executed at t + 1 then
  - $\blacksquare$   $r_1^{t+1}$  will not change.
  - $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1}).$

- 2. If 2 is executed at t + 1 then
  - $r_1^{t+1}$  changes with  $u(1, y^{t+1}) u(2, y^{t+1}).$
  - $\blacksquare$   $r_2^{t+1}$  will not change.

If  $\alpha^{t+1} =_{Def} u(1, y^{t+1}) - u(2, y^{t+1})$  then incremental regret will be either  $(0, -\alpha^{t+1})$  or  $(\alpha^{t+1}, 0)$ .

Suppose in round t + 1 a mixed strategy  $q^{t+1} = (q_1^{t+1}, q_2^{t+1})$  is played.

Suppose there are only two actions, "1" and "2," say.

- 1. If 1 is executed at t + 1 then
  - $\blacksquare$   $r_1^{t+1}$  will not change.
  - $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1})$ .

- 2. If 2 is executed at t + 1 then
  - $r_1^{t+1}$  changes with  $u(1, y^{t+1}) u(2, y^{t+1})$ .
  - $ightharpoonup r_2^{t+1}$  will not change.

If  $\alpha^{t+1} =_{Def} u(1, y^{t+1}) - u(2, y^{t+1})$  then incremental regret will be either  $(0, -\alpha^{t+1})$  or  $(\alpha^{t+1}, 0)$ .

$$E[\Delta r^{t+1}] = ($$

Suppose there are only two actions, "1" and "2," say.

- 1. If 1 is executed at t + 1 then
  - $\blacksquare$   $r_1^{t+1}$  will not change.
  - $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1}).$

- 2. If 2 is executed at t + 1 then
  - $r_1^{t+1}$  changes with  $u(1, y^{t+1}) u(2, y^{t+1}).$
  - $ightharpoonup r_2^{t+1}$  will not change.

If  $\alpha^{t+1} =_{Def} u(1, y^{t+1}) - u(2, y^{t+1})$  then incremental regret will be either  $(0, -\alpha^{t+1})$  or  $(\alpha^{t+1}, 0)$ .

$$E[\Delta r^{t+1}] = (q_1^{t+1} \cdot 0, q_2^{t+1}) \cdot q_2^{t+1} \cdot q_2^{t+1$$

Suppose there are only two actions, "1" and "2," say.

- 1. If 1 is executed at t + 1 then
  - $\blacksquare$   $r_1^{t+1}$  will not change.
  - $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1})$ .

- 2. If 2 is executed at t + 1 then
  - $r_1^{t+1}$  changes with  $u(1, y^{t+1}) u(2, y^{t+1}).$
  - $\blacksquare$   $r_2^{t+1}$  will not change.

If  $\alpha^{t+1} =_{Def} u(1, y^{t+1}) - u(2, y^{t+1})$  then incremental regret will be either  $(0, -\alpha^{t+1})$  or  $(\alpha^{t+1}, 0)$ .

$$E[\Delta r^{t+1}] = (q_1^{t+1} \cdot 0 + q_2^{t+1} \cdot \alpha^{t+1}),$$

Suppose there are only two actions, "1" and "2," say.

- 1. If 1 is executed at t + 1 then
  - $\blacksquare$   $r_1^{t+1}$  will not change.
  - $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1})$ .

- 2. If 2 is executed at t + 1 then
  - $r_1^{t+1}$  changes with  $u(1, y^{t+1}) u(2, y^{t+1})$ .
  - $\blacksquare$   $r_2^{t+1}$  will not change.

If  $\alpha^{t+1} =_{Def} u(1, y^{t+1}) - u(2, y^{t+1})$  then incremental regret will be either  $(0, -\alpha^{t+1})$  or  $(\alpha^{t+1}, 0)$ .

$$E[\Delta r^{t+1}] = (q_1^{t+1} \cdot 0 + q_2^{t+1} \cdot \alpha^{t+1}, q_1^{t+1} \cdot -\alpha^{t+1})$$

Suppose there are only two actions, "1" and "2," say.

- 1. If 1 is executed at t + 1 then
  - $\blacksquare$   $r_1^{t+1}$  will not change.
  - $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1}).$

- 2. If 2 is executed at t + 1 then
  - $r_1^{t+1}$  changes with  $u(1, y^{t+1}) u(2, y^{t+1})$ .
  - $\blacksquare$   $r_2^{t+1}$  will not change.

If  $\alpha^{t+1} =_{Def} u(1, y^{t+1}) - u(2, y^{t+1})$  then incremental regret will be either  $(0, -\alpha^{t+1})$  or  $(\alpha^{t+1}, 0)$ .

$$E[\Delta r^{t+1}] = (q_1^{t+1} \cdot 0 + q_2^{t+1} \cdot \alpha^{t+1}, q_1^{t+1} \cdot -\alpha^{t+1} + q_2^{t+1} \cdot 0)$$

Suppose there are only two actions, "1" and "2," say.

- 1. If 1 is executed at t + 1 then
  - $\blacksquare$   $r_1^{t+1}$  will not change.
  - $r_2^{t+1}$  changes with  $u(2, y^{t+1}) u(1, y^{t+1}).$

- 2. If 2 is executed at t + 1 then
  - $r_1^{t+1}$  changes with  $u(1, y^{t+1}) u(2, y^{t+1})$ .
  - $\blacksquare$   $r_2^{t+1}$  will not change.

If  $\alpha^{t+1} =_{Def} u(1, y^{t+1}) - u(2, y^{t+1})$  then incremental regret will be either  $(0, -\alpha^{t+1})$  or  $(\alpha^{t+1}, 0)$ .

$$E[\Delta r^{t+1}] = (q_1^{t+1} \cdot 0 + q_2^{t+1} \cdot \alpha^{t+1}, q_1^{t+1} \cdot -\alpha^{t+1} + q_2^{t+1} \cdot 0)$$

$$= \alpha^{t+1}(q_2^{t+1}, -q_1^{t+1}).$$

# Why does regret matching work?

Take 
$$q_1^t = q_2^t = 1/2$$
 for all  $t$ . Then

$$E[\bar{r}_1^t + \bar{r}_2^t] = E[\frac{r_1^{t-1} + \Delta r_1^t}{t} + \frac{r_2^{t-1} + \Delta r_2^t}{t}]$$

$$E[\bar{r}_1^t + \bar{r}_2^t] = E\left[\frac{r_1^{t-1} + \Delta r_1^t}{t} + \frac{r_2^{t-1} + \Delta r_2^t}{t}\right]$$

$$= E\left[\frac{r_1^{t-1} - \alpha^t/2}{t} + \frac{r_2^{t-1} + \alpha^t/2}{t}\right]$$

$$E[\bar{r}_1^t + \bar{r}_2^t] = E\left[\frac{r_1^{t-1} + \Delta r_1^t}{t} + \frac{r_2^{t-1} + \Delta r_2^t}{t}\right]$$

$$= E\left[\frac{r_1^{t-1} - \alpha^t/2}{t} + \frac{r_2^{t-1} + \alpha^t/2}{t}\right]$$

$$= E\left[\bar{r}_1^{t-1} + \bar{r}_2^{t-1}\right]$$

$$E[\bar{r}_1^t + \bar{r}_2^t] = E[\frac{r_1^{t-1} + \Delta r_1^t}{t} + \frac{r_2^{t-1} + \Delta r_2^t}{t}]$$

$$= E[\frac{r_1^{t-1} - \alpha^t/2}{t} + \frac{r_2^{t-1} + \alpha^t/2}{t}]$$

$$= E[\bar{r}_1^{t-1} + \bar{r}_2^{t-1}]$$

$$= \bar{r}_1^{t-1} + \bar{r}_2^{t-1}$$

Take  $q_1^t = q_2^t = 1/2$  for all t. Then

$$E[\bar{r}_1^t + \bar{r}_2^t] = E[\frac{r_1^{t-1} + \Delta r_1^t}{t} + \frac{r_2^{t-1} + \Delta r_2^t}{t}]$$

$$= E[\frac{r_1^{t-1} - \alpha^t/2}{t} + \frac{r_2^{t-1} + \alpha^t/2}{t}]$$

$$= E[\bar{r}_1^{t-1} + \bar{r}_2^{t-1}]$$

$$= \bar{r}_1^{t-1} + \bar{r}_2^{t-1}$$

Inductively then

$$E[\bar{r}_1^t + \bar{r}_2^t] = 0,$$

Take  $q_1^t = q_2^t = 1/2$  for all t. Then

$$E[\bar{r}_1^t + \bar{r}_2^t] = E\left[\frac{r_1^{t-1} + \Delta r_1^t}{t} + \frac{r_2^{t-1} + \Delta r_2^t}{t}\right]$$

$$= E\left[\frac{r_1^{t-1} - \alpha^t/2}{t} + \frac{r_2^{t-1} + \alpha^t/2}{t}\right]$$

$$= E\left[\bar{r}_1^{t-1} + \bar{r}_2^{t-1}\right]$$

$$= \bar{r}_1^{t-1} + \bar{r}_2^{t-1}$$

Inductively then

$$E[\bar{r}_1^t + \bar{r}_2^t] = 0,$$

so that  $\lim_{t\to\infty} \bar{r}_1^t + \bar{r}_2^t = 0$  with probability one.

Take  $q_1^t = q_2^t = 1/2$  for all t. Then

$$E[\bar{r}_1^t + \bar{r}_2^t] = E[\frac{r_1^{t-1} + \Delta r_1^t}{t} + \frac{r_2^{t-1} + \Delta r_2^t}{t}]$$

$$= E[\frac{r_1^{t-1} - \alpha^t/2}{t} + \frac{r_2^{t-1} + \alpha^t/2}{t}]$$

$$= E[\bar{r}_1^{t-1} + \bar{r}_2^{t-1}]$$

$$= \bar{r}_1^{t-1} + \bar{r}_2^{t-1}$$

Inductively then

$$E[\bar{r}_1^t + \bar{r}_2^t] = 0,$$

so that  $\lim_{t\to\infty} \bar{r}_1^t + \bar{r}_2^t = 0$  with probability one.

However, the two terms may neutralise each other.

Author: Gerard Vreeswijk. Slides last modified on May  $11^{\mathrm{th}}$ , 2020 at 17:21

Each round t, choose an action that would have minimised regret in the previous round.

- Each round t, choose an action that would have minimised regret in the previous round.
- However: Matching Pennies.

- Each round t, choose an action that would have minimised regret in the previous round.
- However: Matching Pennies.



- $\blacksquare$  Each round t, choose an action that would have minimised regret in the previous round.
- However: Matching Pennies.



• Switch actions if regret in previous round; else stay.

- Each round t, choose an action that would have minimised regret in the previous round.
- However: Matching Pennies.



• Switch actions if regret in previous round; else stay.

 Won't work: suppose you meet an opponent who happens to switch every round as well . . .

- Each round t, choose an action that would have minimised regret in the previous round.
- However: Matching Pennies.



• Switch actions if regret in previous round; else stay.

- Won't work: suppose you meet an opponent who happens to switch every round as well . . .
- Won't work in general: your corrections may by coincidence be out of phase with the path of play of your opponent. Peyton Young:

"Recall that no-regret must hold even when Nature is malevolent." (p. 26)

The objective is to find a (mixed) strategy  $g: H \to \Delta(\{1,2\})$  such that

$$E[\bar{r}^{t+1} \mid r^t, \dots, r^1] < \bar{r}^t \tag{1}$$

The objective is to find a (mixed) strategy  $g: H \to \Delta(\{1,2\})$  such that

$$E[\bar{r}^{t+1} \mid r^t, \dots, r^1] < \bar{r}^t \tag{1}$$

because then Blackwell's approachability theorem can be applied to conclude  $\lim_{t\to\infty} [\bar{r}^t]_+ = 0$ .

The objective is to find a (mixed) strategy  $g: H \to \Delta(\{1,2\})$  such that

$$E[\bar{r}^{t+1} \mid r^t, \dots, r^1] < \bar{r}^t \tag{1}$$

because then Blackwell's approachability theorem can be applied to conclude  $\lim_{t\to\infty} [\bar{r}^t]_+ = 0$ . Since  $\Delta E[r^{t+1}]$  is known, we have

$$E[\bar{r}^{t+1} \mid r^t, \dots, r^1] = E[\frac{r^t + \Delta r^{t+1}}{t+1} \mid r^t, \dots, r^1]$$

The objective is to find a (mixed) strategy  $g: H \to \Delta(\{1,2\})$  such that

$$E[\bar{r}^{t+1} \mid r^t, \dots, r^1] < \bar{r}^t \tag{1}$$

because then Blackwell's approachability theorem can be applied to conclude  $\lim_{t\to\infty} [\bar{r}^t]_+ = 0$ . Since  $\Delta E[r^{t+1}]$  is known, we have

$$E[\bar{r}^{t+1} \mid r^t, \dots, r^1] = E[\frac{r^t + \Delta r^{t+1}}{t+1} \mid r^t, \dots, r^1]$$

$$= \frac{t}{t+1} E[\frac{r^t}{t} \mid r^t, \dots, r^1] + \frac{1}{t+1} E[\Delta r^{t+1} \mid r^t, \dots, r^1]$$

The objective is to find a (mixed) strategy  $g: H \to \Delta(\{1,2\})$  such that

$$E[\bar{r}^{t+1} \mid r^t, \dots, r^1] < \bar{r}^t \tag{1}$$

because then Blackwell's approachability theorem can be applied to conclude  $\lim_{t\to\infty} [\bar{r}^t]_+ = 0$ . Since  $\Delta E[r^{t+1}]$  is known, we have

$$E[\bar{r}^{t+1} \mid r^t, \dots, r^1] = E[\frac{r^t + \Delta r^{t+1}}{t+1} \mid r^t, \dots, r^1]$$

$$= \frac{t}{t+1} E[\frac{r^t}{t} \mid r^t, \dots, r^1] + \frac{1}{t+1} E[\Delta r^{t+1} \mid r^t, \dots, r^1]$$

$$= \frac{t}{t+1} \bar{r}^t + \frac{1}{t+1} E[\Delta r^{t+1} \mid r^t, \dots, r^1]$$

#### Decrease of expected regret

The objective is to find a (mixed) strategy  $g: H \to \Delta(\{1,2\})$  such that

$$E[\bar{r}^{t+1} \mid r^t, \dots, r^1] < \bar{r}^t \tag{1}$$

because then Blackwell's approachability theorem can be applied to conclude  $\lim_{t\to\infty} [\bar{r}^t]_+ = 0$ . Since  $\Delta E[r^{t+1}]$  is known, we have

$$E[\bar{r}^{t+1} \mid r^t, \dots, r^1] = E[\frac{r^t + \Delta r^{t+1}}{t+1} \mid r^t, \dots, r^1]$$

$$= \frac{t}{t+1} E[\frac{r^t}{t} \mid r^t, \dots, r^1] + \frac{1}{t+1} E[\Delta r^{t+1} \mid r^t, \dots, r^1]$$

$$= \frac{t}{t+1} \bar{r}^t + \frac{1}{t+1} E[\Delta r^{t+1} \mid r^t, \dots, r^1]$$

$$= \frac{t}{t+1} \bar{r}^t + \frac{1}{t+1} (-\alpha^{t+1} q_2^{t+1}, \alpha^{t+1} q_1^{t+1})$$

#### Decrease of expected regret

The objective is to find a (mixed) strategy  $g: H \to \Delta(\{1,2\})$  such that

$$E[\bar{r}^{t+1} \mid r^t, \dots, r^1] < \bar{r}^t \tag{1}$$

because then Blackwell's approachability theorem can be applied to conclude  $\lim_{t\to\infty} [\bar{r}^t]_+ = 0$ . Since  $\Delta E[r^{t+1}]$  is known, we have

$$E[\bar{r}^{t+1} \mid r^t, \dots, r^1] = E[\frac{r^t + \Delta r^{t+1}}{t+1} \mid r^t, \dots, r^1]$$

$$= \frac{t}{t+1} E[\frac{r^t}{t} \mid r^t, \dots, r^1] + \frac{1}{t+1} E[\Delta r^{t+1} \mid r^t, \dots, r^1]$$

$$= \frac{t}{t+1} \bar{r}^t + \frac{1}{t+1} E[\Delta r^{t+1} \mid r^t, \dots, r^1]$$

$$= \frac{t}{t+1} \bar{r}^t + \frac{1}{t+1} (-\alpha^{t+1} q_2^{t+1}, \alpha^{t+1} q_1^{t+1})$$

So, the objective is to find a strategy such that  $\alpha^{t+1}(-q_2^{t+1}, q_1^{t+1}) < \bar{r}^t$ .

Author: Gerard Vreeswijk. Slides last modified on May  $11^{\mathrm{th}}$ , 2020 at 17:21

Recall: our objective is  $[\bar{r}^t]_+ \to 0$ .

- Recall: our objective is  $[\bar{r}^t]_+ \to 0$ .
- To this end, choose  $q^{t+1}$  such that

$$E[\Delta r^{t+1}] \perp [\bar{r}^t]_+$$

- Recall: our objective is  $[\bar{r}^t]_+ \to 0$ .
- To this end, choose  $q^{t+1}$  such that

$$E[\Delta r^{t+1}] \perp [\bar{r}^t]_+$$

So:

$$E[\Delta r^{t+1}] \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow (-\alpha^{t+1}q_2^{t+1}, \alpha^{t+1}q_1^{t+1}) \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow \alpha^{t+1}(q_1^{t+1}[\bar{r}_2^t]_+ - q_2^{t+1}[\bar{r}_1^t]_+) = 0$$

$$\Leftrightarrow q_1^{t+1} : q_2^{t+1} = [\bar{r}_1^t]_+ : [\bar{r}_2^t]_+.$$

- Recall: our objective is  $[\bar{r}^t]_+ \to 0$ .
- To this end, choose  $q^{t+1}$  such that

$$E[\Delta r^{t+1}] \perp [\bar{r}^t]_+$$

So:

$$E[\Delta r^{t+1}] \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow (-\alpha^{t+1}q_2^{t+1}, \alpha^{t+1}q_1^{t+1}) \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow \alpha^{t+1}(q_1^{t+1}[\bar{r}_2^t]_+ - q_2^{t+1}[\bar{r}_1^t]_+) = 0$$

$$\Leftrightarrow q_1^{t+1} : q_2^{t+1} = [\bar{r}_1^t]_+ : [\bar{r}_2^t]_+.$$

The last equation amounts to proportional regret matching.

- Recall: our objective is  $[\bar{r}^t]_+ \to 0$ .
- To this end, choose  $q^{t+1}$  such that

$$E[\Delta r^{t+1}] \perp [\bar{r}^t]_+$$

So:

$$E[\Delta r^{t+1}] \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow (-\alpha^{t+1}q_2^{t+1}, \alpha^{t+1}q_1^{t+1}) \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow \alpha^{t+1}(q_1^{t+1}[\bar{r}_2^t]_+ - q_2^{t+1}[\bar{r}_1^t]_+) = 0$$

$$\Leftrightarrow q_1^{t+1} : q_2^{t+1} = [\bar{r}_1^t]_+ : [\bar{r}_2^t]_+.$$

The last equation amounts to proportional regret matching.

- Recall: our objective is  $[\bar{r}^t]_+ \to 0$ .
- To this end, choose  $q^{t+1}$  such that

$$E[\Delta r^{t+1}] \perp [\bar{r}^t]_+$$

So:

$$E[\Delta r^{t+1}] \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow (-\alpha^{t+1}q_2^{t+1}, \alpha^{t+1}q_1^{t+1}) \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow \alpha^{t+1}(q_1^{t+1}[\bar{r}_2^t]_+ - q_2^{t+1}[\bar{r}_1^t]_+) = 0$$

$$\Leftrightarrow q_1^{t+1} : q_2^{t+1} = [\bar{r}_1^t]_+ : [\bar{r}_2^t]_+.$$

The last equation amounts to proportional regret matching.

(Notice that  $\alpha^{t+1}$  has left the stage.)

Boundary cases are obvious and can be treated as follows:

- Recall: our objective is  $[\bar{r}^t]_+ \to 0$ .
- To this end, choose  $q^{t+1}$  such that

$$E[\Delta r^{t+1}] \perp [\bar{r}^t]_+$$

So:

$$E[\Delta r^{t+1}] \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow (-\alpha^{t+1}q_2^{t+1}, \alpha^{t+1}q_1^{t+1}) \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow \alpha^{t+1}(q_1^{t+1}[\bar{r}_2^t]_+ - q_2^{t+1}[\bar{r}_1^t]_+) = 0$$

$$\Leftrightarrow q_1^{t+1} : q_2^{t+1} = [\bar{r}_1^t]_+ : [\bar{r}_2^t]_+.$$

The last equation amounts to proportional regret matching.

- Boundary cases are obvious and can be treated as follows:
  - If  $\bar{r}_1^t \le 0$  and  $\bar{r}_2^t > 0$ , then let  $q^{t+1} =_{Def} (0,1)$ .

- Recall: our objective is  $[\bar{r}^t]_+ \to 0$ .
- To this end, choose  $q^{t+1}$  such that

$$E[\Delta r^{t+1}] \perp [\bar{r}^t]_+$$

So:

$$E[\Delta r^{t+1}] \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow (-\alpha^{t+1}q_2^{t+1}, \alpha^{t+1}q_1^{t+1}) \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow \alpha^{t+1}(q_1^{t+1}[\bar{r}_2^t]_+ - q_2^{t+1}[\bar{r}_1^t]_+) = 0$$

$$\Leftrightarrow q_1^{t+1} : q_2^{t+1} = [\bar{r}_1^t]_+ : [\bar{r}_2^t]_+.$$

The last equation amounts to proportional regret matching.

- Boundary cases are obvious and can be treated as follows:
  - If  $\bar{r}_1^t \le 0$  and  $\bar{r}_2^t > 0$ , then let  $q^{t+1} =_{Def} (0,1)$ .
  - If  $\bar{r}_1^t > 0$  and  $\bar{r}_2^t \le 0$ , then let  $q^{t+1} =_{Def} (1,0)$ .

- Recall: our objective is  $[\bar{r}^t]_+ \to 0$ .
- To this end, choose  $q^{t+1}$  such that

$$E[\Delta r^{t+1}] \perp [\bar{r}^t]_+$$

So:

$$E[\Delta r^{t+1}] \cdot [\bar{r}^t]_+ = 0$$


$$\Leftrightarrow (-\alpha^{t+1}q_2^{t+1}, \alpha^{t+1}q_1^{t+1}) \cdot [\bar{r}^t]_+ = 0$$

$$\Leftrightarrow \alpha^{t+1}(q_1^{t+1}[\bar{r}_2^t]_+ - q_2^{t+1}[\bar{r}_1^t]_+) = 0$$

$$\Leftrightarrow q_1^{t+1} : q_2^{t+1} = [\bar{r}_1^t]_+ : [\bar{r}_2^t]_+.$$


The last equation amounts to proportional regret matching.

- Boundary cases are obvious and can be treated as follows:
  - If  $\bar{r}_1^t \le 0$  and  $\bar{r}_2^t > 0$ , then let  $q^{t+1} =_{Def} (0,1)$ .
  - If  $\bar{r}_1^t > 0$  and  $\bar{r}_2^t \le 0$ , then let  $q^{t+1} =_{Def} (1,0)$ .
  - If all regret is non-positive, then play an action at random.




Expected incremental regret,  $E[\Delta r^{t+1}]$  is made orthogonal to the current regret, independently of the unknown  $\alpha^{t+1}$ .

- $E[\bar{r}^{t+1}]$  is a convex combination of  $\bar{r}_+^t$  and  $E[\Delta r^{t+1}]$ .
- Since  $E[\Delta r^{t+1}] \perp \bar{r}_+^t$ ,  $E[\bar{r}^{t+1}]$  lies closer to the non-positive orthant than  $\bar{r}_+^t$  does, provided t is large.
- Ultimately, the result follows from Blackwell's approachability theorem (Strategic Learning and its Limits, 2004, Ch. 4).




Expected incremental regret,  $E[\Delta r^{t+1}]$  is made orthogonal to the current regret, independently of the unknown  $\alpha^{t+1}$ .

- $E[\bar{r}^{t+1}]$  is a convex combination of  $\bar{r}_+^t$  and  $E[\Delta r^{t+1}]$ .
- Since  $E[\Delta r^{t+1}] \perp \bar{r}_+^t$ ,  $E[\bar{r}^{t+1}]$  lies closer to the non-positive orthant than  $\bar{r}_+^t$  does, provided t is large.
- Ultimately, the result follows from Blackwell's approachability theorem (Strategic Learning and its Limits, 2004, Ch. 4).




Expected incremental regret,  $E[\Delta r^{t+1}]$  is made orthogonal to the current regret, independently of the unknown  $\alpha^{t+1}$ .

- $E[\bar{r}^{t+1}]$  is a convex combination of  $\bar{r}_+^t$  and  $E[\Delta r^{t+1}]$ .
- Since  $E[\Delta r^{t+1}] \perp \bar{r}_+^t$ ,  $E[\bar{r}^{t+1}]$  lies closer to the non-positive orthant than  $\bar{r}_+^t$  does, provided t is large.
- Ultimately, the result follows from Blackwell's approachability theorem (Strategic Learning and its Limits, 2004, Ch. 4).




Expected incremental regret,  $E[\Delta r^{t+1}]$  is made orthogonal to the current regret, independently of the unknown  $\alpha^{t+1}$ .

- $E[\bar{r}^{t+1}]$  is a convex combination of  $\bar{r}_+^t$  and  $E[\Delta r^{t+1}]$ .
- Since  $E[\Delta r^{t+1}] \perp \bar{r}_+^t$ ,  $E[\bar{r}^{t+1}]$  lies closer to the non-positive orthant than  $\bar{r}_+^t$  does, provided t is large.
- Ultimately, the result follows from Blackwell's approachability theorem (Strategic Learning and its Limits, 2004, Ch. 4).




Expected incremental regret,  $E[\Delta r^{t+1}]$  is made orthogonal to the current regret, independently of the unknown  $\alpha^{t+1}$ .

- $E[\bar{r}^{t+1}]$  is a convex combination of  $\bar{r}_+^t$  and  $E[\Delta r^{t+1}]$ .
- Since  $E[\Delta r^{t+1}] \perp \bar{r}_+^t$ ,  $E[\bar{r}^{t+1}]$  lies closer to the non-positive orthant than  $\bar{r}_+^t$  does, provided t is large.
- Ultimately, the result follows from Blackwell's approachability theorem (Strategic Learning and its Limits, 2004, Ch. 4).




Expected incremental regret,  $E[\Delta r^{t+1}]$  is made orthogonal to the current regret, independently of the unknown  $\alpha^{t+1}$ .


- $E[\bar{r}^{t+1}]$  is a convex combination of  $\bar{r}_+^t$  and  $E[\Delta r^{t+1}]$ .
- Since  $E[\Delta r^{t+1}] \perp \bar{r}_+^t$ ,  $E[\bar{r}^{t+1}]$  lies closer to the non-positive orthant than  $\bar{r}_+^t$  does, provided t is large.
- Ultimately, the result follows from Blackwell's approachability theorem (Strategic Learning and its Limits, 2004, Ch. 4).




Expected incremental regret,  $E[\Delta r^{t+1}]$  is made orthogonal to the current regret, independently of the unknown  $\alpha^{t+1}$ .

- $E[\bar{r}^{t+1}]$  is a convex combination of  $\bar{r}_+^t$  and  $E[\Delta r^{t+1}]$ .
- Since  $E[\Delta r^{t+1}] \perp \bar{r}_+^t$ ,  $E[\bar{r}^{t+1}]$  lies closer to the non-positive orthant than  $\bar{r}_+^t$  does, provided t is large.
- Ultimately, the result follows from Blackwell's approachability theorem (Strategic Learning and its Limits, 2004, Ch. 4).






Expected incremental regret,  $E[\Delta r^{t+1}]$  is made orthogonal to the current regret, independently of the unknown  $\alpha^{t+1}$ .




Expected incremental regret,  $E[\Delta r^{t+1}]$  is made orthogonal to the current regret, independently of the unknown  $\alpha^{t+1}$ .



Expected incremental regret,  $E[\Delta r^{t+1}]$  is made orthogonal to the current regret, independently of the unknown  $\alpha^{t+1}$ .


Because at *A* does not know what *B* will play next, this is crucial.

■  $E[\bar{r}^{t+1}]$  is a convex combination of  $\bar{r}_+^t$  and  $E[\Delta r^{t+1}]$ .




Expected incremental regret,  $E[\Delta r^{t+1}]$  is made orthogonal to the current regret, independently of the unknown  $\alpha^{t+1}$ .


- $E[\bar{r}^{t+1}]$  is a convex combination of  $\bar{r}_+^t$  and  $E[\Delta r^{t+1}]$ .
- Since  $E[\Delta r^{t+1}] \perp \bar{r}_+^t$ ,  $E[\bar{r}^{t+1}]$  lies closer to the non-positive orthant than  $\bar{r}_+^t$  does, provided t is large.





Expected incremental regret,  $E[\Delta r^{t+1}]$  is made orthogonal to the current regret, independently of the unknown  $\alpha^{t+1}$ .


- $E[\bar{r}^{t+1}]$  is a convex combination of  $\bar{r}_+^t$  and  $E[\Delta r^{t+1}]$ .
- Since  $E[\Delta r^{t+1}] \perp \bar{r}_+^t$ ,  $E[\bar{r}^{t+1}]$  lies closer to the non-positive orthant than  $\bar{r}_+^t$  does, provided t is large.
- Ultimately, the result follows from Blackwell's approachability theorem (Strategic Learning and its Limits, 2004, Ch. 4).
















































































# Part III: *ϵ*-Greedy Off-policy Regret Matching

 $\epsilon$ -greedy regret matching. Let  $\epsilon > 0$  small.

- 1. **Explore**. Play randomly  $\epsilon$ % of the time.
- 2. Exploit. Else, play off-policy regret matching.

 $\epsilon$ -greedy regret matching. Let  $\epsilon > 0$  small.

- 1. **Explore**. Play randomly  $\epsilon$ % of the time.
- 2. Exploit. Else, play off-policy regret matching.

Define off-policy regret for x in round t as

$$\bar{r}_x^t =_{Def} \bar{u}_x^t(E) - \bar{u}^t$$
, where  $\bar{u}_x^t(E) = \left[ \frac{1}{|E_x|} \sum_{t \in E_x} u(x^t, y^t) \right]$ 

and  $E_x = \{ t \mid \text{player } A \text{ experimented in round } t \text{ and played } x \}.$ 

 $\epsilon$ -greedy regret matching. Let  $\epsilon > 0$  small.

- 1. **Explore**. Play randomly  $\epsilon$ % of the time.
- 2. Exploit. Else, play off-policy regret matching.

Define off-policy regret for x in round t as

$$\bar{r}_x^t =_{Def} \bar{u}_x^t(E) - \bar{u}^t$$
, where  $\bar{u}_x^t(E) = \left[ \frac{1}{|E_x|} \sum_{t \in E_x} u(x^t, y^t) \right]$ 

and  $E_x = \{ t \mid \text{player } A \text{ experimented in round } t \text{ and played } x \}.$ 

■ Proposed as a forecasting heuristic by Foster and Vohra (1993).

 $\epsilon$ -greedy regret matching. Let  $\epsilon > 0$  small.

- 1. **Explore**. Play randomly  $\epsilon$ % of the time.
- 2. Exploit. Else, play off-policy regret matching.

Define off-policy regret for x in round t as

$$\bar{r}_x^t =_{Def} \bar{u}_x^t(E) - \bar{u}^t$$
, where  $\bar{u}_x^t(E) = \left[ \frac{1}{|E_x|} \sum_{t \in E_x} u(x^t, y^t) \right]$ 

and  $E_x = \{ t \mid \text{player } A \text{ experimented in round } t \text{ and played } x \}.$ 

- Proposed as a forecasting heuristic by Foster and Vohra (1993).
- Does not need to know the actions of its opponents.

 $\epsilon$ -greedy regret matching. Let  $\epsilon > 0$  small.

- 1. **Explore**. Play randomly  $\epsilon$ % of the time.
- 2. Exploit. Else, play off-policy regret matching.

Define off-policy regret for x in round t as

$$\bar{r}_x^t =_{Def} \bar{u}_x^t(E) - \bar{u}^t$$
, where  $\bar{u}_x^t(E) = \left[ \frac{1}{|E_x|} \sum_{t \in E_x} u(x^t, y^t) \right]$ 

and  $E_x = \{ t \mid \text{player } A \text{ experimented in round } t \text{ and played } x \}.$ 

- Proposed as a forecasting heuristic by Foster and Vohra (1993).
- Does not need to know the actions of its opponents.
- Turns out to estimate regrets.

**Theorem** (Foster et al., 1998). For all  $\delta > 0$  there exists an  $\epsilon > 0$  such that  $\epsilon$ -greedy regret matching has at most  $\delta$  regret.

**Theorem** (Foster et al., 1998). For all  $\delta > 0$  there exists an  $\epsilon > 0$  such that  $\epsilon$ -greedy regret matching has at most  $\delta$  regret.

If  $\epsilon_t \to 0$  at a rate  $\mathcal{O}(t^{-1/3})$ , there is no regret in the long run.

**Theorem** (Foster et al., 1998). For all  $\delta > 0$  there exists an  $\epsilon > 0$  such that  $\epsilon$ -greedy regret matching has at most  $\delta$  regret.

If  $\epsilon_t \to 0$  at a rate  $\mathcal{O}(t^{-1/3})$ , there is no regret in the long run.

*Proof*. Suppose there are *k* different actions.

**Theorem** (Foster *et al.*, 1998). For all  $\delta > 0$  there exists an  $\epsilon > 0$  such that  $\epsilon$ -greedy regret matching has at most  $\delta$  regret.

If  $\epsilon_t \to 0$  at a rate  $\mathcal{O}(t^{-1/3})$ , there is no regret in the long run.

*Proof.* Suppose there are k different actions. Let  $e^t \in \mathbb{R}^k$  such that

 $e_x^t = ($  player A explores at t and chooses x ) ? 1:0.

**Theorem** (Foster et al., 1998). For all  $\delta > 0$  there exists an  $\epsilon > 0$  such that  $\epsilon$ -greedy regret matching has at most  $\delta$  regret.

If  $\epsilon_t \to 0$  at a rate  $\mathcal{O}(t^{-1/3})$ , there is no regret in the long run.

*Proof.* Suppose there are k different actions. Let  $e^t \in \mathbb{R}^k$  such that

 $e_x^t = ($  player A explores at t and chooses x ) ? 1 : 0.

For each action *i* 

$$Pr(x^t = i \mid A \text{ explores at round } t) = \frac{1}{k}.$$

**Theorem** (Foster *et al.*, 1998). For all  $\delta > 0$  there exists an  $\epsilon > 0$  such that  $\epsilon$ -greedy regret matching has at most  $\delta$  regret.

If  $\epsilon_t \to 0$  at a rate  $\mathcal{O}(t^{-1/3})$ , there is no regret in the long run.

*Proof.* Suppose there are k different actions. Let  $e^t \in \mathbb{R}^k$  such that

 $e_x^t = ($  player A explores at t and chooses x ) ? 1 : 0.

For each action *i* 

$$Pr(x^t = i \mid A \text{ explores at round } t) = \frac{1}{k}.$$

It follows that

$$E[e^t] = \left(\frac{\epsilon}{k}, \dots, \frac{\epsilon}{k}\right).$$

Define

$$z_x^t =_{Def} \left( \frac{\mathbf{k}}{\epsilon} \cdot e_x^t \cdot u(x, y^t) \right) - u(x, y^t).$$

Define

$$z_x^t =_{Def} \left( \frac{\mathbf{k}}{\epsilon} \cdot e_x^t \cdot u(x, y^t) \right) - u(x, y^t).$$

Define

$$z_x^t =_{Def} \left( \frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t) \right) - u(x, y^t).$$

Define

$$z_x^t =_{Def} \left(\frac{\mathbf{k}}{\epsilon} \cdot e_x^t \cdot u(x, y^t)\right) - u(x, y^t).$$

$$E[z_x^t]$$

Define

$$z_x^t =_{Def} \left( \frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t) \right) - u(x, y^t).$$

$$E[z_x^t] = E\left[\left(\frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t)\right) - u(x, y^t)\right]$$

Define

$$z_x^t =_{Def} \left( \frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t) \right) - u(x, y^t).$$

$$E[z_x^t] = E\left[\left(\frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t)\right) - u(x, y^t)\right]$$
$$= \frac{k}{\epsilon} \cdot E\left[e_x^t \cdot u(x, y^t)\right] - E[u(x, y^t)]$$

Define

$$z_x^t =_{Def} \left( \frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t) \right) - u(x, y^t).$$

$$E[z_x^t] = E\left[\left(\frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t)\right) - u(x, y^t)\right]$$

$$= \frac{k}{\epsilon} \cdot E\left[e_x^t \cdot u(x, y^t)\right] - E[u(x, y^t)]$$

$$= \frac{k}{\epsilon} \cdot E[e_x^t] \cdot E[u(x, y^t)] - E[u(x, y^t)] \quad (e^t \text{ and } u^t \text{ are independent})$$

Define

$$z_x^t =_{Def} \left( \frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t) \right) - u(x, y^t).$$

In words,  $z_x^t$  is the difference between the properly magnified empirical payoff for x and the (correct but) virtual payoff for x. It follows that

$$E[z_x^t] = E\left[\left(\frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t)\right) - u(x, y^t)\right]$$

$$= \frac{k}{\epsilon} \cdot E\left[e_x^t \cdot u(x, y^t)\right] - E[u(x, y^t)]$$

$$= \frac{k}{\epsilon} \cdot E[e_x^t] \cdot E[u(x, y^t)] - E[u(x, y^t)] \quad (e^t \text{ and } u^t \text{ are independent})$$

$$= \frac{k}{\epsilon} \cdot \frac{\epsilon}{k} \cdot E[u(x, y^t)] - E[u(x, y^t)]$$

Define

$$z_x^t =_{Def} \left( \frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t) \right) - u(x, y^t).$$

In words,  $z_x^t$  is the difference between the properly magnified empirical payoff for x and the (correct but) virtual payoff for x. It follows that

$$E[z_x^t] = E\left[\left(\frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t)\right) - u(x, y^t)\right]$$

$$= \frac{k}{\epsilon} \cdot E\left[e_x^t \cdot u(x, y^t)\right] - E[u(x, y^t)]$$

$$= \frac{k}{\epsilon} \cdot E[e_x^t] \cdot E[u(x, y^t)] - E[u(x, y^t)] \quad (e^t \text{ and } u^t \text{ are independent})$$

$$= \frac{k}{\epsilon} \cdot \frac{\epsilon}{k} \cdot E[u(x, y^t)] - E[u(x, y^t)]$$

$$= 0.$$



Strong law of large numbers for dependent random variables. Let  $\{w^t\}^t$  be a bounded sequence of possibly dependent random variables in  $R^k$ . Let  $z^t = E[w^t | w^{t-1}, w^{t-2}, \dots, w^1] - w^t$ , and  $\bar{z}^t$  the average of the  $z^t$ 's. Then  $\lim_{t\to\infty} \bar{z}^t = 0$  with probability one.

<sup>&</sup>lt;sup>a</sup>PY refers to Loève, 1978, Book II, Th. 32.E.1.

Strong law of large numbers for dependent random variables. Let  $\{w^t\}^t$  be a bounded sequence of possibly dependent random variables in  $R^k$ . Let  $z^t = E[w^t | w^{t-1}, w^{t-2}, \dots, w^1] - w^t$ , and  $\bar{z}^t$  the average of the  $z^t$ 's. Then  $\lim_{t\to\infty} \bar{z}^t = 0$  with probability one.

<sup>a</sup>PY refers to Loève, 1978, Book II, Th. 32.E.1.

We have

$$z_x^t =_{Def} \left( \frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t) \right) - u(x, y^t).$$

Strong law of large numbers for dependent random variables. Let  $\{w^t\}^t$  be a bounded sequence of possibly dependent random variables in  $R^k$ . Let  $z^t = E[w^t | w^{t-1}, w^{t-2}, \dots, w^1] - w^t$ , and  $\bar{z}^t$  the average of the  $z^t$ 's. Then  $\lim_{t\to\infty} \bar{z}^t = 0$  with probability one.

<sup>a</sup>PY refers to Loève, 1978, Book II, Th. 32.E.1.

We have

$$z_x^t =_{Def} \left( \frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t) \right) - u(x, y^t).$$

and

$$E[z_x^t] = 0.$$

Strong law of large numbers for dependent random variables. Let  $\{w^t\}^t$  be a bounded sequence of possibly dependent random variables in  $R^k$ . Let  $z^t = E[w^t | w^{t-1}, w^{t-2}, \dots, w^1] - w^t$ , and  $\bar{z}^t$  the average of the  $z^t$ 's. Then  $\lim_{t\to\infty} \bar{z}^t = 0$  with probability one.

<sup>a</sup>PY refers to Loève, 1978, Book II, Th. 32.E.1.

We have

$$z_x^t =_{Def} \left( \frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t) \right) - u(x, y^t).$$

and

$$E[z_x^t] = 0.$$

If

$$\bar{z}^t =_{Def} \text{average of } z^s, s \leq t$$

Strong law of large numbers for dependent random variables. Let  $\{w^t\}^t$  be a bounded sequence of possibly dependent random variables in  $R^k$ . Let  $z^t = E[w^t | w^{t-1}, w^{t-2}, \dots, w^1] - w^t$ , and  $\bar{z}^t$  the average of the  $z^t$ 's. Then  $\lim_{t\to\infty} \bar{z}^t = 0$  with probability one.

<sup>a</sup>PY refers to Loève, 1978, Book II, Th. 32.E.1.

We have

$$z_x^t =_{Def} \left( \frac{k}{\epsilon} \cdot e_x^t \cdot u(x, y^t) \right) - u(x, y^t).$$

and

$$E[z_x^t] = 0.$$

If

$$\bar{z}^t =_{Def} \text{average of } z^s, s \leq t$$

then from the strong law of large numbers for dependent random variables it follows that  $\lim_{t\to\infty} \bar{z}^t = 0$  a.s.

$$\bar{z}_{x}^{t} = \underbrace{\frac{1}{t} \sum_{s=1}^{t} \frac{k}{\epsilon} \cdot e_{x}^{s} \cdot u(x, y^{s}) - \bar{u}^{t}}_{\text{scaled}} - \underbrace{\frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}) - \bar{u}^{t}}_{\text{true regret}}$$
scaled
empirical regret

Now write  $\bar{z}_x^t$  as follows (!):

$$\bar{z}_{x}^{t} = \underbrace{\frac{1}{t} \sum_{s=1}^{t} \frac{k}{\epsilon} \cdot e_{x}^{s} \cdot u(x, y^{s}) - \bar{u}^{t}}_{\text{scaled}} - \underbrace{\frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}) - \bar{u}^{t}}_{\text{true regret}}$$
scaled
empirical regret

1. Since  $\lim_{t\to\infty} \bar{z}^t = 0$ , scaled empirical regret converges to true regret a.s.

$$\bar{z}_{x}^{t} = \underbrace{\frac{1}{t} \sum_{s=1}^{t} \frac{k}{\epsilon} \cdot e_{x}^{s} \cdot u(x, y^{s}) - \bar{u}^{t}}_{\text{scaled}} - \underbrace{\frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}) - \bar{u}^{t}}_{\text{true regret}}$$
scaled
empirical regret

- 1. Since  $\lim_{t\to\infty} \bar{z}^t = 0$ , scaled empirical regret converges to true regret a.s.
- 2.  $\epsilon$ % of the time *A* explores.

$$\bar{z}_{x}^{t} = \underbrace{\frac{1}{t} \sum_{s=1}^{t} \frac{k}{\epsilon} \cdot e_{x}^{s} \cdot u(x, y^{s}) - \bar{u}^{t}}_{\text{scaled}} - \underbrace{\frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}) - \bar{u}^{t}}_{\text{true regret}}$$
scaled
empirical regret

- 1. Since  $\lim_{t\to\infty} \bar{z}^t = 0$ , scaled empirical regret converges to true regret a.s.
- 2.  $\epsilon$ % of the time *A* explores.
- 3.  $(1 \epsilon)$ % of the time *A* plays empirical regret

$$\bar{z}_{x}^{t} = \underbrace{\frac{1}{t} \sum_{s=1}^{t} \frac{k}{\epsilon} \cdot e_{x}^{s} \cdot u(x, y^{s}) - \bar{u}^{t}}_{\text{scaled}} - \underbrace{\frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}) - \bar{u}^{t}}_{\text{true regret}}$$
scaled
empirical regret

- 1. Since  $\lim_{t\to\infty} \bar{z}^t = 0$ , scaled empirical regret converges to true regret a.s.
- 2.  $\epsilon$ % of the time *A* explores.
- 3.  $(1 \epsilon)\%$  of the time *A* plays empirical regret  $\rightsquigarrow$  true regret.

$$\bar{z}_{x}^{t} = \underbrace{\frac{1}{t} \sum_{s=1}^{t} \frac{k}{\epsilon} \cdot e_{x}^{s} \cdot u(x, y^{s}) - \bar{u}^{t}}_{\text{scaled}} - \underbrace{\frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}) - \bar{u}^{t}}_{\text{true regret}}$$
scaled
empirical regret

- 1. Since  $\lim_{t\to\infty} \bar{z}^t = 0$ , scaled empirical regret converges to true regret a.s.
- 4. In the long run, empirical regret is within  $2\epsilon$  from true regret.

- 2.  $\epsilon$ % of the time *A* explores.
- 3.  $(1 \epsilon)$ % of the time *A* plays empirical regret  $\rightsquigarrow$  true regret.

$$\bar{z}_{x}^{t} = \underbrace{\frac{1}{t} \sum_{s=1}^{t} \frac{k}{\epsilon} \cdot e_{x}^{s} \cdot u(x, y^{s}) - \bar{u}^{t}}_{\text{scaled}} - \underbrace{\frac{1}{t} \sum_{s=1}^{t} u(x, y^{s}) - \bar{u}^{t}}_{\text{true regret}}$$
scaled
empirical regret

- 1. Since  $\lim_{t\to\infty} \bar{z}^t = 0$ , scaled empirical regret converges to true regret a.s.
- 2.  $\epsilon$ % of the time *A* explores.
- 3.  $(1 \epsilon)\%$  of the time *A* plays empirical regret  $\rightsquigarrow$  true regret.

- 4. In the long run, empirical regret is within  $2\epsilon$  from true regret.
- 5. If  $\epsilon$  is set to  $\delta/2$ , then empirical regret remains within  $2 \cdot \delta/2$  from zero.

Author: Gerard Vreeswijk. Slides last modified on May  $11^{\mathrm{th}}$ , 2020 at 17:21

Regret matching can be traced to Blackwell's approachability theorem and Hannan's notion of universal consistency.

- Regret matching can be traced to Blackwell's approachability theorem and Hannan's notion of universal consistency.
- The diagram on regret matching is taken from Peyton Young, and Foster and Vohra (who formulate the problem from a decision theoretic point of view).

- Regret matching can be traced to Blackwell's approachability theorem and Hannan's notion of universal consistency.
- The diagram on regret matching is taken from Peyton Young, and Foster and Vohra (who formulate the problem from a decision theoretic point of view).
- The regret-matching algorithm and the analysis of its convergence to coarse correlated equilibria (a generalisation of Nash

equilibria) is given by Hart and Mas-Colell.

- Regret matching can be traced to Blackwell's approachability theorem and Hannan's notion of universal consistency.
- The diagram on regret matching is taken from Peyton Young, and Foster and Vohra (who formulate the problem from a decision theoretic point of view).
- The regret-matching algorithm and the analysis of its convergence to coarse correlated equilibria (a generalisation of Nash

equilibria) is given by Hart and Mas-Colell.

Blackwell, D. (1956). "Controlled random walks". *Proc. of the Int. Congress of Mathematicians*, North-Holland Publishing Comp., pp. 336-338.

- Regret matching can be traced to Blackwell's approachability theorem and Hannan's notion of universal consistency.
- The diagram on regret matching is taken from Peyton Young, and Foster and Vohra (who formulate the problem from a decision theoretic point of view).
- The regret-matching algorithm and the analysis of its convergence to coarse correlated equilibria (a generalisation of Nash

equilibria) is given by Hart and Mas-Colell.

Blackwell, D. (1956). "Controlled random walks". *Proc. of the Int. Congress of Mathematicians*, North-Holland Publishing Comp., pp. 336-338.

Hannan, J. F. (1957). "Approximation to Bayes risk in repeated plays". *Contributions to the Theory of Games*, **3**, pp. 97-139.

- Regret matching can be traced to Blackwell's approachability theorem and Hannan's notion of universal consistency.
- The diagram on regret matching is taken from Peyton Young, and Foster and Vohra (who formulate the problem from a decision theoretic point of view).
- The regret-matching algorithm and the analysis of its convergence to coarse correlated equilibria (a generalisation of Nash

equilibria) is given by Hart and Mas-Colell.

Blackwell, D. (1956). "Controlled random walks". *Proc. of the Int. Congress of Mathematicians*, North-Holland Publishing Comp., pp. 336-338.

Hannan, J. F. (1957). "Approximation to Bayes risk in repeated plays". *Contributions to the Theory of Games*, **3**, pp. 97-139.

Hart, S., and Mas-Colell, A. (2000). "A simple adaptive procedure leading to correlated equilibrium". *Econometrica*, **68**, pp. 1127-1150.

- Regret matching can be traced to Blackwell's approachability theorem and Hannan's notion of universal consistency.
- The diagram on regret matching is taken from Peyton Young, and Foster and Vohra (who formulate the problem from a decision theoretic point of view).
- The regret-matching algorithm and the analysis of its convergence to coarse correlated equilibria (a generalisation of Nash

equilibria) is given by Hart and Mas-Colell.

Blackwell, D. (1956). "Controlled random walks". *Proc. of the Int. Congress of Mathematicians*, North-Holland Publishing Comp., pp. 336-338.

Hannan, J. F. (1957). "Approximation to Bayes risk in repeated plays". *Contributions to the Theory of Games*, **3**, pp. 97-139.

Hart, S., and Mas-Colell, A. (2000). "A simple adaptive procedure leading to correlated equilibrium". *Econometrica*, **68**, pp. 1127-1150.

Foster, D., and Vohra, R. (1999). "Regret in the on-line decision problem". GEB: *Games and Economic Behavior*, **29**, pp. 7-36.

Author: Gerard Vreeswijk. Slides last modified on May  $11^{\mathrm{th}}$ , 2020 at 17:21

■ **Fictitious Play**. Monitor actions of opponent(s) and play a best response to most frequent actions. As opposed to no-regret, fictitious play is interested in the opponent's behaviour to predict future play.

- **Fictitious Play**. Monitor actions of opponent(s) and play a best response to most frequent actions. As opposed to no-regret, fictitious play is interested in the opponent's behaviour to predict future play.
- Smoothed fictitious play. With fictitious play, the probability to play sub-optimal responses is zero. Smoothed fictitious play plays sub-optimal responses proportional to their expected payoff, given opponents' play.

- **Fictitious Play**. Monitor actions of opponent(s) and play a best response to most frequent actions. As opposed to no-regret, fictitious play is interested in the opponent's behaviour to predict future play.
- Smoothed fictitious play. With fictitious play, the probability to play sub-optimal responses is zero. Smoothed fictitious play plays sub-optimal responses proportional to their expected payoff, given opponents' play.
- Conditional no-regret. Conditions on particular actions. There is regret if there is a pair of actions (x, x') such that, with hindsight, playing x' was better than playing x.

- **Fictitious Play**. Monitor actions of opponent(s) and play a best response to most frequent actions. As opposed to no-regret, fictitious play is interested in the opponent's behaviour to predict future play.
- Smoothed fictitious play. With fictitious play, the probability to play sub-optimal responses is zero. Smoothed fictitious play plays sub-optimal responses proportional to their expected payoff, given opponents' play.
- Conditional no-regret. Conditions on particular actions. There is regret if there is a pair of actions (x, x') such that, with hindsight, playing x' was better than playing x.
- **Satisficing Play**. While payoffs equal or supersede the average of past payoffs, keep playing the same action.