

# Multi-agent learning

## Repeated games

*Gerard Vreeswijk*, Intelligent Systems Group, Computer Science  
Department, Faculty of Sciences, Utrecht University, The  
Netherlands.

Friday 3<sup>rd</sup> May, 2019

# Repeated games: motivation

# Repeated games: motivation

1. Much *interaction* in MASs can be modelled through *games*.

# Repeated games: motivation

1. Much **interaction** in MASs can be modelled through *games*.
2. Much **learning** in MASs can therefore be modelled through *learning in games*.

# Repeated games: motivation

1. Much **interaction** in MASs can be modelled through *games*.
2. Much **learning** in MASs can therefore be modelled through *learning in games*.
3. Learning in games usually takes place through the (gradual) adaptation of strategies (hence, behaviour) in a **repeated game**.

# Repeated games: motivation

1. Much **interaction** in MASs can be modelled through *games*.
2. Much **learning** in MASs can therefore be modelled through *learning in games*.
3. Learning in games usually takes place through the (gradual) adaptation of strategies (hence, behaviour) in a **repeated game**.
4. In most repeated games, one game (a.k.a. **stage game**) is played repeatedly. Possibilities:

# Repeated games: motivation

1. Much **interaction** in MASs can be modelled through *games*.
2. Much **learning** in MASs can therefore be modelled through *learning in games*.
3. Learning in games usually takes place through the (gradual) adaptation of strategies (hence, behaviour) in a **repeated game**.
4. In most repeated games, one game (a.k.a. **stage game**) is played repeatedly. Possibilities:
  - A **finite** number of times.

# Repeated games: motivation

1. Much **interaction** in MASs can be modelled through *games*.
2. Much **learning** in MASs can therefore be modelled through *learning in games*.
3. Learning in games usually takes place through the (gradual) adaptation of strategies (hence, behaviour) in a **repeated game**.
4. In most repeated games, one game (a.k.a. **stage game**) is played repeatedly. Possibilities:
  - A **finite** number of times.
  - An **indefinite** (same: *indeterminate*) number of times.



# Repeated games: motivation

1. Much **interaction** in MASs can be modelled through *games*.
2. Much **learning** in MASs can therefore be modelled through *learning in games*.
3. Learning in games usually takes place through the (gradual) adaptation of strategies (hence, behaviour) in a **repeated game**.
4. In most repeated games, one game (a.k.a. **stage game**) is played repeatedly. Possibilities:
  - A **finite** number of times.
  - An **indefinite** (same: *indeterminate*) number of times.
  - An **infinite** number of times.

# Repeated games: motivation

1. Much **interaction** in MASs can be modelled through *games*.
2. Much **learning** in MASs can therefore be modelled through *learning in games*.
3. Learning in games usually takes place through the (gradual) adaptation of strategies (hence, behaviour) in a **repeated game**.
4. In most repeated games, one game (a.k.a. **stage game**) is played repeatedly. Possibilities:
  - A **finite** number of times.
  - An **indefinite** (same: *indeterminate*) number of times.
  - An **infinite** number of times.
5. Therefore, familiarity with the basic concepts and results from the theory of repeated games is essential to understand MAL.

# Plan for today

# Plan for today

- NE in normal form games that are repeated a **finite** number of times.

# Plan for today

- NE in normal form games that are repeated a **finite** number of times.
  - Principle of **backward induction**.

# Plan for today

- NE in normal form games that are repeated a **finite** number of times.
  - Principle of **backward induction**.
- NE in normal form games that are repeated an **indefinite** number of times.

# Plan for today

- NE in normal form games that are repeated a **finite** number of times.
  - Principle of **backward induction**.
- NE in normal form games that are repeated an **indefinite** number of times.
  - **Discount factor**. Models the probability of continuation.

# Plan for today

- NE in normal form games that are repeated a **finite** number of times.
  - Principle of **backward induction**.
- NE in normal form games that are repeated an **indefinite** number of times.
  - **Discount factor**. Models the probability of continuation.
  - **Folk theorem**. (Actually many FT's.) Repeated games generally do have infinitely many Nash equilibria.



# Plan for today

- NE in normal form games that are repeated a **finite** number of times.
  - Principle of **backward induction**.
- NE in normal form games that are repeated an **indefinite** number of times.
  - **Discount factor**. Models the probability of continuation.
  - **Folk theorem**. (Actually many FT's.) Repeated games generally do have infinitely many Nash equilibria.
  - **Trigger strategy**.

# Plan for today

- NE in normal form games that are repeated a **finite** number of times.
  - Principle of **backward induction**.
- NE in normal form games that are repeated an **indefinite** number of times.
  - **Discount factor**. Models the probability of continuation.
  - **Folk theorem**. (Actually many FT's.) Repeated games generally do have infinitely many Nash equilibria.
  - **Trigger strategy**. “on-path” vs. “off-path” play

# Plan for today

- NE in normal form games that are repeated a **finite** number of times.
  - Principle of **backward induction**.
- NE in normal form games that are repeated an **indefinite** number of times.
  - **Discount factor**. Models the probability of continuation.
  - **Folk theorem**. (Actually many FT's.) Repeated games generally do have infinitely many Nash equilibria.
  - **Trigger strategy**. “on-path” vs. “off-path” play, “**minmax**” as a threat.

# Plan for today

- NE in normal form games that are repeated a **finite** number of times.
  - Principle of **backward induction**.
- NE in normal form games that are repeated an **indefinite** number of times.
  - **Discount factor**. Models the probability of continuation.
  - **Folk theorem**. (Actually many FT's.) Repeated games generally do have infinitely many Nash equilibria.
  - **Trigger strategy**. “on-path” vs. “off-path” play, “**minmax**” as a threat.

This presentation draws heavily on (Peters, 2008).

# Plan for today

- NE in normal form games that are repeated a **finite** number of times.
  - Principle of **backward induction**.
- NE in normal form games that are repeated an **indefinite** number of times.
  - **Discount factor**. Models the probability of continuation.
  - **Folk theorem**. (Actually many FT's.) Repeated games generally do have infinitely many Nash equilibria.
  - **Trigger strategy**. “on-path” vs. “off-path” play, “**minmax**” as a threat.

This presentation draws heavily on (Peters, 2008).

\* H. Peters (2008): *Game Theory: A Multi-Leveled Approach*. Springer, ISBN: 978-3-540-69290-4. Ch. 8: Repeated games.

# **Part I:**

# **Nash equilibria**

# **Part I:**

## **Nash equilibria in normal form games**

# **Part I: Nash equilibria in normal form games that are repeated**



**Part I:**  
**Nash equilibria**  
**in normal form games**  
**that are repeated**  
**a *finite* number of times**

# Nash equilibria in playing the PD twice

Prisoners' Dilemma		<i>Other:</i>	
		Cooperate	Defect
You:	Cooperate	(3, 3)	(0, 5)
	Defect	(5, 0)	(1, 1)

# Nash equilibria in playing the PD twice

Prisoners' Dilemma		Other:	
		Cooperate	Defect
You:	Cooperate	(3,3)	(0,5)
	Defect	(5,0)	(1,1)

- Even if mixed strategies are allowed, the PD possesses **one Nash equilibrium**, viz. (D,D) with payoffs (1,1).

# Nash equilibria in playing the PD twice

Prisoners' Dilemma		Other:	
		Cooperate	Defect
You:	Cooperate	(3,3)	(0,5)
	Defect	(5,0)	(1,1)

- Even if mixed strategies are allowed, the PD possesses one Nash equilibrium, viz. (D,D) with payoffs (1,1).
- This equilibrium is Pareto sub-optimal.

# Nash equilibria in playing the PD twice

Prisoners' Dilemma		Other:	
		Cooperate	Defect
You:	Cooperate	(3,3)	(0,5)
	Defect	(5,0)	(1,1)


- Even if mixed strategies are allowed, the PD possesses **one Nash equilibrium**, viz. (D,D) with payoffs (1,1).
- This equilibrium is **Pareto sub-optimal**.
- Does the situation change if two parties get to play the Prisoners' Dilemma **two times** in succession?

# Nash equilibria in playing the PD twice


Prisoners' Dilemma		Other:	
		Cooperate	Defect
You:	Cooperate	(3,3)	(0,5)
	Defect	(5,0)	(1,1)

- Even if mixed strategies are allowed, the PD possesses **one Nash equilibrium**, viz. (D,D) with payoffs (1,1).
- This equilibrium is **Pareto sub-optimal**.
- Does the situation change if two parties get to play the Prisoners' Dilemma **two times** in succession?
- The following diagram (hopefully) shows that playing the PD two times in succession does **not** yield an essentially new NE.

# Nash equilibria in playing the PD twice



# Nash equilibria in playing the PD twice



P.S. This is just a payoff tree, not a game in extensive form!



# Nash equilibria in playing the PD twice

In normal form:

		<i>Other:</i>			
		CC	CD	DC	DD
You:	CC	(6, 6)	(3, 8)	(3, 8)	(0, 10)
	CD	(8, 3)	(4, 4)	(5, 5)	(1, 6)
	DC	(8, 3)	(5, 5)	(4, 4)	(1, 6)
	DD	(10, 0)	(6, 1)	(6, 1)	(2, 2)

# Nash equilibria in playing the PD twice

In normal form:

		<i>Other:</i>			
		CC	CD	DC	DD
You:	CC	(6, 6)	(3, 8)	(3, 8)	(0, 10)
	CD	(8, 3)	(4, 4)	(5, 5)	(1, 6)
	DC	(8, 3)	(5, 5)	(4, 4)	(1, 6)
	DD	(10, 0)	(6, 1)	(6, 1)	(2, 2)

- The action profile (*DD*, *DD*) is the only Nash equilibrium.

# Nash equilibria in playing the PD twice

In normal form:

		<i>Other:</i>			
		CC	CD	DC	DD
You:	CC	(6, 6)	(3, 8)	(3, 8)	(0, 10)
	CD	(8, 3)	(4, 4)	(5, 5)	(1, 6)
	DC	(8, 3)	(5, 5)	(4, 4)	(1, 6)
	DD	(10, 0)	(6, 1)	(6, 1)	(2, 2)

- The action profile (*DD*, *DD*) is the only Nash equilibrium.
- With 3 successive games, we obtain a  $2^3 \times 2^3$  matrix, where the action profile (*DDD*, *DDD*) still would be the only Nash equilibrium.

# Nash equilibria in playing the PD twice

In normal form:

		<i>Other:</i>			
		CC	CD	DC	DD
You:	CC	(6, 6)	(3, 8)	(3, 8)	(0, 10)
	CD	(8, 3)	(4, 4)	(5, 5)	(1, 6)
	DC	(8, 3)	(5, 5)	(4, 4)	(1, 6)
	DD	(10, 0)	(6, 1)	(6, 1)	(2, 2)

- The action profile ( $DD, DD$ ) is the only Nash equilibrium.
- With 3 successive games, we obtain a  $2^3 \times 2^3$  matrix, where the action profile ( $DDD, DDD$ ) still would be the only Nash equilibrium.
- Generalise to  $N$  repetitions: ( $DD^{N-1}, DD^{N-1}$ ) still is the only Nash equilibrium in a repeated game where the PD is played  $N$  times in succession.

# **Part II:**

# **Nash equilibria**

# **Part II:**

## **Nash equilibria in normal form games**

# **Part II: Nash equilibria in normal form games that are repeated**

**Part II:**  
**Nash equilibria**  
**in normal form games**  
**that are repeated**  
**an indefinite number of times**



# Terminology: finite, indefinite, infinite

To repeat an experiment ...



# Terminology: finite, indefinite, infinite

To repeat an experiment ...

■ ... **ten** times.



# Terminology: finite, indefinite, infinite

To repeat an experiment ...

■ ... **ten** times. That's hopefully clear.



# Terminology: finite, indefinite, infinite

To repeat an experiment ...

- ... **ten** times. That's hopefully clear.
- ... a **finite** number of times.



# Terminology: finite, indefinite, infinite

To repeat an experiment ...

- ... **ten** times. That's hopefully clear.
- ... a **finite** number of times. May mean: a **fixed number of times**, where the number of repetitions is determined beforehand.



# Terminology: finite, indefinite, infinite

To repeat an experiment ...

- ... **ten** times. That's hopefully clear.
- ... a **finite** number of times. May mean: a **fixed number of times**, where the number of repetitions is determined beforehand. Or it may mean: **an indefinite number of times**.



# Terminology: finite, indefinite, infinite

To repeat an experiment ...

- ... **ten** times. That's hopefully clear.
- ... a **finite** number of times. May mean: a **fixed number of times**, where the number of repetitions is determined beforehand. Or it may mean: **an indefinite number of times**. Depends on the context!



# Terminology: finite, indefinite, infinite

To repeat an experiment ...

- ... **ten** times. That's hopefully clear.
- ... a **finite** number of times. May mean: a **fixed number of times**, where the number of repetitions is determined beforehand. Or it may mean: **an indefinite number of times**. Depends on the context!
- ... an **indefinite** number of times.





# Terminology: finite, indefinite, infinite

To repeat an experiment ...

- ... **ten** times. That's hopefully clear.
- ... a **finite** number of times. May mean: a **fixed number of times**, where the number of repetitions is determined beforehand. Or it may mean: **an indefinite number of times**. Depends on the context!
- ... an **indefinite** number of times. Means: a finite number of times, but nothing is known beforehand about the number of repetitions.



# Terminology: finite, indefinite, infinite

To repeat an experiment ...

- ... **ten** times. That's hopefully clear.
- ... a **finite** number of times. May mean: a **fixed number of times**, where the number of repetitions is determined beforehand. Or it may mean: **an indefinite number of times**. Depends on the context!
- ... an **indefinite** number of times. Means: a finite number of times, but nothing is known beforehand about the number of repetitions.
- ... an **infinite** number of times.



# Terminology: finite, indefinite, infinite

To repeat an experiment ...

- ... **ten** times. That's hopefully clear.
- ... a **finite** number of times. May mean: a **fixed number of times**, where the number of repetitions is determined beforehand. Or it may mean: **an indefinite number of times**. Depends on the context!
- ... an **indefinite** number of times. Means: a finite number of times, but nothing is known beforehand about the number of repetitions.
- ... an **infinite** number of times. When throwing a dice this must mean a **countably infinite** number of times.



# Indefinite number of repetitions

# Indefinite number of repetitions

- A Pareto-suboptimal outcome can be avoided in case the following three conditions are met.

# Indefinite number of repetitions

- A Pareto-suboptimal outcome can be avoided in case the following three conditions are met.
  1. The Prisoners' Dilemma is repeated an **indefinite** number of times (rounds).

# Indefinite number of repetitions

- A Pareto-suboptimal outcome can be avoided in case the following three conditions are met.
  1. The Prisoners' Dilemma is repeated an **indefinite** number of times (rounds).
  2. A so-called **discount factor**  $\delta \in [0, 1]$  determines **the probability of continuing the game** after each round.

# Indefinite number of repetitions

- A Pareto-suboptimal outcome can be avoided in case the following three conditions are met.
  1. The Prisoners' Dilemma is repeated an **indefinite** number of times (rounds).
  2. A so-called **discount factor**  $\delta \in [0, 1]$  determines **the probability of continuing the game** after each round.
  3. The probability to continue,  $\delta$ , is large enough.



# Indefinite number of repetitions

- A Pareto-suboptimal outcome can be avoided in case the following three conditions are met.
  1. The Prisoners' Dilemma is repeated an **indefinite** number of times (rounds).
  2. A so-called **discount factor**  $\delta \in [0, 1]$  determines **the probability of continuing the game** after each round.
  3. The probability to continue,  $\delta$ , is large enough.
- Under these conditions suddenly **infinitely many** Nash equilibria exist. This is sometimes called an **embarrassment of richness** (Peters, 2008).

# Indefinite number of repetitions

- A Pareto-suboptimal outcome can be avoided in case the following three conditions are met.
  1. The Prisoners' Dilemma is repeated an **indefinite** number of times (rounds).
  2. A so-called **discount factor**  $\delta \in [0, 1]$  determines **the probability of continuing the game** after each round.
  3. The probability to continue,  $\delta$ , is large enough.
- Under these conditions suddenly **infinitely many** Nash equilibria exist. This is sometimes called an **embarrassment of richness** (Peters, 2008).
- Various **Folk theorems** state the existence of multiple equilibria in games that are repeated an indefinite number of times.

# Indefinite number of repetitions

- A Pareto-suboptimal outcome can be avoided in case the following three conditions are met.
  1. The Prisoners' Dilemma is repeated an **indefinite** number of times (rounds).
  2. A so-called **discount factor**  $\delta \in [0, 1]$  determines **the probability of continuing the game** after each round.
  3. The probability to continue,  $\delta$ , is large enough.
- Under these conditions suddenly **infinitely many** Nash equilibria exist. This is sometimes called an **embarrassment of richness** (Peters, 2008).
- Various **Folk theorems** state the existence of multiple equilibria in games that are repeated an indefinite number of times.
- Here we discuss one version of “the” Folk Theorem.

# Family of Folk Theorems

There actually exist many Folk Theorems.

# Family of Folk Theorems

There actually exist many Folk Theorems.

- **Horizon.** The game may be repeated indefinitely (present case)

# Family of Folk Theorems

There actually exist many Folk Theorems.

- **Horizon.** The game may be repeated indefinitely (present case) or there may be an upper bound to the number of plays.

# Family of Folk Theorems

There actually exist many Folk Theorems.

- **Horizon.** The game may be repeated indefinitely (present case) or there may be an upper bound to the number of plays.
- **Information.** Players may act on the basis of **CKR** (present case)

# Family of Folk Theorems

There actually exist many Folk Theorems.

- **Horizon.** The game may be repeated indefinitely (present case) or there may be an upper bound to the number of plays.
- **Information.** Players may act on the basis of **CKR** (present case), or certain parts of the history may be hidden.



# Family of Folk Theorems

There actually exist many Folk Theorems.

- **Horizon.** The game may be repeated indefinitely (present case) or there may be an upper bound to the number of plays.
- **Information.** Players may act on the basis of **CKR** (present case), or certain parts of the history may be hidden.
- **Reward.** Players may collect their payoff through a discount factor (present case)

# Family of Folk Theorems

There actually exist many Folk Theorems.

- **Horizon.** The game may be repeated indefinitely (present case) or there may be an upper bound to the number of plays.
- **Information.** Players may act on the basis of **CKR** (present case), or certain parts of the history may be hidden.
- **Reward.** Players may collect their payoff through a discount factor (present case) or through average rewards.

# Family of Folk Theorems

There actually exist many Folk Theorems.

- **Horizon.** The game may be repeated indefinitely (present case) or there may be an upper bound to the number of plays.
- **Information.** Players may act on the basis of **CKR** (present case), or certain parts of the history may be hidden.
- **Reward.** Players may collect their payoff through a discount factor (present case) or through average rewards.
- **Subgame perfectness.** Subgame perfect equilibria (present case)

# Family of Folk Theorems

There actually exist many Folk Theorems.

- **Horizon.** The game may be repeated indefinitely (present case) or there may be an upper bound to the number of plays.
- **Information.** Players may act on the basis of **CKR** (present case), or certain parts of the history may be hidden.
- **Reward.** Players may collect their payoff through a discount factor (present case) or through average rewards.
- **Subgame perfectness.** Subgame perfect equilibria (present case) or plain Nash equilibria.

# Family of Folk Theorems

There actually exist many Folk Theorems.

- **Horizon.** The game may be repeated indefinitely (present case) or there may be an upper bound to the number of plays.
- **Information.** Players may act on the basis of **CKR** (present case), or certain parts of the history may be hidden.
- **Reward.** Players may collect their payoff through a discount factor (present case) or through average rewards.
- **Subgame perfectness.** Subgame perfect equilibria (present case) or plain Nash equilibria.
- **Equilibrium.** We may be interested in Nash equilibria (present case)

# Family of Folk Theorems

There actually exist many Folk Theorems.

- **Horizon.** The game may be repeated indefinitely (present case) or there may be an upper bound to the number of plays.
- **Information.** Players may act on the basis of **CKR** (present case), or certain parts of the history may be hidden.
- **Reward.** Players may collect their payoff through a discount factor (present case) or through average rewards.
- **Subgame perfectness.** Subgame perfect equilibria (present case) or plain Nash equilibria.
- **Equilibrium.** We may be interested in Nash equilibria (present case), or other types of equilibria, such as so-called  $\epsilon$ -Nash equilibria or so-called *correlated equilibria*.

# The concept of a repeated game

# The concept of a repeated game

- Let  $G$  be a game in normal form.



# The concept of a repeated game

- Let  $G$  be a game in normal form.
- The repeated game  $G^*(\delta)$ , is  $G$ , played an indefinite number of times

# The concept of a repeated game

- Let  $G$  be a game in normal form.
- The repeated game  $G^*(\delta)$ , is  $G$ , played an indefinite number of times, where  $\delta$  represents the probability that the game will be played another time.

# The concept of a repeated game

- Let  $G$  be a game in normal form.
- The repeated game  $G^*(\delta)$ , is  $G$ , played an indefinite number of times, where  $\delta$  represents the probability that the game will be played another time.

Exercise: give  $P\{ G^*(\delta) \text{ lasts at least } t \text{ rounds} \}$ .

# The concept of a repeated game

- Let  $G$  be a game in normal form.
- The repeated game  $G^*(\delta)$ , is  $G$ , played an indefinite number of times, where  $\delta$  represents the probability that the game will be played another time.

Exercise: give  $P\{ G^*(\delta) \text{ lasts at least } t \text{ rounds} \}$ . Answer:  $\delta^t$ .

# The concept of a repeated game

- Let  $G$  be a game in normal form.
- The repeated game  $G^*(\delta)$ , is  $G$ , played an indefinite number of times, where  $\delta$  represents the probability that the game will be played another time.

Exercise: give  $P\{ G^*(\delta) \text{ lasts at least } t \text{ rounds} \}$ . Answer:  $\delta^t$ .

- $G$  is called the stage game.

# The concept of a repeated game

- Let  $G$  be a game in normal form.
- The repeated game  $G^*(\delta)$ , is  $G$ , played an indefinite number of times, where  $\delta$  represents the probability that the game will be played another time.

Exercise: give  $P\{ G^*(\delta) \text{ lasts at least } t \text{ rounds} \}$ . Answer:  $\delta^t$ .

- $G$  is called the stage game.
- A history  $h$  of length  $t$  of a repeated game is a sequence of action profiles of length  $t$ .

# The concept of a repeated game

- Let  $G$  be a game in normal form.
- The **repeated game**  $G^*(\delta)$ , is  $G$ , played an indefinite number of times, where  $\delta$  represents the probability that the game will be played another time.

Exercise: give  $P\{ G^*(\delta) \text{ lasts at least } t \text{ rounds} \}$ . Answer:  $\delta^t$ .

- $G$  is called the **stage game**.
- A **history**  $h$  of length  $t$  of a repeated game is a sequence of action profiles of length  $t$ .

Example: (for the prisoner's dilemma):

Row player:	C	D	D	D	C	C	D	D	D	D
Column player:	C	D	D	D	D	D	D	C	D	D
	0	1	2	3	4	5	6	7	8	9

# The concept of a repeated game (II)



# The concept of a repeated game (II)

- The set of all possible histories (of any length) is denoted by  $H$ .

# The concept of a repeated game (II)

- The set of all possible histories (of any length) is denoted by  $H$ .
- A **strategy** for Player  $i$  is a function  $s_i : H \rightarrow \Delta\{C, D\}$  such that

$$\Pr(\text{Player } i \text{ plays } C \text{ in round } |h| + 1 \mid h) = s_i(h)(C).$$

# The concept of a repeated game (II)

- The set of all possible histories (of any length) is denoted by  $H$ .

- A **strategy** for Player  $i$  is a function  $s_i : H \rightarrow \Delta\{C, D\}$  such that

$$\Pr(\text{Player } i \text{ plays } C \text{ in round } |h| + 1 \mid h) = s_i(h)(C).$$

- A **strategy profile**  $s$  is a combination of strategies, one for each player.

# The concept of a repeated game (II)

- The set of all possible histories (of any length) is denoted by  $H$ .
- A **strategy** for Player  $i$  is a function  $s_i : H \rightarrow \Delta\{C, D\}$  such that

$$\Pr(\text{Player } i \text{ plays } C \text{ in round } |h| + 1 \mid h) = s_i(h)(C).$$

- A **strategy profile**  $s$  is a combination of strategies, one for each player.
- The **expected payoff** for player  $i$  given  $s$  can be computed. It is

$$\text{Expected payoff}_i(s) = \sum_{t=0}^{\infty} \delta^t \text{Expected payoff}_{i,t}(s).$$

Example on next page.

# The concept of a repeated game (II)

Repeated from the previous slide:

The **expected payoff** for player  $i$  given  $s$  can be computed. It is

$$\text{Expected payoff}_i(s) = \sum_{t=0}^{\infty} \delta^t \text{Expected payoff}_{i,t}(s).$$

# The concept of a repeated game (II)

Repeated from the previous slide:

The **expected payoff** for player  $i$  given  $s$  can be computed. It is

$$\text{Expected payoff}_i(s) = \sum_{t=0}^{\infty} \delta^t \text{Expected payoff}_{i,t}(s).$$

*Example:* prisoner's dilemma, strategy Player 1 is  $s_1 =$  "always cooperate 80%";

# The concept of a repeated game (II)

Repeated from the previous slide:

The **expected payoff** for player  $i$  given  $s$  can be computed. It is

$$\text{Expected payoff}_i(s) = \sum_{t=0}^{\infty} \delta^t \text{Expected payoff}_{i,t}(s).$$

*Example:* prisoner's dilemma, strategy Player 1 is  $s_1 =$  "always cooperate 80%"; strategy Player 2 is  $s_2 =$  "always cooperate 70%";

# The concept of a repeated game (II)

Repeated from the previous slide:

The **expected payoff** for player  $i$  given  $s$  can be computed. It is

$$\text{Expected payoff}_i(s) = \sum_{t=0}^{\infty} \delta^t \text{Expected payoff}_{i,t}(s).$$

*Example:* prisoner's dilemma, strategy Player 1 is  $s_1 =$  "always cooperate 80%"; strategy Player 2 is  $s_2 =$  "always cooperate 70%";  $\delta = 1/2$ .



# The concept of a repeated game (II)

Repeated from the previous slide:

The **expected payoff** for player  $i$  given  $s$  can be computed. It is

$$\text{Expected payoff}_i(s) = \sum_{t=0}^{\infty} \delta^t \text{Expected payoff}_{i,t}(s).$$

*Example:* prisoner's dilemma, strategy Player 1 is  $s_1 =$  "always cooperate 80%"; strategy Player 2 is  $s_2 =$  "always cooperate 70%";  $\delta = 1/2$ .

$$\begin{aligned} \text{Expected payoff}_1(s) &= \sum_{t=0}^{\infty} \left[ \left( \frac{1}{2} \right)^t [0.8(0.7 \cdot 3 + 0.3 \cdot 0) + 0.2(0.7 \cdot 5 + 0.3 \cdot 1)] \right] \\ &= \frac{1}{1 - 1/2} [\dots] \approx \frac{1}{1 - 1/2} 2.44 = 2 \times 2.44 = 4.88. \end{aligned}$$

# Subgame perfect equilibria of $G^*(\delta)$ : $D^*$

**Definition.** A strategy profile  $s$  for  $G^*(\delta)$  is a **subgame-perfect Nash equilibrium** if (1) it is a Nash equilibrium of this repeated game, and (2) for every subgame (i.e., tail game) of this repeated game,  $s$  restricted to that subgame is a Nash equilibrium as well.

# Subgame perfect equilibria of $G^*(\delta)$ : $D^*$

**Definition.** A strategy profile  $s$  for  $G^*(\delta)$  is a **subgame-perfect Nash equilibrium** if (1) it is a Nash equilibrium of this repeated game, and (2) for every subgame (i.e., tail game) of this repeated game,  $s$  restricted to that subgame is a Nash equilibrium as well.

Consider the strategy of iterated defection  $D^*$ : “always defect, no matter what”.<sup>1</sup>

# Subgame perfect equilibria of $G^*(\delta)$ : $D^*$

**Definition.** A strategy profile  $s$  for  $G^*(\delta)$  is a **subgame-perfect Nash equilibrium** if (1) it is a Nash equilibrium of this repeated game, and (2) for every subgame (i.e., tail game) of this repeated game,  $s$  restricted to that subgame is a Nash equilibrium as well.

Consider the strategy of iterated defection  $D^*$ : “always defect, no matter what”.<sup>1</sup>

**Claim.** *The strategy profile  $(D^*, D^*)$  is a subgame perfect equilibrium in  $G^*(\delta)$ .*

# Subgame perfect equilibria of $G^*(\delta)$ : $D^*$

**Definition.** A strategy profile  $s$  for  $G^*(\delta)$  is a **subgame-perfect Nash equilibrium** if (1) it is a Nash equilibrium of this repeated game, and (2) for every subgame (i.e., tail game) of this repeated game,  $s$  restricted to that subgame is a Nash equilibrium as well.

Consider the strategy of iterated defection  $D^*$ : “always defect, no matter what”.<sup>1</sup>

**Claim.** *The strategy profile  $(D^*, D^*)$  is a subgame perfect equilibrium in  $G^*(\delta)$ .*

*Proof.* Consider any tailgame starting at round  $t \geq 0$ .

# Subgame perfect equilibria of $G^*(\delta)$ : $D^*$

**Definition.** A strategy profile  $s$  for  $G^*(\delta)$  is a **subgame-perfect Nash equilibrium** if (1) it is a Nash equilibrium of this repeated game, and (2) for every subgame (i.e., tail game) of this repeated game,  $s$  restricted to that subgame is a Nash equilibrium as well.

Consider the strategy of iterated defection  $D^*$ : “always defect, no matter what”.<sup>1</sup>

**Claim.** *The strategy profile  $(D^*, D^*)$  is a subgame perfect equilibrium in  $G^*(\delta)$ .*

*Proof.* Consider any tailgame starting at round  $t \geq 0$ . We are done if we can show that  $(D^*, D^*)$  is a NE for this subgame.

# Subgame perfect equilibria of $G^*(\delta)$ : $D^*$

**Definition.** A strategy profile  $s$  for  $G^*(\delta)$  is a **subgame-perfect Nash equilibrium** if (1) it is a Nash equilibrium of this repeated game, and (2) for every subgame (i.e., tail game) of this repeated game,  $s$  restricted to that subgame is a Nash equilibrium as well.

Consider the strategy of iterated defection  $D^*$ : “always defect, no matter what”.<sup>1</sup>

**Claim.** *The strategy profile  $(D^*, D^*)$  is a subgame perfect equilibrium in  $G^*(\delta)$ .*

*Proof.* Consider any tailgame starting at round  $t \geq 0$ . We are done if we can show that  $(D^*, D^*)$  is a NE for this subgame. This is true: given that one player always defects, it never pays off for the other player to play  $C$  at any time.

# Subgame perfect equilibria of $G^*(\delta)$ : $D^*$

**Definition.** A strategy profile  $s$  for  $G^*(\delta)$  is a **subgame-perfect Nash equilibrium** if (1) it is a Nash equilibrium of this repeated game, and (2) for every subgame (i.e., tail game) of this repeated game,  $s$  restricted to that subgame is a Nash equilibrium as well.

Consider the strategy of iterated defection  $D^*$ : “always defect, no matter what”.<sup>1</sup>

**Claim.** *The strategy profile  $(D^*, D^*)$  is a subgame perfect equilibrium in  $G^*(\delta)$ .*

*Proof.* Consider any tailgame starting at round  $t \geq 0$ . We are done if we can show that  $(D^*, D^*)$  is a NE for this subgame. This is true: given that one player always defects, it never pays off for the other player to play  $C$  at any time. Therefore, everyone sticks to  $D^*$ .

<sup>1</sup>A notation like  $D^*$  or (worse)  $D^\infty$  is suggestive. Mathematically it makes no sense, but intuitively it does.



# **Part III:**

## **Trigger strategies**

# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

# Cost of deviating in Round $N$

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ .

# Cost of deviating in Round $N$

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1}$$

# Cost of deviating in Round $N$

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3$$

# Cost of deviating in Round $N$

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5$$

# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1$$



# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1$$

By playing  $C$  throughout, he *could* have earned

# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1$$

By playing  $C$  throughout, he *could* have earned  $\sum_{t=0}^{\infty} \delta^t \cdot 3$

# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1$$

By playing  $C$  throughout, he *could* have earned  $\sum_{t=0}^{\infty} \delta^t \cdot 3$

# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1$$

By playing  $C$  throughout, he *could* have earned  $\sum_{t=0}^{\infty} \delta^t \cdot 3$  which means he forfeited

# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1$$

By playing  $C$  throughout, he *could* have earned  $\sum_{t=0}^{\infty} \delta^t \cdot 3$  which means he forfeited

$$\left[ \sum_{t=0}^{\infty} \delta^t \cdot 3 \right]$$

# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1$$

By playing  $C$  throughout, he *could* have earned  $\sum_{t=0}^{\infty} \delta^t \cdot 3$  which means he forfeited

$$\left[ \sum_{t=0}^{\infty} \delta^t \cdot 3 \right] - \left[ \sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1 \right]$$

# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1$$

By playing  $C$  throughout, he *could* have earned  $\sum_{t=0}^{\infty} \delta^t \cdot 3$  which means he forfeited

$$\left[ \sum_{t=0}^{\infty} \delta^t \cdot 3 \right] - \left[ \sum_{t=0}^{N-1} \delta^t \cdot 3 \right]$$

# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1$$

By playing  $C$  throughout, he *could* have earned  $\sum_{t=0}^{\infty} \delta^t \cdot 3$  which means he forfeited

$$\left[ \sum_{t=0}^{\infty} \delta^t \cdot 3 \right] - \left[ \sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 \right]$$



# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1$$

By playing  $C$  throughout, he *could* have earned  $\sum_{t=0}^{\infty} \delta^t \cdot 3$  which means he forfeited

$$\left[ \sum_{t=0}^{\infty} \delta^t \cdot 3 \right] - \left[ \sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1 \right]$$

# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1$$

By playing  $C$  throughout, he *could* have earned  $\sum_{t=0}^{\infty} \delta^t \cdot 3$  which means he forfeited

$$\left[ \sum_{t=0}^{\infty} \delta^t \cdot 3 \right] - \left[ \sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1 \right] = -2\delta^N$$

# Cost of deviating in Round N

Consider the so-called **trigger strategy**  $T$ : “always play  $C$  unless  $D$  has been played at least once. In that case play  $D$  forever”.

**Claim.** *The strategy profile  $(T, T)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

*Proof.* Suppose one player starts to defect at Round  $N$ . By doing so he expects a payoff of

$$\sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1$$

By playing  $C$  throughout, he *could* have earned  $\sum_{t=0}^{\infty} \delta^t \cdot 3$  which means he forfeited

$$\left[ \sum_{t=0}^{\infty} \delta^t \cdot 3 \right] - \left[ \sum_{t=0}^{N-1} \delta^t \cdot 3 + \delta^N \cdot 5 + \sum_{t=N+1}^{\infty} \delta^t \cdot 1 \right] = -2\delta^N + 2 \sum_{t=N+1}^{\infty} \delta^t.$$

# Cost of deviating in Round N

So if

$$-2\delta^N + 2 \sum_{t=N+1}^{\infty} \delta^t > 0$$

there is forfeit from payoff.

# Cost of deviating in Round N

So if

$$-2\delta^N + 2 \sum_{t=N+1}^{\infty} \delta^t > 0$$

there is forfeit from payoff. This is when

$$\sum_{t=N+1}^{\infty} \delta^t > \delta^N$$

# Cost of deviating in Round N

So if

$$-2\delta^N + 2 \sum_{t=N+1}^{\infty} \delta^t > 0$$

there is forfeit from payoff. This is when

$$\sum_{t=N+1}^{\infty} \delta^t > \delta^N$$

$$\Leftrightarrow \delta^{N+1} \sum_{t=0}^{\infty} \delta^t > \delta^N$$

# Cost of deviating in Round N

So if

$$-2\delta^N + 2 \sum_{t=N+1}^{\infty} \delta^t > 0$$

there is forfeit from payoff. This is when

$$\sum_{t=N+1}^{\infty} \delta^t > \delta^N$$

$$\Leftrightarrow \delta^{N+1} \sum_{t=0}^{\infty} \delta^t > \delta^N \quad \Leftrightarrow \quad \delta = 1 \text{ xor } \delta^{N+1} \frac{1}{1-\delta} > \delta^N$$

# Cost of deviating in Round N

So if

$$-2\delta^N + 2 \sum_{t=N+1}^{\infty} \delta^t > 0$$

there is forfeit from payoff. This is when

$$\sum_{t=N+1}^{\infty} \delta^t > \delta^N$$

$$\Leftrightarrow \delta^{N+1} \sum_{t=0}^{\infty} \delta^t > \delta^N \quad \Leftrightarrow \quad \delta = 1 \text{ xor } \delta^{N+1} \frac{1}{1-\delta} > \delta^N$$

$$\Leftrightarrow \delta = 1 \text{ xor } \delta \frac{1}{1-\delta} > 1, \delta \neq 0$$



# Cost of deviating in Round N

So if

$$-2\delta^N + 2 \sum_{t=N+1}^{\infty} \delta^t > 0$$

there is forfeit from payoff. This is when

$$\sum_{t=N+1}^{\infty} \delta^t > \delta^N$$

$$\Leftrightarrow \delta^{N+1} \sum_{t=0}^{\infty} \delta^t > \delta^N \quad \Leftrightarrow \quad \delta = 1 \text{ xor } \delta^{N+1} \frac{1}{1-\delta} > \delta^N$$

$$\Leftrightarrow \delta = 1 \text{ xor } \delta \frac{1}{1-\delta} > 1, \delta \neq 0 \quad \Leftrightarrow \quad \delta > 1 - \delta$$

# Cost of deviating in Round N

So if

$$-2\delta^N + 2 \sum_{t=N+1}^{\infty} \delta^t > 0$$

there is forfeit from payoff. This is when

$$\sum_{t=N+1}^{\infty} \delta^t > \delta^N$$

$$\Leftrightarrow \delta^{N+1} \sum_{t=0}^{\infty} \delta^t > \delta^N \quad \Leftrightarrow \quad \delta = 1 \text{ xor } \delta^{N+1} \frac{1}{1-\delta} > \delta^N$$

$$\Leftrightarrow \delta = 1 \text{ xor } \delta \frac{1}{1-\delta} > 1, \delta \neq 0 \quad \Leftrightarrow \quad \delta > 1 - \delta \quad \Leftrightarrow \quad \delta > \frac{1}{2}.$$

# Cost of deviating in Round N

So if

$$-2\delta^N + 2 \sum_{t=N+1}^{\infty} \delta^t > 0$$

there is forfeit from payoff. This is when

$$\sum_{t=N+1}^{\infty} \delta^t > \delta^N$$

$$\Leftrightarrow \delta^{N+1} \sum_{t=0}^{\infty} \delta^t > \delta^N \quad \Leftrightarrow \quad \delta = 1 \text{ xor } \delta^{N+1} \frac{1}{1-\delta} > \delta^N$$

$$\Leftrightarrow \delta = 1 \text{ xor } \delta \frac{1}{1-\delta} > 1, \delta \neq 0 \quad \Leftrightarrow \quad \delta > 1 - \delta \quad \Leftrightarrow \quad \delta > \frac{1}{2}.$$

Therefore, if  $\delta > 1/2$  every player forfeits payoff by deviating from  $T$ .  $\square$

# Example: an alternating trigger strategy

Yet another subgame perfect equilibrium:

# Example: an alternating trigger strategy

Yet another subgame perfect equilibrium:

**Informal definition of strategies.**  $A$  and  $B$  tacitly agree to alternate actions, i.e.,

$$(C, D), (D, C), (C, D), (D, C) \dots$$

# Example: an alternating trigger strategy

Yet another subgame perfect equilibrium:

**Informal definition of strategies.**  $A$  and  $B$  tacitly agree to alternate actions, i.e.,

$$(C, D), (D, C), (C, D), (D, C) \dots$$

If one of them deviates, the other party plays  $D$  forever.

# Example: an alternating trigger strategy

Yet another subgame perfect equilibrium:

**Informal definition of strategies.**  $A$  and  $B$  tacitly agree to alternate actions, i.e.,

$$(C, D), (D, C), (C, D), (D, C) \dots$$

If one of them deviates, the other party plays  $D$  forever. (The party who originally deviated plays  $D$  forever thereafter as well.)

# Example: an alternating trigger strategy

Yet another subgame perfect equilibrium:

**Informal definition of strategies.**  $A$  and  $B$  tacitly agree to alternate actions, i.e.,

$$(C, D), (D, C), (C, D), (D, C) \dots$$

If one of them deviates, the other party plays  $D$  forever. (The party who originally deviated plays  $D$  forever thereafter as well.) **CKR** is at work here!



# Example: an alternating trigger strategy

Yet another subgame perfect equilibrium:

**Informal definition of strategies.**  $A$  and  $B$  tacitly agree to alternate actions, i.e.,

$$(C, D), (D, C), (C, D), (D, C) \dots$$

If one of them deviates, the other party plays  $D$  forever. (The party who originally deviated plays  $D$  forever thereafter as well.) **CKR** is at work here!

Let  $A$  be the strategy that plays  $C$  in Round 1. Let  $B$  be the other strategy.

# Example: an alternating trigger strategy

Yet another subgame perfect equilibrium:

**Informal definition of strategies.**  $A$  and  $B$  tacitly agree to alternate actions, i.e.,

$$(C, D), (D, C), (C, D), (D, C) \dots$$

If one of them deviates, the other party plays  $D$  forever. (The party who originally deviated plays  $D$  forever thereafter as well.) **CKR** is at work here!

Let  $A$  be the strategy that plays  $C$  in Round 1. Let  $B$  be the other strategy.

**Claim.** *The strategy profile  $(A, B)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

# Example: an alternating trigger strategy

Yet another subgame perfect equilibrium:

**Informal definition of strategies.**  $A$  and  $B$  tacitly agree to alternate actions, i.e.,

$$(C, D), (D, C), (C, D), (D, C) \dots$$

If one of them deviates, the other party plays  $D$  forever. (The party who originally deviated plays  $D$  forever thereafter as well.) **CKR** is at work here!

Let  $A$  be the strategy that plays  $C$  in Round 1. Let  $B$  be the other strategy.

**Claim.** *The strategy profile  $(A, B)$  is a subgame perfect equilibrium in  $G^*(\delta)$ , provided the probability of continuation,  $\delta$ , is sufficiently large.*

An analysis of this situation and a proof of this claim can be found in (Peters, 2008), pp. 104-105.\*

\*H. Peters (2008): *Game Theory: A Multi-Leveled Approach*. Springer, ISBN: 978-3-540-69290-4.

# Generalisation of trigger strategies

# Generalisation of trigger strategies

- **Every** convex combination<sup>2</sup> of payoffs

$$\alpha_1(3,3) + \alpha_2(0,5) + \alpha_3(5,0) + \alpha_4(1,1)$$

can be established by smartly picking appropriate strategy patterns.

# Generalisation of trigger strategies

- **Every** convex combination<sup>2</sup> of payoffs

$$\alpha_1(3,3) + \alpha_2(0,5) + \alpha_3(5,0) + \alpha_4(1,1)$$

can be established by smartly picking appropriate strategy patterns.  
E.g.: “We play 4 times  $(C,C)$ . Then we play 7 times  $(C,D), (D,C)$ , then 4 times  $(C,C)$  and so on”.

# Generalisation of trigger strategies

- **Every** convex combination<sup>2</sup> of payoffs

$$\alpha_1(3,3) + \alpha_2(0,5) + \alpha_3(5,0) + \alpha_4(1,1)$$

can be established by smartly picking appropriate strategy patterns.  
E.g.: “We play 4 times  $(C,C)$ . Then we play 7 times  $(C,D), (D,C)$ , then 4 times  $(C,C)$  and so on”.

- Ensure that  $(C,C)$  occurs (in the long run) in  $\alpha_1$ ,  $(C,D)$  in  $\alpha_2$ ,  $(D,C)$  in  $\alpha_3$ , and  $(D,D)$  in  $\alpha_4$  percent of the stages.

# Generalisation of trigger strategies

- **Every** convex combination<sup>2</sup> of payoffs

$$\alpha_1(3,3) + \alpha_2(0,5) + \alpha_3(5,0) + \alpha_4(1,1)$$

can be established by smartly picking appropriate strategy patterns.  
E.g.: “We play 4 times  $(C,C)$ . Then we play 7 times  $(C,D), (D,C)$ , then 4 times  $(C,C)$  and so on”.

- Ensure that  $(C,C)$  occurs (in the long run) in  $\alpha_1$ ,  $(C,D)$  in  $\alpha_2$ ,  $(D,C)$  in  $\alpha_3$ , and  $(D,D)$  in  $\alpha_4$  percent of the stages.
- As long as these limiting average payoffs exceed  $\text{payoff}(\{D,D\})$  for each player (which is 1), associated trigger strategies can be formulated that lead to these payoffs and trigger eternal play of  $(D,D)$  after a deviation.



# Generalisation of trigger strategies

- **Every** convex combination<sup>2</sup> of payoffs

$$\alpha_1(3,3) + \alpha_2(0,5) + \alpha_3(5,0) + \alpha_4(1,1)$$


can be established by smartly picking appropriate strategy patterns.  
E.g.: “We play 4 times  $(C,C)$ . Then we play 7 times  $(C,D), (D,C)$ , then 4 times  $(C,C)$  and so on”.

- Ensure that  $(C,C)$  occurs (in the long run) in  $\alpha_1$ ,  $(C,D)$  in  $\alpha_2$ ,  $(D,C)$  in  $\alpha_3$ , and  $(D,D)$  in  $\alpha_4$  percent of the stages.
- As long as these limiting average payoffs exceed  $\text{payoff}(\{D,D\})$  for each player (which is 1), associated trigger strategies can be formulated that lead to these payoffs and trigger eternal play of  $(D,D)$  after a deviation.
- For  $\delta$  high enough, these strategies again form a SGP NE.


---

<sup>2</sup>Meaning  $\alpha_i \geq 0$  and  $\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 = 1$ .

# Folk theorem for SGP NE in a repeated PD




# Folk theorem for SGP NE in a repeated PD



profiles).

1. **Feasible payoffs** (striped):  
payoff combos that can be  
obtained by jointly repeating  
patterns of actions (more  
accurately: patterns of action

# Folk theorem for SGP NE in a repeated PD




1. **Feasible payoffs** (striped):  
payoff combos that can be  
obtained by jointly repeating  
patterns of actions (more  
accurately: patterns of action

profiles).

2. **Enforceable payoffs** (shaded):  
everyone resides above  
punishment **minmax**.

# Folk theorem for SGP NE in a repeated PD




1. **Feasible payoffs** (striped):  
payoff combos that can be  
obtained by jointly repeating  
patterns of actions (more  
accurately: patterns of action

profiles).

2. **Enforceable payoffs** (shaded):  
everyone resides above  
punishment **minmax**.

For every payoff pair  $(x, y)$   
in  $(1) \cap (2)$ ,

# Folk theorem for SGP NE in a repeated PD




1. **Feasible payoffs** (striped):  
payoff combos that can be  
obtained by jointly repeating  
patterns of actions (more  
accurately: patterns of action

profiles).

2. **Enforceable payoffs** (shaded):  
everyone resides above  
punishment **minmax**.

For every payoff pair  $(x, y)$   
in  $(1) \cap (2)$ , there is a  
 $\delta(x, y) \in (0, 1)$ ,

# Folk theorem for SGP NE in a repeated PD




1. **Feasible payoffs** (striped):  
payoff combos that can be  
obtained by jointly repeating  
patterns of actions (more  
accurately: patterns of action

profiles).

2. **Enforceable payoffs** (shaded):  
everyone resides above  
punishment **minmax**.

For every payoff pair  $(x, y)$   
in  $(1) \cap (2)$ , there is a  
 $\delta(x, y) \in (0, 1)$ , such that for  
all  $\delta \geq \delta(x, y)$

# Folk theorem for SGP NE in a repeated PD



1. **Feasible payoffs** (striped):  
payoff combos that can be obtained by jointly repeating patterns of actions (more accurately: patterns of action

profiles).

2. **Enforceable payoffs** (shaded):  
everyone resides above punishment **minmax**.

For every payoff pair  $(x, y)$  in  $(1) \cap (2)$ , there is a  $\delta(x, y) \in (0, 1)$ , such that for all  $\delta \geq \delta(x, y)$  the payoff  $(x, y)$  can be obtained as the limiting average in a subgame perfect equilibrium of  $G^*(\delta)$ .



# **Part IV:**

## **non-SGP Nash equilibria**

# Existence of non-SGP Nash equilibria



# Existence of non-SGP Nash equilibria

- We have seen that many subgame perfect equilibria exist for repeated games.



# Existence of non-SGP Nash equilibria

- We have seen that many subgame perfect equilibria exist for repeated games.
- What about the **existence of non-SGP Nash equilibria in repeated games**, i.e., equilibria that are not necessarily subgame perfect?



# Existence of non-SGP Nash equilibria

- We have seen that many subgame perfect equilibria exist for repeated games.
- What about the **existence of non-SGP Nash equilibria in repeated games**, i.e., equilibria that are not necessarily subgame perfect?
- Without the requirement of subgame perfection, **deviations can be punished more severely**: the equilibrium does not have to induce SGPs.







# Existence of non-SGP Nash equilibria

- We have seen that many subgame perfect equilibria exist for repeated games.
- What about the **existence of non-SGP Nash equilibria in repeated games**, i.e., equilibria that are not necessarily subgame perfect?
- Without the requirement of subgame perfection, **deviations can be punished more severely**: the equilibrium does not have to induce SGPs.
- However, non-SGPs implies **threats that are not credible**.



# Game Theory: A Critical [what?]



# Example: A repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)



# Example: A repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	( $1, 1$ )	( $0, 0$ )
	Down ( $D$ )	( $0, 0$ )	( $-1, 4$ )

1. For row,  $U$  is a dominating strategy.

# Example: A repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

1. For row,  $U$  is a dominating strategy.
2. The pure profile  $(U, L)$  is the only mixed strategy profile that is a NE.

# Example: A repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

1. For row,  $U$  is a dominating strategy.
2. The pure profile  $(U, L)$  is the only mixed strategy profile that is a NE.
3. Define trigger-strategies  $(T1, T2)$  such that the pattern  $[(D, R), (U, L)^3]^*$  is played indefinitely.

# Example: A repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

1. For row,  $U$  is a dominating strategy.
2. The pure profile  $(U, L)$  is the only mixed strategy profile that is a NE.
3. Define trigger-strategies  $(T1, T2)$  such that the pattern  $[(D, R), (U, L)^3]^*$  is played indefinitely.

If this pattern is violated, both parties fall back to punishment strategies:

# Example: A repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

1. For row,  $U$  is a dominating strategy.
2. The pure profile  $(U, L)$  is the only mixed strategy profile that is a NE.
3. Define trigger-strategies  $(T1, T2)$  such that the pattern  $[(D, R), (U, L)^3]^*$  is played indefinitely.

If this pattern is violated, both parties fall back to punishment strategies:

- The punishment strategy of row is mixed  $(0.8, 0.2)^*$ .

# Example: A repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

1. For row,  $U$  is a dominating strategy.
2. The pure profile  $(U, L)$  is the only mixed strategy profile that is a NE.
3. Define trigger-strategies  $(T1, T2)$  such that the pattern  $[(D, R), (U, L)^3]^*$  is played indefinitely.

If this pattern is violated, both parties fall back to punishment strategies:

- The punishment strategy of row is mixed  $(0.8, 0.2)^*$ .
- The punishment strategy of col is pure  $R^*$ .

# Example: A repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

1. For row,  $U$  is a dominating strategy.
2. The pure profile  $(U, L)$  is the only mixed strategy profile that is a NE.
3. Define trigger-strategies  $(T1, T2)$  such that the pattern  $[(D, R), (U, L)^3]^*$  is played indefinitely.

If this pattern is violated, both parties fall back to punishment strategies:

- The punishment strategy of row is mixed  $(0.8, 0.2)^*$ .
- The punishment strategy of col is pure  $R^*$ .

This combination of strategies is **not** a NE.

# Example: A repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

1. For row,  $U$  is a dominating strategy.
2. The pure profile  $(U, L)$  is the only mixed strategy profile that is a NE.
3. Define trigger-strategies  $(T1, T2)$  such that the pattern  $[(D, R), (U, L)^3]^*$  is played indefinitely.

If this pattern is violated, both parties fall back to punishment strategies:

- The punishment strategy of row is mixed  $(0.8, 0.2)^*$ .
- The punishment strategy of col is pure  $R^*$ .

This combination of strategies is **not** a NE. (For  $R^*$  induces  $U^*$ .)



# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

**Claim.** *The combination of trigger strategies  $(T1, T2)$  is a Nash-equilibrium for large enough  $\delta \in [0, 1]$ .*

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

**Claim.** *The combination of trigger strategies  $(T1, T2)$  is a Nash-equilibrium for large enough  $\delta \in [0, 1]$ .*

- $T1 \Rightarrow T2$ . If row plays (the non-degenerated part of)  $T1$ , then col must play  $T2$ , for  $T2$  is a best response to  $T1$ .

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(−1, 4)

**Claim.** *The combination of trigger strategies  $(T1, T2)$  is a Nash-equilibrium for large enough  $\delta \in [0, 1]$ .*

- $T1 \Rightarrow T2$ . If row plays (the non-degenerated part of)  $T1$ , then col must play  $T2$ , for  $T2$  is a best response to  $T1$ .
- $T2 \Rightarrow T1$ . If at all, the best moment for row to deviate is at  $D$ , for that would give row an incidental advantage of 1. After that row's opponent falls back to  $R^*$ .

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

**Claim.** *The combination of trigger strategies  $(T1, T2)$  is a Nash-equilibrium for large enough  $\delta \in [0, 1]$ .*

- $T1 \Rightarrow T2$ . If row plays (the non-degenerated part of)  $T1$ , then col must play  $T2$ , for  $T2$  is a best response to  $T1$ .
- $T2 \Rightarrow T1$ . If at all, the best moment for row to deviate is at  $D$ , for that would give row an incidental advantage of 1. After that row's opponent falls back to  $R^*$ .

Total payoff for row: 0 (for cheating)

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

**Claim.** *The combination of trigger strategies  $(T1, T2)$  is a Nash-equilibrium for large enough  $\delta \in [0, 1]$ .*

- $T1 \Rightarrow T2$ . If row plays (the non-degenerated part of)  $T1$ , then col must play  $T2$ , for  $T2$  is a best response to  $T1$ .
- $T2 \Rightarrow T1$ . If at all, the best moment for row to deviate is at  $D$ , for that would give row an incidental advantage of 1. After that row's opponent falls back to  $R^*$ .

Total payoff for row: 0 (for cheating) + 0 +  $\dots$  + 0 (for being punished by col).

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

- $T2 \Rightarrow T1$  (continued). Total payoff for row player: 0 (for cheating) +  $0 + \dots + 0$  (for being punished by the column player).



# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( $L$ )	Right ( $R$ )
Row:	Up ( $U$ )	(1, 1)	(0, 0)
	Down ( $D$ )	(0, 0)	(-1, 4)

- $T2 \Rightarrow T1$  (continued). Total payoff for row player: 0 (for cheating) +  $0 + \dots + 0$  (for being punished by the column player). Payoff for row player if he was loyal:

$$(-1 + 1 \cdot \delta + 1 \cdot \delta^2 + 1 \cdot \delta^3)$$

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( <i>L</i> )	Right ( <i>R</i> )
Row:	Up ( <i>U</i> )	(1, 1)	(0, 0)
	Down ( <i>D</i> )	(0, 0)	(-1, 4)

- $T2 \Rightarrow T1$  (continued). Total payoff for row player: 0 (for cheating) +  $0 + \dots + 0$  (for being punished by the column player). Payoff for row player if he was loyal:

$$(-1 + 1 \cdot \delta + 1 \cdot \delta^2 + 1 \cdot \delta^3) + (-1 \cdot \delta^4 + 1 \cdot \delta^5 + 1 \cdot \delta^6 + 1 \cdot \delta^7)$$

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( <i>L</i> )	Right ( <i>R</i> )
Row:	Up ( <i>U</i> )	(1, 1)	(0, 0)
	Down ( <i>D</i> )	(0, 0)	(-1, 4)

- $T2 \Rightarrow T1$  (continued). Total payoff for row player: 0 (for cheating) +  $0 + \dots + 0$  (for being punished by the column player). Payoff for row player if he was loyal:

$$(-1 + 1 \cdot \delta + 1 \cdot \delta^2 + 1 \cdot \delta^3) + (-1 \cdot \delta^4 + 1 \cdot \delta^5 + 1 \cdot \delta^6 + 1 \cdot \delta^7) + \dots$$

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( <i>L</i> )	Right ( <i>R</i> )
Row:	Up ( <i>U</i> )	(1, 1)	(0, 0)
	Down ( <i>D</i> )	(0, 0)	(-1, 4)

- $T2 \Rightarrow T1$  (continued). Total payoff for row player: 0 (for cheating) +  $0 + \dots + 0$  (for being punished by the column player). Payoff for row player if he was loyal:

$$(-1 + 1 \cdot \delta + 1 \cdot \delta^2 + 1 \cdot \delta^3) + (-1 \cdot \delta^4 + 1 \cdot \delta^5 + 1 \cdot \delta^6 + 1 \cdot \delta^7) + \dots$$

$$= \sum_{k=0}^{\infty} \delta^k$$

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( <i>L</i> )	Right ( <i>R</i> )
Row:	Up ( <i>U</i> )	(1, 1)	(0, 0)
	Down ( <i>D</i> )	(0, 0)	(-1, 4)

- $T2 \Rightarrow T1$  (continued). Total payoff for row player: 0 (for cheating) +  $0 + \dots + 0$  (for being punished by the column player). Payoff for row player if he was loyal:

$$(-1 + 1 \cdot \delta + 1 \cdot \delta^2 + 1 \cdot \delta^3) + (-1 \cdot \delta^4 + 1 \cdot \delta^5 + 1 \cdot \delta^6 + 1 \cdot \delta^7) + \dots$$

$$= \sum_{k=0}^{\infty} \delta^k - 2 \sum_{k=0}^{\infty} \delta^{4k}$$

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( <i>L</i> )	Right ( <i>R</i> )
Row:	Up ( <i>U</i> )	(1, 1)	(0, 0)
	Down ( <i>D</i> )	(0, 0)	(-1, 4)

- $T2 \Rightarrow T1$  (continued). Total payoff for row player: 0 (for cheating) +  $0 + \dots + 0$  (for being punished by the column player). Payoff for row player if he was loyal:

$$(-1 + 1 \cdot \delta + 1 \cdot \delta^2 + 1 \cdot \delta^3) + (-1 \cdot \delta^4 + 1 \cdot \delta^5 + 1 \cdot \delta^6 + 1 \cdot \delta^7) + \dots$$

$$= \sum_{k=0}^{\infty} \delta^k - 2 \sum_{k=0}^{\infty} \delta^{4k} = \frac{1}{1 - \delta}$$

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( <i>L</i> )	Right ( <i>R</i> )
Row:	Up ( <i>U</i> )	(1, 1)	(0, 0)
	Down ( <i>D</i> )	(0, 0)	(-1, 4)

- $T2 \Rightarrow T1$  (continued). Total payoff for row player: 0 (for cheating) +  $0 + \dots + 0$  (for being punished by the column player). Payoff for row player if he was loyal:

$$(-1 + 1 \cdot \delta + 1 \cdot \delta^2 + 1 \cdot \delta^3) + (-1 \cdot \delta^4 + 1 \cdot \delta^5 + 1 \cdot \delta^6 + 1 \cdot \delta^7) + \dots$$

$$= \sum_{k=0}^{\infty} \delta^k - 2 \sum_{k=0}^{\infty} \delta^{4k} = \frac{1}{1-\delta} - 2 \frac{1}{1-\delta^4}.$$

# Example: a repeated game with a non-SGP NE

Some game		Col:	
		Left ( <i>L</i> )	Right ( <i>R</i> )
Row:	Up ( <i>U</i> )	(1, 1)	(0, 0)
	Down ( <i>D</i> )	(0, 0)	(-1, 4)

- $T2 \Rightarrow T1$  (continued). Total payoff for row player: 0 (for cheating) +  $0 + \dots + 0$  (for being punished by the column player). Payoff for row player if he was loyal:

$$(-1 + 1 \cdot \delta + 1 \cdot \delta^2 + 1 \cdot \delta^3) + (-1 \cdot \delta^4 + 1 \cdot \delta^5 + 1 \cdot \delta^6 + 1 \cdot \delta^7) + \dots$$

$$= \sum_{k=0}^{\infty} \delta^k - 2 \sum_{k=0}^{\infty} \delta^{4k} = \frac{1}{1-\delta} - 2 \frac{1}{1-\delta^4}.$$

This expression is positive only if  $\delta \geq 0.54$ . (Solve 3rd-degree equation.)



# Example: A repeated game with a non-SGP NE

Col:	$L$	$R$
Row: $U$	$(1, 1)$	$(0, 0)$
$D$	$(0, 0)$	$(-1, 4)$

# Example: A repeated game with a non-SGP NE

Col:	$L$	$R$
Row: $U$	$(1, 1)$	$(0, 0)$
$D$	$(0, 0)$	$(-1, 4)$

- Col can punish row maximally by playing  $R^*$ . How row can punish col is less obvious.

# Example: A repeated game with a non-SGP NE

Col:	$L$	$R$
Row: $U$	$(1, 1)$	$(0, 0)$
$D$	$(0, 0)$	$(-1, 4)$

- Col can punish row maximally by playing  $R^*$ . How row can punish col is less obvious.
- If row plays  $D^*$  then col will play  $R^*$ . If row plays  $U^*$ , then col will play  $L^*$ .

# Example: A repeated game with a non-SGP NE

Col:	$L$	$R$
Row: $U$	$(1, 1)$	$(0, 0)$
$D$	$(0, 0)$	$(-1, 4)$

- Col can punish row maximally by playing  $R^*$ . How row can punish col is less obvious.
- If row plays  $D^*$  then col will play  $R^*$ . If row plays  $U^*$ , then col will play  $L^*$ .
- Row can punish col even more by playing a **minmax** strategy.

# Example: A repeated game with a non-SGP NE

Col:	L	R
Row: U	(1, 1)	(0, 0)
D	(0, 0)	(-1, 4)

expected payoff by choosing the right mix  $(l, r)$ :

$$\max_{l,r} ul \cdot 1 + dr \cdot 4$$

- Col can punish row maximally by playing  $R^*$ . How row can punish col is less obvious.
- If row plays  $D^*$  then col will play  $R^*$ . If row plays  $U^*$ , then col will play  $L^*$ .
- Row can punish col even more by playing a **minmax** strategy.  
Given row's mixed strategy  $(u, d)$ , col maximises his

# Example: A repeated game with a non-SGP NE

Col:	L	R
Row: U	(1, 1)	(0, 0)
D	(0, 0)	(-1, 4)

- Col can punish row maximally by playing  $R^*$ . How row can punish col is less obvious.
- If row plays  $D^*$  then col will play  $R^*$ . If row plays  $U^*$ , then col will play  $L^*$ .
- Row can punish col even more by playing a **minmax** strategy.  
Given row's mixed strategy  $(u, d)$ , col maximises his

expected payoff by choosing the right mix  $(l, r)$ :

$$\max_{l,r} ul \cdot 1 + dr \cdot 4$$

$$= \max_l ul + 4(1 - u)(1 - l)$$

# Example: A repeated game with a non-SGP NE

Col:	L	R
Row: U	(1, 1)	(0, 0)
D	(0, 0)	(-1, 4)

- Col can punish row maximally by playing  $R^*$ . How row can punish col is less obvious.
- If row plays  $D^*$  then col will play  $R^*$ . If row plays  $U^*$ , then col will play  $L^*$ .
- Row can punish col even more by playing a **minmax** strategy.  
Given row's mixed strategy

$(u, d)$ , col maximises his expected payoff by choosing the right mix  $(l, r)$ :

$$\max_{l,r} ul \cdot 1 + dr \cdot 4$$

$$= \max_l ul + 4(1 - u)(1 - l)$$

$$= \max_l (5u - 4)l + 4 - 4u.$$

# Example: A repeated game with a non-SGP NE

Col:	L	R
Row: U	(1, 1)	(0, 0)
D	(0, 0)	(-1, 4)

- Col can punish row maximally by playing  $R^*$ . How row can punish col is less obvious.
- If row plays  $D^*$  then col will play  $R^*$ . If row plays  $U^*$ , then col will play  $L^*$ .
- Row can punish col even more by playing a **minmax** strategy. Given row's mixed strategy

$(u, d)$ , col maximises his expected payoff by choosing the right mix  $(l, r)$ :

$$\max_{l,r} ul \cdot 1 + dr \cdot 4$$

$$= \max_l ul + 4(1 - u)(1 - l)$$


$$= \max_l (5u - 4)l + 4 - 4u.$$

- If  $5u - 4 = 0$ , it does not matter what col chooses for  $l$ —his expected payoff is always  $4 - 4(4/5) = 4/5$ .




# Example: A repeated game with a non-SGP NE

Draw a picture of the payoff surface of col.




# Example: A repeated game with a non-SGP NE



Draw a picture of the payoff surface of col.

- If  $5u - 4 = 0$ , it does not matter what col chooses for  $l$ . He expects  $4/5$ .


# Example: A repeated game with a non-SGP NE



Draw a picture of the payoff surface of col.

- If  $5u - 4 = 0$ , it does not matter what col chooses for  $l$ . He expects  $4/5$ .
- If  $5u - 4 > 0$  col will play  $l = 1$ , and expects to earn  $u$  and  $u > 4/5$ .


# Example: A repeated game with a non-SGP NE



Draw a picture of the payoff surface of col.

- If  $5u - 4 = 0$ , it does not matter what col chooses for  $l$ . He expects  $4/5$ .
- If  $5u - 4 > 0$  col will play  $l = 1$ , and expects to earn  $u$  and  $u > 4/5$ .
- If  $5u - 4 < 0$  col will play  $l = 0$ , and expects to earn  $4 - 4u > 4 - 4(4/5) = 4/5$ .

# Example: A repeated game with a non-SGP NE



Draw a picture of the payoff surface of col.

- If  $5u - 4 = 0$ , it does not matter what col chooses for  $l$ . He expects  $4/5$ .
- If  $5u - 4 > 0$  col will play  $l = 1$ , and expects to earn  $u$  and  $u > 4/5$ .
- If  $5u - 4 < 0$  col will play  $l = 0$ , and expects to earn  $4 - 4u > 4 - 4(4/5) = 4/5$ .

These calculations are done by hand, and do not easily generalise to higher dimensions.

# What next?

Now that we know that infinitely many equilibria exist in repeated games (an **embarrassment of richness**), there are a number of ways in which we may proceed.

# What next?

Now that we know that infinitely many equilibria exist in repeated games (an **embarrassment of richness**), there are a number of ways in which we may proceed.

- **Reinforcement Learning.** Agents simply execute the action(s) with maximal rewards in the past.

# What next?

Now that we know that infinitely many equilibria exist in repeated games (an **embarrassment of richness**), there are a number of ways in which we may proceed.

- **Reinforcement Learning.** Agents simply execute the action(s) with maximal rewards in the past.
- **No-regret learning.** Agents execute the action(s) with maximal virtual rewards in the past.



# What next?

Now that we know that infinitely many equilibria exist in repeated games (an **embarrassment of richness**), there are a number of ways in which we may proceed.

- **Reinforcement Learning.** Agents simply execute the action(s) with maximal rewards in the past.
- **No-regret learning.** Agents execute the action(s) with maximal virtual rewards in the past.
- **Fictitious Play.** Sample the actions of opponent(s) and play a best response.

# What next?

Now that we know that infinitely many equilibria exist in repeated games (an **embarrassment of richness**), there are a number of ways in which we may proceed.

- **Reinforcement Learning.** Agents simply execute the action(s) with maximal rewards in the past.
- **No-regret learning.** Agents execute the action(s) with maximal virtual rewards in the past.
- **Fictitious Play.** Sample the actions of opponent(s) and play a best response.
- **Gradient Dynamics.** This is to approximate NE of single-shot games (stage games) through gradient ascent (hill-climbing).