

Multi-agent learning

Bayesian play

Gerard Vreeswijk, Intelligent Software Systems, Computer Science
Department, Faculty of Sciences, Utrecht University, The
Netherlands.

Thursday 21st May, 2020

Plan for today

Preparation

Plan for today

Preparation

- Bayes' rule.

Plan for today

Preparation

- Bayes' rule.
- Examples of Bayesian play.

Plan for today

Preparation

- Bayes' rule.
- Examples of Bayesian play.
- Demo.

Plan for today

Preparation

- Bayes' rule.
- Examples of Bayesian play.
- Demo.

Formalism

Plan for today

Preparation

- Bayes' rule.
- Examples of Bayesian play.
- Demo.

Formalism

- Jordan's framework: reply rule
+ forecasting rule = predictive
learning rule.

Plan for today

Preparation

- Bayes' rule.
- Examples of Bayesian play.
- Demo.

Formalism

- Jordan's framework: reply rule + forecasting rule = predictive learning rule.
- Beliefs about reply rules of other players, Kuhn's result.

Plan for today

Preparation

- Bayes' rule.
- Examples of Bayesian play.
- Demo.

Formalism

- Jordan's framework: reply rule + forecasting rule = predictive learning rule.
- Beliefs about reply rules of other players, Kuhn's result.
- True distribution of play vs. subjective distribution of play.

Plan for today

Preparation

- Bayes' rule.
- Examples of Bayesian play.
- Demo.

Formalism

- Jordan's framework: reply rule + forecasting rule = predictive learning rule.
- Beliefs about reply rules of other players, Kuhn's result.
- True distribution of play vs. subjective distribution of play.

Main results

Plan for today

Preparation

- Bayes' rule.
- Examples of Bayesian play.
- Demo.

Formalism

- Jordan's framework: reply rule + forecasting rule = predictive learning rule.
- Beliefs about reply rules of other players, Kuhn's result.
- True distribution of play vs. subjective distribution of play.

Main results

- Domination of measures (a.k.a. absolute continuity).

Plan for today

Preparation

- Bayes' rule.
- Examples of Bayesian play.
- Demo.

Formalism

- Jordan's framework: reply rule + forecasting rule = predictive learning rule.
- Beliefs about reply rules of other players, Kuhn's result.
- True distribution of play vs. subjective distribution of play.

Main results

- Domination of measures (a.k.a. absolute continuity).
- Theorem of Blackwell and Dubins (1992).

Plan for today

Preparation

- Bayes' rule.
- Examples of Bayesian play.
- Demo.

Formalism

- Jordan's framework: reply rule + forecasting rule = predictive learning rule.
- Beliefs about reply rules of other players, Kuhn's result.
- True distribution of play vs. subjective distribution of play.

Main results

- Domination of measures (a.k.a. absolute continuity).
- Theorem of Blackwell and Dubins (1992).
- Notion of ϵ -closeness.

Plan for today

Preparation

- Bayes' rule.
- Examples of Bayesian play.
- Demo.

Formalism

- Jordan's framework: reply rule + forecasting rule = predictive learning rule.
- Beliefs about reply rules of other players, Kuhn's result.
- True distribution of play vs. subjective distribution of play.

Main results

- Domination of measures (a.k.a. absolute continuity).
- Theorem of Blackwell and Dubins (1992).
- Notion of ϵ -closeness.
- Theorem of Kalai and Lehrer (1993):

Plan for today

Preparation

- Bayes' rule.
- Examples of Bayesian play.
- Demo.

Formalism

- Jordan's framework: reply rule + forecasting rule = predictive learning rule.
- Beliefs about reply rules of other players, Kuhn's result.
- True distribution of play vs. subjective distribution of play.

Main results

- Domination of measures (a.k.a. absolute continuity).
- Theorem of Blackwell and Dubins (1992).
- Notion of ϵ -closeness.
- Theorem of Kalai and Lehrer (1993): If a player gives all potential play paths a small positive probability ("grain of truth"), then, eventually, his/her subjective beliefs will be ϵ -close to the actual realisation of play.

Key publication

Kalai & Lehrer (1993). "Rational learning leads to Nash equilibrium". *Econometrica*, Vol. **61**, No. 5, pp 1019-1045.

Key publication

Kalai & Lehrer (1993). “Rational learning leads to Nash equilibrium”. *Econometrica*, Vol. **61**, No. 5, pp 1019-1045.

Scholarly resources

Young (2004): *Strategic Learning and its Limits*, Oxford UP. Ch. 7: “Bayesian Learning”.

Shoham *et al.* (2009): *Multi-agent Systems*. Ch. 7: “Learning and Teaching”. Sec. 7.3: “Rational Learning”.

Key publication

Kalai & Lehrer (1993). “Rational learning leads to Nash equilibrium”. *Econometrica*, Vol. 61, No. 5, pp 1019-1045.

Scholarly resources

Young (2004): *Strategic Learning and its Limits*, Oxford UP. Ch. 7: “Bayesian Learning”.

Shoham *et al.* (2009): *Multi-agent Systems*. Ch. 7: “Learning and Teaching”. Sec. 7.3: “Rational Learning”.

Practical computer science / AI application

Zeng & Sycara (1996): *Bayesian Learning in Negotiation* in: Working Notes of the AAAI Spring Symposium on Adaptation, Co-Evolution and Learning in Multiagent Systems, Stanford, CA.

Part I:

Elementary probability and Bayes' theorem

Conditional probability

Conditional probability

- If $\Pr\{F\} \neq 0$ then the conditional probability given F is defined as

Conditional probability

- If $\Pr\{F\} \neq 0$ then the conditional probability given F is defined as

$$\Pr\{E|F\}$$

Conditional probability

- If $\Pr\{F\} \neq 0$ then the conditional probability given F is defined as

$$\Pr\{E|F\} =_{Def}$$

Conditional probability

- If $\Pr\{F\} \neq 0$ then the conditional probability given F is defined as

$$\Pr\{E|F\} =_{Def} \Pr\{EF\}$$

Conditional probability

- If $\Pr\{F\} \neq 0$ then the conditional probability given F is defined as

$$\Pr\{E|F\} =_{Def} \frac{\Pr\{EF\}}{\Pr\{F\}}$$

Conditional probability

- If $\Pr\{F\} \neq 0$ then the conditional probability given F is defined as

$$\Pr\{E|F\} =_{Def} \frac{\Pr\{EF\}}{\Pr\{F\}}.$$

Conditional probability

- If $\Pr\{F\} \neq 0$ then the conditional probability given F is defined as

$$\Pr\{E|F\} =_{Def} \frac{\Pr\{EF\}}{\Pr\{F\}}.$$

It can be shown that $\Pr\{\cdot | F\}$ is a probability measure on F .

Conditional probability

- If $\Pr\{F\} \neq 0$ then the conditional probability given F is defined as

$$\Pr\{E|F\} =_{Def} \frac{\Pr\{EF\}}{\Pr\{F\}}.$$

It can be shown that $\Pr\{\cdot | F\}$ is a probability measure on F .

- Writing $\Pr\{EF\}$ as $\Pr\{F|E\}\Pr\{E\}$ yields Bayes' rule:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\Pr\{F\}}.$$

Conditional probability

- If $\Pr\{F\} \neq 0$ then the conditional probability given F is defined as

$$\Pr\{E|F\} =_{Def} \frac{\Pr\{EF\}}{\Pr\{F\}}.$$

It can be shown that $\Pr\{\cdot | F\}$ is a probability measure on F .

- Writing $\Pr\{EF\}$ as $\Pr\{F|E\}\Pr\{E\}$ yields Bayes' rule:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\Pr\{F\}}.$$

$\Pr\{E\}$ is called the prior probability of E .

Conditional probability

- If $\Pr\{F\} \neq 0$ then the conditional probability given F is defined as

$$\Pr\{E|F\} =_{Def} \frac{\Pr\{EF\}}{\Pr\{F\}}.$$

It can be shown that $\Pr\{\cdot | F\}$ is a probability measure on F .

- Writing $\Pr\{EF\}$ as $\Pr\{F|E\}\Pr\{E\}$ yields Bayes' rule:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\Pr\{F\}}.$$

$\Pr\{E\}$ is called the prior probability of E .

Typical exercise: “given $\Pr\{E\}$, $\Pr\{F\}$, and $\Pr\{F|E\}$, compute $\Pr\{E|F\}$ ”. (E.g., E = “influenza”, F = “fever”.)

Conditional probability

- If $\Pr\{F\} \neq 0$ then the conditional probability given F is defined as

$$\Pr\{E|F\} =_{Def} \frac{\Pr\{EF\}}{\Pr\{F\}}.$$

It can be shown that $\Pr\{\cdot|F\}$ is a probability measure on F .

- Writing $\Pr\{EF\}$ as $\Pr\{F|E\}\Pr\{E\}$ yields Bayes' rule:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\Pr\{F\}}.$$

$\Pr\{E\}$ is called the prior probability of E .

Typical exercise: “given $\Pr\{E\}$, $\Pr\{F\}$, and $\Pr\{F|E\}$, compute $\Pr\{E|F\}$ ”. (E.g., E = “influenza”, F = “fever”.)

- It is customary to marginalise the denominator through E as well:

$$\Pr\{F\} = \Pr\{F|E\}\Pr\{E\} + \Pr\{F|E^c\}\Pr\{E^c\},$$

Conditional probability

- If $\Pr\{F\} \neq 0$ then the conditional probability given F is defined as

$$\Pr\{E|F\} =_{Def} \frac{\Pr\{EF\}}{\Pr\{F\}}.$$

It can be shown that $\Pr\{\cdot | F\}$ is a probability measure on F .

- Writing $\Pr\{EF\}$ as $\Pr\{F|E\}\Pr\{E\}$ yields Bayes' rule:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\Pr\{F\}}.$$

$\Pr\{E\}$ is called the prior probability of E .

Typical exercise: “given $\Pr\{E\}$, $\Pr\{F\}$, and $\Pr\{F|E\}$, compute $\Pr\{E|F\}$ ”. (E.g., E = “influenza”, F = “fever”.)

- It is customary to marginalise the denominator through E as well:

$$\Pr\{F\} = \Pr\{F|E\}\Pr\{E\} + \Pr\{F|E^c\}\Pr\{E^c\},$$

so that

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\Pr\{F|E\}\Pr\{E\} + \Pr\{F|E^c\}\Pr\{E^c\}}.$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\}$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} =$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\} \Pr\{E\}}{\Pr\{F\}}$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\Pr\{F\}}$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\Pr\{F\}}$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int}$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}}$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}\Pr\{E\}}$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}\Pr\{E\}dE}$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}\Pr\{E\}dE}$$

$$f_X(x|Y = y)$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}\Pr\{E\}dE}$$

$$f_X(x|Y = y) = \frac{f_{XY}(x, y)}{f_Y(y)}$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}\Pr\{E\}dE}$$

$$f_X(x|Y = y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{f_Y(y|X = x) f_X(x)}{f_Y(y)}$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}\Pr\{E\}dE}$$

$$f_X(x|Y = y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{f_Y(y|X = x) f_X(x)}{f_Y(y)} = \frac{f_Y(y|X = x) f_X(x)}{\int f_Y(y|X = x) f_X(x) dx}.$$

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}\Pr\{E\}dE}$$

$$f_X(x|Y=y) = \frac{f_{XY}(x,y)}{f_Y(y)} = \frac{f_Y(y|X=x) f_X(x)}{f_Y(y)} = \frac{f_Y(y|X=x) f_X(x)}{\int f_Y(y|X=x) f_X(x) dx}.$$

As in the discrete case, these terms have standard names.

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}\Pr\{E\}dE}$$

$$f_X(x|Y=y) = \frac{f_{XY}(x,y)}{f_Y(y)} = \frac{f_Y(y|X=x) f_X(x)}{f_Y(y)} = \frac{f_Y(y|X=x) f_X(x)}{\int f_Y(y|X=x) f_X(x) dx}.$$

As in the discrete case, these terms have standard names.

■ $f_{XY}(x,y)$ is the **joint density** of X and Y .

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}\Pr\{E\}dE}$$

$$f_X(x|Y=y) = \frac{f_{XY}(x,y)}{f_Y(y)} = \frac{f_Y(y|X=x) f_X(x)}{f_Y(y)} = \frac{f_Y(y|X=x) f_X(x)}{\int f_Y(y|X=x) f_X(x) dx}.$$

As in the discrete case, these terms have standard names.

- $f_{XY}(x,y)$ is the **joint density** of X and Y .
- $f_X(x)$ is the **prior density** of X .

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}\Pr\{E\}dE}$$

$$f_X(x|Y = y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{f_Y(y|X = x) f_X(x)}{f_Y(y)} = \frac{f_Y(y|X = x) f_X(x)}{\int f_Y(y|X = x) f_X(x) dx}.$$

As in the discrete case, these terms have standard names.

- $f_{XY}(x, y)$ is the **joint density** of X and Y .
- $f_X(x)$ is the **prior density** of X .
- $f_X(x|Y = y)$ is the **posterior density** of X given $Y = y$.

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}\Pr\{E\}dE}$$

$$f_X(x|Y = y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{f_Y(y|X = x) f_X(x)}{f_Y(y)} = \frac{f_Y(y|X = x) f_X(x)}{\int f_Y(y|X = x) f_X(x) dx}.$$

As in the discrete case, these terms have standard names.

- $f_{XY}(x, y)$ is the **joint density** of X and Y .
- $f_X(x)$ is the **prior density** of X .
- $f_X(x|Y = y)$ is the **posterior density** of X given $Y = y$.
- $f_Y(y|X = x)$ is the **posterior density** of Y given $X = x$.

Bayes' theorem for continuous random variables

Bayes' theorem for continuous random variables:

$$\Pr\{E|F\} = \frac{\Pr\{F|E\}\Pr\{E\}}{\int \Pr\{F|E\}\Pr\{E\}dE}$$

$$f_X(x|Y=y) = \frac{f_{XY}(x,y)}{f_Y(y)} = \frac{f_Y(y|X=x) f_X(x)}{f_Y(y)} = \frac{f_Y(y|X=x) f_X(x)}{\int f_Y(y|X=x) f_X(x) dx}.$$

As in the discrete case, these terms have standard names.

- $f_{XY}(x,y)$ is the **joint density** of X and Y .
- $f_X(x)$ is the **prior density** of X .
- $f_X(x|Y=y)$ is the **posterior density** of X given $Y=y$.
- $f_Y(y|X=x)$ is the **posterior density** of Y given $X=x$.
- $f_X(x)$ and $f_Y(y)$ are **marginal densities** of X and Y .

Part Ia: Bayes' rule applied to guess opponent's reply rules

Use Bayes' rule to compute $\Pr\{g = g_i \mid h\}$ for every i

Use Bayes' rule to compute $\Pr\{g = g_i \mid h\}$ for every i

- Let $G = \{g_1, \dots, g_n\}$
denote possible
reply rules of the
opponent.

Use Bayes' rule to compute $\Pr\{g = g_i \mid h\}$ for every i

- Let $G = \{g_1, \dots, g_n\}$ denote possible reply rules of the opponent.
- Let g denote the actual reply rule of the opponent.

Use Bayes' rule to compute $\Pr\{g = g_i \mid h\}$ for every i

- Let $G = \{g_1, \dots, g_n\}$ denote possible reply rules of the opponent.
- Let g denote the actual reply rule of the opponent.
- Give every event $g = g_i$ the same prior probability $\Pr\{g = g_i\} = 1/n$.

Use Bayes' rule to compute $\Pr\{g = g_i \mid h\}$ for every i

- Let $G = \{g_1, \dots, g_n\}$ denote possible reply rules of the opponent.
- Let g denote the actual reply rule of the opponent.
- Give every event $g = g_i$ the same prior probability $\Pr\{g = g_i\} = 1/n$.
- We'd like to know $\Pr\{g = g_i \mid h\}$ for every i .

Use Bayes' rule to compute $\Pr\{g = g_i \mid h\}$ for every i

- Let $G = \{g_1, \dots, g_n\}$ denote possible reply rules of the opponent.
- Bayes' rule for $g = g_i$'s posterior probability:
$$\Pr\{g = g_i \mid h\} = \frac{\Pr(h \mid g = g_i)\Pr\{g = g_i\}}{\Pr\{h\}}.$$
- Let g denote the actual reply rule of the opponent.
- Give every event $g = g_i$ the same prior probability
$$\Pr\{g = g_i\} = 1/n.$$
- We'd like to know $\Pr\{g = g_i \mid h\}$ for every i .

Use Bayes' rule to compute $\Pr\{g = g_i \mid h\}$ for every i

- Let $G = \{g_1, \dots, g_n\}$ denote possible reply rules of the opponent.
- Bayes' rule for $g = g_i$'s posterior probability:
$$\Pr\{g = g_i \mid h\} = \frac{\Pr(h \mid g = g_i)\Pr\{g = g_i\}}{\Pr\{h\}}.$$
- Let g denote the actual reply rule of the opponent.
- If h is marginalised by the events $g = g_i$
$$\Pr\{h\} = \sum_{j=1}^n \Pr\{h \mid g = g_j\}\Pr\{g = g_j\},$$
- Give every event $g = g_i$ the same prior probability
$$\Pr\{g = g_i\} = 1/n.$$
- We'd like to know $\Pr\{g = g_i \mid h\}$ for every i .

Use Bayes' rule to compute $\Pr\{g = g_i \mid h\}$ for every i

- Let $G = \{g_1, \dots, g_n\}$ denote possible reply rules of the opponent.
- Bayes' rule for $g = g_i$'s posterior probability:
$$\Pr\{g = g_i \mid h\} = \frac{\Pr(h \mid g = g_i)\Pr\{g = g_i\}}{\Pr\{h\}}.$$

- Let g denote the actual reply rule of the opponent.
- If h is marginalised by the events $g = g_i$

$$\Pr\{h\} = \sum_{j=1}^n \Pr\{h \mid g = g_j\}\Pr\{g = g_j\},$$

- Give every event $g = g_i$ the same prior probability

$$\Pr\{g = g_i\} = 1/n.$$

- We'd like to know $\Pr\{g = g_i \mid h\}$ for every i .

we obtain

$$\Pr\{g = g_i \mid h\} = \frac{\Pr\{h \mid g = g_i\}\Pr\{g = g_i\}}{\sum_{j=1}^n \Pr\{h \mid g = g_j\}\Pr\{g = g_j\}}$$

Use Bayes' rule to compute $\Pr\{g = g_i \mid h\}$ for every i

- Let $G = \{g_1, \dots, g_n\}$ denote possible reply rules of the opponent.
- Bayes' rule for $g = g_i$'s posterior probability:
$$\Pr\{g = g_i \mid h\} = \frac{\Pr(h \mid g = g_i)\Pr\{g = g_i\}}{\Pr\{h\}}.$$

- Let g denote the actual reply rule of the opponent.

- Give every event $g = g_i$ the same prior probability
 $\Pr\{g = g_i\} = 1/n.$

- We'd like to know $\Pr\{g = g_i \mid h\}$ for every i .

- If h is marginalised by the events $g = g_i$

$$\Pr\{h\} = \sum_{j=1}^n \Pr\{h \mid g = g_j\}\Pr\{g = g_j\},$$

we obtain

$$\Pr\{g = g_i \mid h\} = \frac{\Pr\{h \mid g = g_i\}\Pr\{g = g_i\}}{\sum_{j=1}^n \Pr\{h \mid g = g_j\}\Pr\{g = g_j\}}$$

- Because of identical priors and normalization, effectively
 $\Pr\{g = g_i \mid h\} \propto \Pr\{h \mid g = g_i\}.$

Part II: Demo and examples

Demo and examples

Demo and examples

1. **Demo.** Learn reactive reply rules, such as

- **All-C**: always cooperate.
- **Unforgiving** (“unforgiving”): cooperate until opponent defects, then defect forever.
- **C-90%**: cooperate 90% of the time (randomly).
- **Tit-for-tat**: mimic opponent’s moves.
- **Josh 10%**: play tit-for-tat 90% of the time, defect 10% of the time.
- **Majority**: respond with the action most played by the opponent.
- **Eatherly**: mirror the (projected) mixed strategy of the opponent.
- ...


Demo and examples

1. **Demo.** Learn reactive reply rules, such as


- **All-C**: always cooperate.
- **Unforgiving** (“unforgiving”): cooperate until opponent defects, then defect forever.
- **C-90%**: cooperate 90% of the time (randomly).
- **Tit-for-tat**: mimic opponent’s moves.
- **Josh 10%**: play tit-for-tat 90% of the time, defect 10% of the time.
- **Majority**: respond with the action most played by the opponent.
- **Eatherly**: mirror the (projected) mixed strategy of the opponent.
- ...

2. **Examples.** Learn reply rules in the repeated prisoners’ dilemma; learn reply rules in the coordination game.


Example: learning on reactive strategies




Example: learning on reactive strategies




Example: learning on reactive strategies




Example: learning on reactive strategies




Example: learning on reactive strategies




Example: learning on reactive strategies



Example: learning on reactive strategies




Example: learning on reactive strategies




Example: learning on reactive strategies




Example: learning on reactive strategies




Example: learning on reactive strategies




Example: learning on reactive strategies



Example: learning on reactive strategies



Example: learning on reactive strategies



Explanation of $\Pr\{h \mid s_2 = \text{Joss-10\%}\}$

Event	Description
$s_2 = \text{Joss-10\%}$	Player 2's strategy is Joss-10%
$X_1^{t-1} = C$	Player 1 cooperated in the previous round
$X_2^t = C$	Player 2 cooperates in the current round
ζ^t	Joss-10% randomises (hence defects)

Explanation of $\Pr\{h \mid s_2 = \text{Joss-10\%}\}$

Event	Description
$s_2 = \text{Joss-10\%}$	Player 2's strategy is Joss-10%
$X_1^{t-1} = C$	Player 1 cooperated in the previous round
$X_2^t = C$	Player 2 cooperates in the current round
ζ^t	Joss-10% randomises (hence defects)

Round:	1	2	3	4	5	6	7
Player 1:	C	C	C	D	D	C	C
Player 2 x (and h):	C	D	C	C	D	C	C
$\Pr\{X_2^t = x \mid s_2 = \text{Joss-10\%}\}:$	0.9	0.1	0.9	0.9	1	0	0
$\Pr\{h_2^t = h \mid s_2 = \text{Joss-10\%}\}:$	0.9	0.09	0.081	0.0729	0.0729	0	0

Explanation of $\Pr\{h \mid s_2 = \text{Joss-10\%}\}$

Event	Description
$s_2 = \text{Joss-10\%}$	Player 2's strategy is Joss-10%
$X_1^{t-1} = C$	Player 1 cooperated in the previous round
$X_2^t = C$	Player 2 cooperates in the current round
ζ^t	Joss-10% randomises (hence defects)

Round:	1	2	3	4	5	6	7
Player 1:	C	C	C	D	D	C	C
Player 2 x (and h):	C	D	C	C	D	C	C
$\Pr\{X_2^t = x \mid s_2 = \text{Joss-10\%}\}$:	0.9	0.1	0.9	0.9	1	0	0
$\Pr\{h_2^t = h \mid s_2 = \text{Joss-10\%}\}$:	0.9	0.09	0.081	0.0729	0.0729	0	0

$$\begin{aligned}
 \Pr\{h_2^t = C, h \mid X_1^{t-1} = C\} &= (\Pr\{X_2^t = C \mid X_1^{t-1} = C, \zeta^t\} \Pr\{\zeta^t\} + \\
 &\quad \Pr\{X_2^t = C \mid X_1^{t-1} = C, \bar{\zeta}^t\} \Pr\{\bar{\zeta}^t\}) \Pr\{h\} \\
 &= (0 \cdot 0.1 + 1 \cdot 0.9) \Pr\{h\} \\
 &= 0.9 \Pr\{h\}.
 \end{aligned}$$

Example: the disrupted unforgiving reply rule

Define

$g_t =_{Def}$ Play unforgiving before t , defect unconditionally at t and later.

Example: the disrupted unforgiving reply rule

Define

$g_t =_{Def}$ Play unforgiving before t , defect unconditionally at t and later.

Player 1 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Example: the disrupted unforgiving reply rule

Define

$g_t =_{Def}$ Play unforgiving before t , defect unconditionally at t and later.

Player 1 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round	g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
-------	-------	-------	-------	-------	-------	-------	---------	------------

Example: the disrupted unforgiving reply rule

Define

$g_t =_{Def}$ Play unforgiving before t , defect unconditionally at t and later.

Player 1 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round	g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
0.	1/4	1/8	1/16	1/32	1/64	1/128	\dots	1/2

Example: the disrupted unforgiving reply rule

Define

$g_t =_{Def}$ Play unforgiving before t , defect unconditionally at t and later.

Player 1 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
0.		$1/4$	$1/8$	$1/16$	$1/32$	$1/64$	$1/128$	\dots	$1/2$
1.	CC	0	$1/6$	$1/12$	$1/24$	$1/48$	$1/96$	\dots	$2/3$

Example: the disrupted unforgiving reply rule

Define

$g_t =_{Def}$ Play unforgiving before t , defect unconditionally at t and later.

Player 1 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
0.		$1/4$	$1/8$	$1/16$	$1/32$	$1/64$	$1/128$	\dots	$1/2$
1.	CC	0	$1/6$	$1/12$	$1/24$	$1/48$	$1/96$	\dots	$2/3$
2.	CC	0	0	$1/10$	$1/20$	$1/40$	$1/80$	\dots	$4/5$

Example: the disrupted unforgiving reply rule

Define

$g_t =_{Def}$ Play unforgiving before t , defect unconditionally at t and later.

Player 1 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
0.		$1/4$	$1/8$	$1/16$	$1/32$	$1/64$	$1/128$	\dots	$1/2$
1.	CC	0	$1/6$	$1/12$	$1/24$	$1/48$	$1/96$	\dots	$2/3$
2.	CC	0	0	$1/10$	$1/20$	$1/40$	$1/80$	\dots	$4/5$
3.	CC	0	0	0	$1/18$	$1/36$	$1/72$	\dots	$8/9$

Example: the disrupted unforgiving reply rule

Define

$g_t =_{Def}$ Play unforgiving before t , defect unconditionally at t and later.

Player 1 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
0.		$1/4$	$1/8$	$1/16$	$1/32$	$1/64$	$1/128$	\dots	$1/2$
1.	CC	0	$1/6$	$1/12$	$1/24$	$1/48$	$1/96$	\dots	$2/3$
2.	CC	0	0	$1/10$	$1/20$	$1/40$	$1/80$	\dots	$4/5$
3.	CC	0	0	0	$1/18$	$1/36$	$1/72$	\dots	$8/9$
4.	CC	0	0	0	0	$1/34$	$1/68$	\dots	$16/17$

Example: the disrupted unforgiving reply rule

Define

$g_t =_{Def}$ Play unforgiving before t , defect unconditionally at t and later.

Player 1 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
0.		1/4	1/8	1/16	1/32	1/64	1/128	\dots	1/2
1.	CC	0	1/6	1/12	1/24	1/48	1/96	\dots	2/3
2.	CC	0	0	1/10	1/20	1/40	1/80	\dots	4/5
3.	CC	0	0	0	1/18	1/36	1/72	\dots	8/9
4.	CC	0	0	0	0	1/34	1/68	\dots	16/17
5.	CD	0	0	0	0	1	0	\dots	0

Example: the disrupted unforgiving reply rule

Define

$g_t =_{Def}$ Play unforgiving before t , defect unconditionally at t and later.

Player 1 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
0.		$1/4$	$1/8$	$1/16$	$1/32$	$1/64$	$1/128$	\dots	$1/2$
1.	CC	0	$1/6$	$1/12$	$1/24$	$1/48$	$1/96$	\dots	$2/3$
2.	CC	0	0	$1/10$	$1/20$	$1/40$	$1/80$	\dots	$4/5$
3.	CC	0	0	0	$1/18$	$1/36$	$1/72$	\dots	$8/9$
4.	CC	0	0	0	0	$1/34$	$1/68$	\dots	$16/17$
5.	CD	0	0	0	0	1	0	\dots	0
6.	DD	0	0	0	0	1	0	\dots	0

Example: the disrupted unforgiving reply rule

Define

$g_t =_{Def}$ Play unforgiving before t , defect unconditionally at t and later.

Player 1 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
0.		1/4	1/8	1/16	1/32	1/64	1/128	\dots	1/2
1.	CC	0	1/6	1/12	1/24	1/48	1/96	\dots	2/3
2.	CC	0	0	1/10	1/20	1/40	1/80	\dots	4/5
3.	CC	0	0	0	1/18	1/36	1/72	\dots	8/9
4.	CC	0	0	0	0	1/34	1/68	\dots	16/17
5.	CD	0	0	0	0	1	0	\dots	0
6.	DD	0	0	0	0	1	0	\dots	0
\vdots		\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\dots	\vdots

Disrupted unforgiving strategy: opponent's perspective

Same game (and same realisation of play) but now how beliefs of Player 2 evolve.

Disrupted unforgiving strategy: opponent's perspective

Same game (and same realisation of play) but now how beliefs of Player 2 evolve.

Player 2 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Disrupted unforgiving strategy: opponent's perspective

Same game (and same realisation of play) but now how beliefs of Player 2 evolve.

Player 2 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round	g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
-------	-------	-------	-------	-------	-------	-------	---------	------------

Disrupted unforgiving strategy: opponent's perspective

Same game (and same realisation of play) but now how beliefs of Player 2 evolve.

Player 2 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round	g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
	$1/4$	$1/8$	$1/16$	$1/32$	$1/64$	$1/128$	\dots	$1/2$

Disrupted unforgiving strategy: opponent's perspective

Same game (and same realisation of play) but now how beliefs of Player 2 evolve.

Player 2 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
		$1/4$	$1/8$	$1/16$	$1/32$	$1/64$	$1/128$	\dots	$1/2$
1.	CC	0	$1/6$	$1/12$	$1/24$	$1/48$	$1/96$	\dots	$2/3$

Disrupted unforgiving strategy: opponent's perspective

Same game (and same realisation of play) but now how beliefs of Player 2 evolve.

Player 2 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
		$1/4$	$1/8$	$1/16$	$1/32$	$1/64$	$1/128$	\dots	$1/2$
1.	CC	0	$1/6$	$1/12$	$1/24$	$1/48$	$1/96$	\dots	$2/3$
2.	CC	0	0	$1/10$	$1/20$	$1/40$	$1/80$	\dots	$4/5$

Disrupted unforgiving strategy: opponent's perspective

Same game (and same realisation of play) but now how beliefs of Player 2 evolve.

Player 2 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
		$1/4$	$1/8$	$1/16$	$1/32$	$1/64$	$1/128$	\dots	$1/2$
1.	CC	0	$1/6$	$1/12$	$1/24$	$1/48$	$1/96$	\dots	$2/3$
2.	CC	0	0	$1/10$	$1/20$	$1/40$	$1/80$	\dots	$4/5$
3.	CC	0	0	0	$1/18$	$1/36$	$1/72$	\dots	$8/9$

Disrupted unforgiving strategy: opponent's perspective

Same game (and same realisation of play) but now how beliefs of Player 2 evolve.

Player 2 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
		$1/4$	$1/8$	$1/16$	$1/32$	$1/64$	$1/128$	\dots	$1/2$
1.	CC	0	$1/6$	$1/12$	$1/24$	$1/48$	$1/96$	\dots	$2/3$
2.	CC	0	0	$1/10$	$1/20$	$1/40$	$1/80$	\dots	$4/5$
3.	CC	0	0	0	$1/18$	$1/36$	$1/72$	\dots	$8/9$
4.	CC	0	0	0	0	$1/34$	$1/68$	\dots	$16/17$

Disrupted unforgiving strategy: opponent's perspective

Same game (and same realisation of play) but now how beliefs of Player 2 evolve.

Player 2 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
		1/4	1/8	1/16	1/32	1/64	1/128	\dots	1/2
1.	CC	0	1/6	1/12	1/24	1/48	1/96	\dots	2/3
2.	CC	0	0	1/10	1/20	1/40	1/80	\dots	4/5
3.	CC	0	0	0	1/18	1/36	1/72	\dots	8/9
4.	CC	0	0	0	0	1/34	1/68	\dots	16/17
5.	CD	0	0	0	0	0	1/66	\dots	32/33

Disrupted unforgiving strategy: opponent's perspective

Same game (and same realisation of play) but now how beliefs of Player 2 evolve.

Player 2 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
		1/4	1/8	1/16	1/32	1/64	1/128	\dots	1/2
1.	CC	0	1/6	1/12	1/24	1/48	1/96	\dots	2/3
2.	CC	0	0	1/10	1/20	1/40	1/80	\dots	4/5
3.	CC	0	0	0	1/18	1/36	1/72	\dots	8/9
4.	CC	0	0	0	0	1/34	1/68	\dots	16/17
5.	CD	0	0	0	0	0	1/66	\dots	32/33
6.	DD	0	0	0	0	0	1/66	\dots	32/33

Disrupted unforgiving strategy: opponent's perspective

Same game (and same realisation of play) but now how beliefs of Player 2 evolve.

Player 2 gives positive prior probability to strategies $\{g_t\}_{t \geq 1} \cup \{g_\infty\}$.

Round		g_1	g_2	g_3	g_4	g_5	g_6	\dots	g_∞
		$1/4$	$1/8$	$1/16$	$1/32$	$1/64$	$1/128$	\dots	$1/2$
1.	CC	0	$1/6$	$1/12$	$1/24$	$1/48$	$1/96$	\dots	$2/3$
2.	CC	0	0	$1/10$	$1/20$	$1/40$	$1/80$	\dots	$4/5$
3.	CC	0	0	0	$1/18$	$1/36$	$1/72$	\dots	$8/9$
4.	CC	0	0	0	0	$1/34$	$1/68$	\dots	$16/17$
5.	CD	0	0	0	0	0	$1/66$	\dots	$32/33$
6.	DD	0	0	0	0	0	$1/66$	\dots	$32/33$
\vdots		\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\dots	\vdots

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

$$S = \{ L^* : \text{stay left forever} , \\ R^* : \text{stay right forever} , \\ F^* : \text{play mixed } 0.5 \text{ forever} \}.$$

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

$$S = \{ L^* : \text{stay left forever} , \\ R^* : \text{stay right forever} , \\ F^* : \text{play mixed } 0.5 \text{ forever} \}.$$

Suppose

	1's prior	2's prior
L^*	0.6	0.2
R^*	0.3	0.5
F^*	0.1	0.3.

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

$$S = \{ L^* : \text{stay left forever} , \\ R^* : \text{stay right forever} , \\ F^* : \text{play mixed } 0.5 \text{ forever} \}.$$

Suppose

	1's prior	2's prior
L^*	0.6	0.2
R^*	0.3	0.5
F^*	0.1	0.3.

After

$$h = [(L, R), (R, L)]$$

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

Player 1 reasons:

$S = \{ L^* : \text{stay left forever} ,$
 $R^* : \text{stay right forever} ,$
 $F^* : \text{play mixed } 0.5 \text{ forever} \}.$

Suppose

	1's prior	2's prior
L^*	0.6	0.2
R^*	0.3	0.5
F^*	0.1	0.3.

After

$$h = [(L, R), (R, L)]$$

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

Player 1 reasons:

$$\Pr\{R^*|h\}$$

$S = \{ L^* : \text{stay left forever} ,$
 $R^* : \text{stay right forever} ,$
 $F^* : \text{play mixed } 0.5 \text{ forever} \}.$

Suppose

	1's prior	2's prior
L^*	0.6	0.2
R^*	0.3	0.5
F^*	0.1	0.3.

After

$$h = [(L, R), (R, L)]$$

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

$S = \{ L^* : \text{stay left forever} ,$
 $R^* : \text{stay right forever} ,$
 $F^* : \text{play mixed } 0.5 \text{ forever} \}.$

Suppose

	1's prior	2's prior
L^*	0.6	0.2
R^*	0.3	0.5
F^*	0.1	0.3.

After

$$h = [(L, R), (R, L)]$$

Player 1 reasons:

$$\Pr\{R^*|h\} = \frac{\Pr\{h|R^*\}\Pr\{R^*\}}{\sum_{j=1}^3 \Pr\{h|s_j\}\Pr\{s_j\}}$$

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

$S = \{ L^* : \text{stay left forever} ,$
 $R^* : \text{stay right forever} ,$
 $F^* : \text{play mixed } 0.5 \text{ forever} \}.$

Suppose

	1's prior	2's prior
L^*	0.6	0.2
R^*	0.3	0.5
F^*	0.1	0.3.

After

$$h = [(L, R), (R, L)]$$

Player 1 reasons:

$$\begin{aligned}\Pr\{R^*|h\} &= \frac{\Pr\{h|R^*\}\Pr\{R^*\}}{\sum_{j=1}^3 \Pr\{h|s_j\}\Pr\{s_j\}} \\ &= \frac{0 \cdot 0.3}{0 \cdot 0.6 + 0 \cdot 0.3 + (\frac{1}{2})^2 \cdot 0.1}\end{aligned}$$

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

$S = \{ L^* : \text{stay left forever} ,$
 $R^* : \text{stay right forever} ,$
 $F^* : \text{play mixed } 0.5 \text{ forever} \}.$

Suppose

	1's prior	2's prior
L^*	0.6	0.2
R^*	0.3	0.5
F^*	0.1	0.3.

After

$$h = [(L, R), (R, L)]$$

Player 1 reasons:

$$\begin{aligned}\Pr\{R^*|h\} &= \frac{\Pr\{h|R^*\}\Pr\{R^*\}}{\sum_{j=1}^3 \Pr\{h|s_j\}\Pr\{s_j\}} \\ &= \frac{0 \cdot 0.3}{0 \cdot 0.6 + 0 \cdot 0.3 + (\frac{1}{2})^2 \cdot 0.1} \\ &= 0.\end{aligned}$$

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

$S = \{ L^* : \text{stay left forever} ,$
 $R^* : \text{stay right forever} ,$
 $F^* : \text{play mixed } 0.5 \text{ forever} \}.$

Suppose

	1's prior	2's prior
L^*	0.6	0.2
R^*	0.3	0.5
F^*	0.1	0.3.

After

$$h = [(L, R), (R, L)]$$

Player 1 reasons:

$$\begin{aligned}\Pr\{R^*|h\} &= \frac{\Pr\{h|R^*\}\Pr\{R^*\}}{\sum_{j=1}^3 \Pr\{h|s_j\}\Pr\{s_j\}} \\ &= \frac{0 \cdot 0.3}{0 \cdot 0.6 + 0 \cdot 0.3 + (\frac{1}{2})^2 \cdot 0.1} \\ &= 0.\end{aligned}$$

$$[\Pr\{(L, \cdot), (R, \cdot)|F^*\} = (\frac{1}{2})^2] \text{ and}$$

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

$S = \{ L^* : \text{stay left forever} ,$
 $R^* : \text{stay right forever} ,$
 $F^* : \text{play mixed } 0.5 \text{ forever} \}.$

Suppose

	1's prior	2's prior
L^*	0.6	0.2
R^*	0.3	0.5
F^*	0.1	0.3.

After

$$h = [(L, R), (R, L)]$$

Player 1 reasons:

$$\begin{aligned}\Pr\{R^*|h\} &= \frac{\Pr\{h|R^*\}\Pr\{R^*\}}{\sum_{j=1}^3 \Pr\{h|s_j\}\Pr\{s_j\}} \\ &= \frac{0 \cdot 0.3}{0 \cdot 0.6 + 0 \cdot 0.3 + (\frac{1}{2})^2 \cdot 0.1} \\ &= 0.\end{aligned}$$

$$[\Pr\{(L, \cdot), (R, \cdot)|F^*\} = (\frac{1}{2})^2] \text{ and}$$

$$\Pr\{F^*|h\}$$

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

$S = \{ L^* : \text{stay left forever} ,$
 $R^* : \text{stay right forever} ,$
 $F^* : \text{play mixed } 0.5 \text{ forever} \}.$

Suppose

	1's prior	2's prior
L^*	0.6	0.2
R^*	0.3	0.5
F^*	0.1	0.3.

After

$$h = [(L, R), (R, L)]$$

Player 1 reasons:

$$\begin{aligned}\Pr\{R^*|h\} &= \frac{\Pr\{h|R^*\}\Pr\{R^*\}}{\sum_{j=1}^3 \Pr\{h|s_j\}\Pr\{s_j\}} \\ &= \frac{0 \cdot 0.3}{0 \cdot 0.6 + 0 \cdot 0.3 + (\frac{1}{2})^2 \cdot 0.1} \\ &= 0.\end{aligned}$$

$$[\Pr\{(L, \cdot), (R, \cdot)|F^*\} = (\frac{1}{2})^2] \text{ and}$$

$$\Pr\{F^*|h\} = \frac{\Pr\{h|F^*\}\Pr\{F^*\}}{\sum_{j=1}^3 \Pr\{h|s_j\}\Pr\{s_j\}}$$

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

$S = \{ L^* : \text{stay left forever}, \\ R^* : \text{stay right forever}, \\ F^* : \text{play mixed } 0.5 \text{ forever} \}.$

Suppose

	1's prior	2's prior
L^*	0.6	0.2
R^*	0.3	0.5
F^*	0.1	0.3.

After

$$h = [(L, R), (R, L)]$$

Player 1 reasons:

$$\begin{aligned} \Pr\{R^*|h\} &= \frac{\Pr\{h|R^*\}\Pr\{R^*\}}{\sum_{j=1}^3 \Pr\{h|s_j\}\Pr\{s_j\}} \\ &= \frac{0 \cdot 0.3}{0 \cdot 0.6 + 0 \cdot 0.3 + (\frac{1}{2})^2 \cdot 0.1} \\ &= 0. \end{aligned}$$

$$[\Pr\{(L, \cdot), (R, \cdot)|F^*\} = (\frac{1}{2})^2] \text{ and}$$

$$\begin{aligned} \Pr\{F^*|h\} &= \frac{\Pr\{h|F^*\}\Pr\{F^*\}}{\sum_{j=1}^3 \Pr\{h|s_j\}\Pr\{s_j\}} \\ &= \frac{(\frac{1}{2})^2 \cdot 0.1}{0 \cdot 0.6 + 0 \cdot 0.3 + (\frac{1}{2})^2 \cdot 0.1} \end{aligned}$$

Example: Coordination game

Suppose Player 1 and 2 play the coordination game, and deem the following strategies possible:

$S = \{ L^* : \text{stay left forever}, \\ R^* : \text{stay right forever}, \\ F^* : \text{play mixed } 0.5 \text{ forever} \}.$

Suppose

	1's prior	2's prior
L^*	0.6	0.2
R^*	0.3	0.5
F^*	0.1	0.3.

After

$$h = [(L, R), (R, L)]$$

Player 1 reasons:

$$\begin{aligned}\Pr\{R^*|h\} &= \frac{\Pr\{h|R^*\}\Pr\{R^*\}}{\sum_{j=1}^3 \Pr\{h|s_j\}\Pr\{s_j\}} \\ &= \frac{0 \cdot 0.3}{0 \cdot 0.6 + 0 \cdot 0.3 + (\frac{1}{2})^2 \cdot 0.1} \\ &= 0.\end{aligned}$$

$$[\Pr\{(L, \cdot), (R, \cdot)|F^*\} = (\frac{1}{2})^2] \text{ and}$$

$$\begin{aligned}\Pr\{F^*|h\} &= \frac{\Pr\{h|F^*\}\Pr\{F^*\}}{\sum_{j=1}^3 \Pr\{h|s_j\}\Pr\{s_j\}} \\ &= \frac{(\frac{1}{2})^2 \cdot 0.1}{0 \cdot 0.6 + 0 \cdot 0.3 + (\frac{1}{2})^2 \cdot 0.1} \\ &= 1.\end{aligned}$$

Extend the support of possible strategies

Problem. If Player 2 indeed plays F^* , then Player 1 has solved the problem.

Extend the support of possible strategies

Problem. If Player 2 indeed plays F^* , then Player 1 has solved the problem.

But Player 2 need not play F^* :

Extend the support of possible strategies

Problem. If Player 2 indeed plays F^* , then Player 1 has solved the problem.

But Player 2 need not play F^* :

- Like Player 1, Player 2 also has deduced that its opponent plays F^* .

Extend the support of possible strategies

Problem. If Player 2 indeed plays F^* , then Player 1 has solved the problem.

But Player 2 need not play F^* :

- Like Player 1, Player 2 also has deduced that its opponent plays F^* .
- Any action is a best reply to F^* . (Verify, if necessary.)

Extend the support of possible strategies

Problem. If Player 2 indeed plays F^* , then Player 1 has solved the problem.

But Player 2 need not play F^* :

- Like Player 1, Player 2 also has deduced that its opponent plays F^* .
- Any action is a best reply to F^* . (Verify, if necessary.)
- So Player 2 could in fact play any strategy after it observed h .

Extend the support of possible strategies

Problem. If Player 2 indeed plays F^* , then Player 1 has solved the problem.

But Player 2 need not play F^* :

- Like Player 1, Player 2 also has deduced that its opponent plays F^* .
- Any action is a best reply to F^* . (Verify, if necessary.)
- So Player 2 could in fact play any strategy after it observed h .

Solution. Extend S by letting $S =_{Def} \{s_p \mid 0 \leq p \leq 1\}$. This gives us all stationary mixed strategies.

Extend the support of possible strategies

Problem. If Player 2 indeed plays F^* , then Player 1 has solved the problem.

But Player 2 need not play F^* :

- Like Player 1, Player 2 also has deduced that its opponent plays F^* .
- Any action is a best reply to F^* . (Verify, if necessary.)
- So Player 2 could in fact play any strategy after it observed h .

Solution. Extend S by letting $S =_{Def} \{s_p \mid 0 \leq p \leq 1\}$. This gives us all stationary mixed strategies. In particular, $F^* = s_{0.5}$.

Extend the support of possible strategies

Problem. If Player 2 indeed plays F^* , then Player 1 has solved the problem.

But Player 2 need not play F^* :

- Like Player 1, Player 2 also has deduced that its opponent plays F^* .
- Any action is a best reply to F^* . (Verify, if necessary.)
- So Player 2 could in fact play any strategy after it observed h .

Solution. Extend S by letting $S =_{Def} \{s_p \mid 0 \leq p \leq 1\}$. This gives us all stationary mixed strategies. In particular, $F^* = s_{0.5}$.

Bayes' theorem for continuous random variables gives

$$f(s = s_p | h) = \frac{\Pr\{h | s = s_p\} f(s_p)}{\int_{p=0}^{p=1} \Pr\{h | s = s_p\} f(s_p) dp}$$

Extend the support of possible strategies

Problem. If Player 2 indeed plays F^* , then Player 1 has solved the problem.

But Player 2 need not play F^* :

- Like Player 1, Player 2 also has deduced that its opponent plays F^* .
- Any action is a best reply to F^* . (Verify, if necessary.)
- So Player 2 could in fact play any strategy after it observed h .

Solution. Extend S by letting $S =_{Def} \{s_p \mid 0 \leq p \leq 1\}$. This gives us all stationary mixed strategies. In particular, $F^* = s_{0.5}$.

Bayes' theorem for continuous random variables gives

$$f(s = s_p | h) = \frac{\Pr\{h | s = s_p\} f(s_p)}{\int_{p=0}^{p=1} \Pr\{h | s = s_p\} f(s_p) dp}$$

Now, $\text{BR}(s_p | h) \cap S \neq \emptyset$, so that from round to round play vs. prediction is a closed system.

Then give all possible play paths positive priors?

Then give all possible play paths positive priors?

- Enter **measure theory**, which is a mathematical discipline in and of itself ...

Then give all possible play paths positive priors?

- Enter **measure theory**, which is a mathematical discipline in and of itself ...
- With a diagonalisation argument it can be shown that **the set of all realisations** (paths of play), Ω , is uncountable.

Then give all possible play paths positive priors?

- Enter **measure theory**, which is a mathematical discipline in and of itself ...
- With a diagonalisation argument it can be shown that **the set of all realisations** (paths of play), Ω , is uncountable.
- **Question:** is it possible to have a probability measure

$$\mu : \Omega \rightarrow [0, 1]$$

such that $\mu\{E\} > 0$ for all non-empty $E \subseteq \Omega$?

Then give all possible play paths positive priors?

- Enter **measure theory**, which is a mathematical discipline in and of itself ...
- With a diagonalisation argument it can be shown that **the set of all realisations** (paths of play), Ω , is uncountable.
- **Question:** is it possible to have a probability measure

$$\mu : \Omega \rightarrow [0, 1]$$

such that $\mu\{E\} > 0$ for all non-empty $E \subseteq \Omega$?

Know that a probability

measure must be closed under complement and countable union.

Then give all possible play paths positive priors?

- Enter **measure theory**, which is a mathematical discipline in and of itself ...
- With a diagonalisation argument it can be shown that **the set of all realisations** (paths of play), Ω , is uncountable.
- **Question:** is it possible to have a probability measure

$$\mu : \Omega \rightarrow [0, 1]$$

such that $\mu\{E\} > 0$ for all non-empty $E \subseteq \Omega$?

Know that a probability

measure must be closed under complement and countable union. (This is known as a **σ -algebra**).

Then give all possible play paths positive priors?

- Enter **measure theory**, which is a mathematical discipline in and of itself ...
- With a diagonalisation argument it can be shown that **the set of all realisations** (paths of play), Ω , is uncountable.
- **Question:** is it possible to have a probability measure

$$\mu : \Omega \rightarrow [0, 1]$$

such that $\mu\{E\} > 0$ for all non-empty $E \subseteq \Omega$?

Know that a probability

measure must be closed under complement and countable union. (This is known as a **σ -algebra**).

- **Answer: no.** The set 2^Ω is “too large” \Rightarrow problems and paradoxes. (Banach-Tarski.)

Then give all possible play paths positive priors?

- Enter **measure theory**, which is a mathematical discipline in and of itself ...
- With a diagonalisation argument it can be shown that **the set of all realisations** (paths of play), Ω , is uncountable.
- **Question:** is it possible to have a probability measure

$$\mu : \Omega \rightarrow [0, 1]$$

such that $\mu\{E\} > 0$ for all non-empty $E \subseteq \Omega$?

Know that a probability

measure must be closed under complement and countable union. (This is known as a **σ -algebra**).

- **Answer: no.** The set 2^Ω is “too large” \Rightarrow problems and paradoxes. (Banach-Tarski.)



Then give all possible play paths positive priors?

- Enter **measure theory**, which is a mathematical discipline in and of itself ...
- With a diagonalisation argument it can be shown that **the set of all realisations** (paths of play), Ω , is uncountable.
- **Question:** is it possible to have a probability measure

$$\mu : \Omega \rightarrow [0, 1]$$

such that $\mu\{E\} > 0$ for all non-empty $E \subseteq \Omega$?

Know that a probability

measure must be closed under complement and countable union. (This is known as a **σ -algebra**).

- **Answer: no.** The set 2^Ω is “too large” \Rightarrow problems and paradoxes. (Banach-Tarski.)



- **Solution:** extend μ of H to μ on a **σ -algebra** of Ω .

Part III: Formalism

Probabilities on action profiles

Probabilities on action profiles

- $\Delta[X]$: possible probability distributions over X .

Probabilities on action profiles

- $\Delta[X]$: possible probability distributions over X .
- $\Delta[X_i]$: possible mixed strategies for Player i

Probabilities on action profiles

- $\Delta[X]$: possible probability distributions over X .
- $\Delta[X_i]$: possible mixed strategies for Player i , often abbreviated to Δ_i .

Probabilities on action profiles

- $\Delta[X]$: possible probability distributions over X .
- $\Delta[X_i]$: possible mixed strategies for Player i , often abbreviated to Δ_i .
- The set of possible probability distributions over all counter-profiles

$$\Delta[X_{-i}], \quad (1)$$

Probabilities on action profiles

- $\Delta[X]$: possible probability distributions over X .
- $\Delta[X_i]$: possible mixed strategies for Player i , often abbreviated to Δ_i .
- The set of possible probability distributions over all counter-profiles

$$\Delta[X_{-i}], \quad (1)$$

(where X_{-i} is shorthand for $\prod_{j \neq i} X_j$)

Probabilities on action profiles

- $\Delta[X]$: possible probability distributions over X .
- $\Delta[X_i]$: possible mixed strategies for Player i , often abbreviated to Δ_i .
- The set of possible probability distributions over all counter-profiles

$$\Delta[X_{-i}], \quad (1)$$

(where X_{-i} is shorthand for $\prod_{j \neq i} X_j$)
allows that actions may be
dependent:

$$\Pr\{x_1 = a_1, \dots, x_n = a_n\} \neq \Pr\{x_1 = a_1\} \times \dots \times \Pr\{x_n = a_n\}.$$

Probabilities on action profiles

- $\Delta[X]$: possible probability distributions over X .
- $\Delta[X_i]$: possible mixed strategies for Player i , often abbreviated to Δ_i .
- The set of possible probability distributions over all counter-profiles

$$\Delta[X_{-i}], \quad (1)$$

(where X_{-i} is shorthand for $\prod_{j \neq i} X_j$)
allows that actions may be
dependent:

$$\Pr\{x_1 = a_1, \dots, x_n = a_n\} \neq \Pr\{x_1 = a_1\} \times \dots \times \Pr\{x_n = a_n\}.$$

- The product of possible mixed strategies

$$\prod_{j \neq i} \Delta_j. \quad (2)$$

contains much less
information, actions are
necessarily independent:

$$\Pr\{x_1 = a_1, \dots, x_n = a_n\} = \Pr\{x_1 = a_1\} \times \dots \times \Pr\{x_n = a_n\}.$$

Probabilities on action profiles

- $\Delta[X]$: possible probability distributions over X .
- $\Delta[X_i]$: possible mixed strategies for Player i , often abbreviated to Δ_i .
- The set of possible probability distributions over all counter-profiles

$$\Delta[X_{-i}], \quad (1)$$

(where X_{-i} is shorthand for $\prod_{j \neq i} X_j$) allows that actions may be **dependent**:

$$\Pr\{x_1 = a_1, \dots, x_n = a_n\} \neq \Pr\{x_1 = a_1\} \times \dots \times \Pr\{x_n = a_n\}.$$

- The product of possible mixed strategies

$$\prod_{j \neq i} \Delta_j. \quad (2)$$

contains much less information, actions are necessarily independent:

$$\Pr\{x_1 = a_1, \dots, x_n = a_n\} = \Pr\{x_1 = a_1\} \times \dots \times \Pr\{x_n = a_n\}.$$

- The abbreviation Δ_{-i} may now mean (1) or (2).

Probabilities on action profiles

- $\Delta[X]$: possible probability distributions over X .
- $\Delta[X_i]$: possible mixed strategies for Player i , often abbreviated to Δ_i .
- The set of possible probability distributions over all counter-profiles

$$\Delta[X_{-i}], \quad (1)$$

(where X_{-i} is shorthand for $\prod_{j \neq i} X_j$)
allows that actions may be
dependent:

$$\Pr\{x_1 = a_1, \dots, x_n = a_n\} \neq \Pr\{x_1 = a_1\} \times \dots \times \Pr\{x_n = a_n\}.$$

- The product of possible mixed strategies

$$\prod_{j \neq i} \Delta_j. \quad (2)$$

contains much less
information, actions are
necessarily independent:

$$\Pr\{x_1 = a_1, \dots, x_n = a_n\} = \Pr\{x_1 = a_1\} \times \dots \times \Pr\{x_n = a_n\}.$$

- The abbreviation Δ_{-i} may now mean (1) or (2).

Often, (2) is meant.

Framework for predictive learning (Jordan, 1993)

Framework for predictive learning (Jordan, 1993)

- A forecasting rule for player i is a function that maps every history to a probability distribution over counter-action profiles:

$$f_i : H \rightarrow \Delta_{-i}.$$

Framework for predictive learning (Jordan, 1993)

- A forecasting rule for player i is a function that maps every history to a probability distribution over counter-action profiles:

$$f_i : H \rightarrow \Delta_{-i}.$$

- Abbreviate $\Delta[X_i]$ to Δ_i .

Framework for predictive learning (Jordan, 1993)

- A **forecasting rule** for player i is a function that maps every history to a probability distribution over counter-action profiles:

$$f_i : H \rightarrow \Delta_{-i}.$$

- Abbreviate $\Delta[X_i]$ to Δ_i .

A **reply rule** for player i is a function that maps a history to a probability distribution over i 's own actions in the next round:

$$g_i : H \rightarrow \Delta_i.$$

Framework for predictive learning (Jordan, 1993)

- A **forecasting rule** for player i is a function that maps every history to a probability distribution over counter-action profiles:
- A **predictive learning rule** for player i is the combination of a forecasting rule and a reply rule. This is typically written as (f_i, g_i) .

$$f_i : H \rightarrow \Delta_{-i}.$$

- Abbreviate $\Delta[X_i]$ to Δ_i .

A **reply rule** for player i is a function that maps a history to a probability distribution over i 's own actions in the next round:

$$g_i : H \rightarrow \Delta_i.$$

Framework for predictive learning (Jordan, 1993)

- A **forecasting rule** for player i is a function that maps every history to a probability distribution over counter-action profiles:
- A **predictive learning rule** for player i is the combination of a forecasting rule and a reply rule. This is typically written as (f_i, g_i) .

$$f_i : H \rightarrow \Delta_{-i}.$$

- Abbreviate $\Delta[X_i]$ to Δ_i .

A **reply rule** for player i is a function that maps a history to a probability distribution over i 's own actions in the next round:

$$g_i : H \rightarrow \Delta_i.$$

- The idea is that g_i uses f_{i-1} .

Framework for predictive learning (Jordan, 1993)

- A **forecasting rule** for player i is a function that maps every history to a probability distribution over counter-action profiles:

$$f_i : H \rightarrow \Delta_{-i}.$$

- Abbreviate $\Delta[X_i]$ to Δ_i .

A **reply rule** for player i is a function that maps a history to a probability distribution over i 's own actions in the next round:

$$g_i : H \rightarrow \Delta_i.$$

- A **predictive learning rule** for player i is the combination of a forecasting rule and a reply rule. This is typically written as (f_i, g_i) .

- The idea is that g_i uses f_{i-1} .
- Forecasting and reply functions are **deterministic**.

Framework for predictive learning (Jordan, 1993)

- A **forecasting rule** for player i is a function that maps every history to a probability distribution over counter-action profiles:

$$f_i : H \rightarrow \Delta_{-i}.$$

- Abbreviate $\Delta[X_i]$ to Δ_i .

A **reply rule** for player i is a function that maps a history to a probability distribution over i 's own actions in the next round:

$$g_i : H \rightarrow \Delta_i.$$

- A **predictive learning rule** for player i is the combination of a forecasting rule and a reply rule. This is typically written as (f_i, g_i) .

- The idea is that g_i uses f_{i-1} .
- Forecasting and reply functions are **deterministic**.
- Reinforcement and regret do not fit. They are not involved with prediction.

Framework for predictive learning (Jordan, 1993)

- A **forecasting rule** for player i is a function that maps every history to a probability distribution over counter-action profiles:

$$f_i : H \rightarrow \Delta_{-i}.$$

- Abbreviate $\Delta[X_i]$ to Δ_i .

A **reply rule** for player i is a function that maps a history to a probability distribution over i 's own actions in the next round:

$$g_i : H \rightarrow \Delta_i.$$

- A **predictive learning rule** for player i is the combination of a forecasting rule and a reply rule. This is typically written as (f_i, g_i) .

- The idea is that g_i uses f_{i-1} .
- Forecasting and reply functions are **deterministic**.
- Reinforcement and regret do not fit. They are not involved with prediction.
- Fictitious play and Bayesian play do fit.

Beliefs about reply rules of other players: ν_i

Beliefs about reply rules of other players: ν_i

- The beliefs of Player i about the reply rules of other players is represented by a probability distribution ν_i over the reply rules of other players. It is therefore an element of

$$\Delta[\Pi_{j \neq i} \Delta_j^H]. \quad (3)$$

Beliefs about reply rules of other players: ν_i

- The beliefs of Player i about the reply rules of other players is represented by a probability distribution ν_i over the reply rules of other players. It is therefore an element of

$$\Delta[\Pi_{j \neq i} \Delta_j^H]. \quad (3)$$

- If Player i assumes that its opponents operate independently, then (3) reduces to

$$\Pi_{j \neq i} \Delta[\Delta_j^H]. \quad (4)$$

Beliefs about reply rules of other players: ν_i

- The **beliefs** of Player i about the reply rules of other players is represented by a probability distribution ν_i over the reply rules of other players. It is therefore an element of

$$\Delta[\Pi_{j \neq i} \Delta_j^H]. \quad (3)$$

- If Player i assumes that its opponents operate independently, then (3) reduces to

$$\Pi_{j \neq i} \Delta[\Delta_j^H]. \quad (4)$$

Theorem (Kuhn, 1953). Every distribution over mixed strategies of opponents can be represented by a single set of mixed strategies of opponents

Beliefs about reply rules of other players: ν_i

- The beliefs of Player i about the reply rules of other players is represented by a probability distribution ν_i over the reply rules of other players. It is therefore an element of

$$\Delta[\Pi_{j \neq i} \Delta_j^H]. \quad (3)$$

- If Player i assumes that its opponents operate independently, then (3) reduces to

$$\Pi_{j \neq i} \Delta[\Delta_j^H]. \quad (4)$$

Theorem (Kuhn, 1953). Every distribution over mixed strategies of opponents can be represented by a single set of mixed strategies of opponents

Proof: by conditional expectation. (Cf. Kalai and Lehrer, 1993, p. 1031.)

Beliefs about reply rules of other players: ν_i

- The beliefs of Player i about the reply rules of other players is represented by a probability distribution ν_i over the reply rules of other players. It is therefore an element of

$$\Delta[\Pi_{j \neq i} \Delta_j^H]. \quad (3)$$

- If Player i assumes that its opponents operate independently, then (3) reduces to

$$\Pi_{j \neq i} \Delta[\Delta_j^H]. \quad (4)$$

- **Theorem (Kuhn, 1953).** Every distribution over mixed strategies of opponents can be represented by a single set of mixed strategies of opponents

Proof: by conditional expectation. (Cf. Kalai and Lehrer, 1993, p. 1031.)

Therefore, (4) further reduces to

$$\Pi_{j \neq i} \Delta_j^H.$$

Beliefs about reply rules of other players: ν_i

- The **beliefs** of Player i about the reply rules of other players is represented by a probability distribution ν_i over the reply rules of other players. It is therefore an element of

$$\Delta[\Pi_{j \neq i} \Delta_j^H]. \quad (3)$$

- If Player i assumes that its opponents operate independently, then (3) reduces to

$$\Pi_{j \neq i} \Delta[\Delta_j^H]. \quad (4)$$

- **Theorem (Kuhn, 1953).** Every distribution over mixed strategies of opponents can be represented by a single set of mixed strategies of opponents

Proof: by conditional expectation. (Cf. Kalai and Lehrer, 1993, p. 1031.)

Therefore, (4) further reduces to

$$\Pi_{j \neq i} \Delta_j^H.$$

Often, this is abbreviated to Δ_{-i}^H .

Idea of Kuhn's proof

distribution		λ_1	λ_2	λ_3	\dots	λ_r	\dots	
strategies	\mathbf{G}	\mathbf{g}_1^j	\mathbf{g}_2^j	\mathbf{g}_3^j	\dots	\mathbf{g}_r^j	\dots	\mathbf{g}^j
histories	h_1	q_1^1	q_2^1	q_3^1	\dots	q_r^1	\dots	$\lambda_1 q_1^1 + \dots + \lambda_r q_r^1 + \dots$
	h_2	q_1^2	q_2^2	\dots	\dots	q_r^2	\dots	$\lambda_1 q_1^2 + \dots + \lambda_r q_r^2 + \dots$
	h_3	q_1^3	\dots	\dots	\dots	q_r^3	\dots	$\lambda_1 q_1^3 + \dots + \lambda_r q_r^3 + \dots$
	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots	\vdots

Idea of Kuhn's proof

distribution		λ_1	λ_2	λ_3	\dots	λ_r	\dots	
strategies	\mathbf{G}	\mathbf{g}_1^j	\mathbf{g}_2^j	\mathbf{g}_3^j	\dots	\mathbf{g}_r^j	\dots	\mathbf{g}^j
histories	h_1	q_1^1	q_2^1	q_3^1	\dots	q_r^1	\dots	$\lambda_1 q_1^1 + \dots + \lambda_r q_r^1 + \dots$
	h_2	q_1^2	q_2^2	\dots	\dots	q_r^2	\dots	$\lambda_1 q_1^2 + \dots + \lambda_r q_r^2 + \dots$
	h_3	q_1^3	\dots	\dots	\dots	q_r^3	\dots	$\lambda_1 q_1^3 + \dots + \lambda_r q_r^3 + \dots$
	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots	\vdots

Remarks:

Idea of Kuhn's proof

distribution		λ_1	λ_2	λ_3	\dots	λ_r	\dots	
strategies	\mathbf{G}	\mathbf{g}_1^j	\mathbf{g}_2^j	\mathbf{g}_3^j	\dots	\mathbf{g}_r^j	\dots	\mathbf{g}^j
histories	h_1	q_1^1	q_2^1	q_3^1	\dots	q_r^1	\dots	$\lambda_1 q_1^1 + \dots + \lambda_r q_r^1 + \dots$
	h_2	q_1^2	q_2^2	\dots	\dots	q_r^2	\dots	$\lambda_1 q_1^2 + \dots + \lambda_r q_r^2 + \dots$
	h_3	q_1^3	\dots	\dots	\dots	q_r^3	\dots	$\lambda_1 q_1^3 + \dots + \lambda_r q_r^3 + \dots$
	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots	\vdots

Remarks:

1. This table suggests that S is countable, but S may be uncountable. (For example, if the prior is a β -distribution.)

Idea of Kuhn's proof

distribution		λ_1	λ_2	λ_3	\dots	λ_r	\dots	
strategies	\mathbf{G}	\mathbf{g}_1^j	\mathbf{g}_2^j	\mathbf{g}_3^j	\dots	\mathbf{g}_r^j	\dots	\mathbf{g}^j
histories	h_1	q_1^1	q_2^1	q_3^1	\dots	q_r^1	\dots	$\lambda_1 q_1^1 + \dots + \lambda_r q_r^1 + \dots$
	h_2	q_1^2	q_2^2	\dots	\dots	q_r^2	\dots	$\lambda_1 q_1^2 + \dots + \lambda_r q_r^2 + \dots$
	h_3	q_1^3	\dots	\dots	\dots	q_r^3	\dots	$\lambda_1 q_1^3 + \dots + \lambda_r q_r^3 + \dots$
	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots	\vdots

Remarks:

1. This table suggests that S is countable, but S may be uncountable. (For example, if the prior is a β -distribution.)
2. If S is uncountable, the sums on the right are integrals.

Idea of Kuhn's proof

distribution		λ_1	λ_2	λ_3	\dots	λ_r	\dots	
strategies	\mathbf{G}	\mathbf{g}_1^j	\mathbf{g}_2^j	\mathbf{g}_3^j	\dots	\mathbf{g}_r^j	\dots	\mathbf{g}^j
histories	h_1	q_1^1	q_2^1	q_3^1	\dots	q_r^1	\dots	$\lambda_1 q_1^1 + \dots + \lambda_r q_r^1 + \dots$
	h_2	q_1^2	q_2^2	\dots	\dots	q_r^2	\dots	$\lambda_1 q_1^2 + \dots + \lambda_r q_r^2 + \dots$
	h_3	q_1^3	\dots	\dots	\dots	q_r^3	\dots	$\lambda_1 q_1^3 + \dots + \lambda_r q_r^3 + \dots$
	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots	\vdots

Remarks:

1. This table suggests that S is countable, but S may be uncountable. (For example, if the prior is a β -distribution.)
2. If S is uncountable, the sums on the right are integrals.
3. It must be checked that every $\lambda_1 q_1^j + \dots + \lambda_r q_r^j + \dots$ is a legitimate mixed strategy.

Idea of Kuhn's proof

distribution		λ_1	λ_2	λ_3	\dots	λ_r	\dots	
strategies	\mathbf{G}	\mathbf{g}_1^j	\mathbf{g}_2^j	\mathbf{g}_3^j	\dots	\mathbf{g}_r^j	\dots	\mathbf{g}^j
histories	h_1	q_1^1	q_2^1	q_3^1	\dots	q_r^1	\dots	$\lambda_1 q_1^1 + \dots + \lambda_r q_r^1 + \dots$
	h_2	q_1^2	q_2^2	\dots	\dots	q_r^2	\dots	$\lambda_1 q_1^2 + \dots + \lambda_r q_r^2 + \dots$
	h_3	q_1^3	\dots	\dots	\dots	q_r^3	\dots	$\lambda_1 q_1^3 + \dots + \lambda_r q_r^3 + \dots$
	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots	\vdots

Remarks:

1. This table suggests that S is countable, but S may be uncountable. (For example, if the prior is a β -distribution.)
2. If S is uncountable, the sums on the right are integrals.
3. It must be checked that every $\lambda_1 q_1^j + \dots + \lambda_r q_r^j + \dots$ is a legitimate mixed strategy. The sum may then be abbreviated by, e.g., q^j .

Behavioural strategy and conditional forecast

Behavioural strategy and conditional forecast

- A behavioural strategy σ_i is a reply rule conditioned by beliefs.

Behavioural strategy and conditional forecast

- A behavioural strategy σ_i is a reply rule conditioned by beliefs. Put differently, it maps beliefs ν_i to a reply rule g_i :

$$\sigma_i : \Delta_{-i}^H \rightarrow \Delta_i^H : \nu_i \mapsto g_i.$$

Behavioural strategy and conditional forecast

- A behavioural strategy σ_i is a reply rule conditioned by beliefs. Put differently, it maps beliefs ν_i to a reply rule g_i :

$$\sigma_i : \Delta_{-i}^H \rightarrow \Delta_i^H : \nu_i \mapsto g_i.$$

So

$$\sigma_i \in (\Delta_i^H)^{\Delta_{-i}^H}.$$

Behavioural strategy and conditional forecast

- A behavioural strategy σ_i is a reply rule conditioned by beliefs. Put differently, it maps beliefs ν_i to a reply rule g_i :

$$\sigma_i : \Delta_{-i}^H \rightarrow \Delta_i^H : \nu_i \mapsto g_i.$$

So

$$\sigma_i \in (\Delta_i^H)^{\Delta_{-i}^H}.$$

Given a fixed ν_i , a behavioural strategy σ_i induces a reply rule g_i through

$$g_i =_{\text{Def}} \sigma_i(\nu_i).$$

Behavioural strategy and conditional forecast

- A behavioural strategy σ_i is a reply rule conditioned by beliefs. Put differently, it maps beliefs ν_i to a reply rule g_i :

$$\sigma_i : \Delta_{-i}^H \rightarrow \Delta_i^H : \nu_i \mapsto g_i.$$

So

$$\sigma_i \in (\Delta_i^H)^{\Delta_{-i}^H}.$$

Given a fixed ν_i , a behavioural strategy σ_i induces a reply rule g_i through

$$g_i =_{\text{Def}} \sigma_i(\nu_i).$$

- A conditional forecast τ_i is a forecast conditioned by beliefs.

Behavioural strategy and conditional forecast

- A **behavioural strategy** σ_i is a reply rule conditioned by beliefs. Put differently, it maps beliefs ν_i to a reply rule g_i :

$$\sigma_i : \Delta_{-i}^H \rightarrow \Delta_i^H : \nu_i \mapsto g_i.$$

So

$$\sigma_i \in (\Delta_i^H)^{\Delta_{-i}^H}.$$

Given a fixed ν_i , a behavioural strategy σ_i induces a reply rule g_i through

$$g_i =_{\text{Def}} \sigma_i(\nu_i).$$

- A **conditional forecast** τ_i is a forecast conditioned by beliefs. Put differently, it maps beliefs ν_i to a forecasting rule f_i :

$$\tau_i : \Delta_{-i}^H \rightarrow \Delta_{-i}^H : \nu_i \mapsto f_i.$$

Behavioural strategy and conditional forecast

- A behavioural strategy σ_i is a reply rule conditioned by beliefs. Put differently, it maps beliefs ν_i to a reply rule g_i :

$$\sigma_i : \Delta_{-i}^H \rightarrow \Delta_i^H : \nu_i \mapsto g_i.$$

So

$$\sigma_i \in (\Delta_i^H)^{\Delta_{-i}^H}.$$

Given a fixed ν_i , a behavioural strategy σ_i induces a reply rule g_i through

$$g_i =_{\text{Def}} \sigma_i(\nu_i).$$

- A conditional forecast τ_i is a forecast conditioned by beliefs. Put differently, it maps beliefs ν_i to a forecasting rule f_i :

$$\tau_i : \Delta_{-i}^H \rightarrow \Delta_{-i}^H : \nu_i \mapsto f_i.$$

So

$$\tau_i \in (\Delta_{-i}^H)^{\Delta_{-i}^H}.$$

Behavioural strategy and conditional forecast

- A **behavioural strategy** σ_i is a reply rule conditioned by beliefs. Put differently, it maps beliefs ν_i to a reply rule g_i :

$$\sigma_i : \Delta_{-i}^H \rightarrow \Delta_i^H : \nu_i \mapsto g_i.$$

So

$$\sigma_i \in (\Delta_i^H)^{\Delta_{-i}^H}.$$

Given a fixed ν_i , a behavioural strategy σ_i induces a reply rule g_i through

$$g_i =_{\text{Def}} \sigma_i(\nu_i).$$

- A **conditional forecast** τ_i is a forecast conditioned by beliefs. Put differently, it maps beliefs ν_i to a forecasting rule f_i :

$$\tau_i : \Delta_{-i}^H \rightarrow \Delta_{-i}^H : \nu_i \mapsto f_i.$$


So

$$\tau_i \in (\Delta_{-i}^H)^{\Delta_{-i}^H}.$$

Summary:

$$\begin{array}{c} H \times \Delta_{-i} \xrightarrow{\tau_i} \Delta_{-i} \\ \vdots \\ H \times \Delta_{-i} \xrightarrow{\sigma_i} \Delta_i \end{array}$$

Behav. strategy and belief of opponent's behav. strategy



Fictitious play compared to Bayesian play

Fictitious play compared to Bayesian play

- A forecasting rule for player i is a function that maps every history to a probability distribution over counter-action profiles.

Fictitious play compared to Bayesian play

- A forecasting rule for player i is a function that maps every history to a probability distribution over counter-action profiles.
- With fictitious play, the forecasting function, f , is

$$f_i(h) =_{Def} \prod_{j \neq i} \phi^j.$$

Fictitious play compared to Bayesian play

- A forecasting rule for player i is a function that maps every history to a probability distribution over counter-action profiles.
- With fictitious play, the forecasting function, f , is

$$f_i(h) =_{Def} \prod_{j \neq i} \phi^j.$$


- With Bayesian play, forecasting is much more subtler.

Fictitious play compared to Bayesian play

- A forecasting rule for player i is a function that maps every history to a probability distribution over counter-action profiles.
- With fictitious play, the forecasting function, f , is

$$f_i(h) =_{Def} \prod_{j \neq i} \phi^j.$$

- With Bayesian play, forecasting is much more subtler. There, a forecast is a probability distribution over the reply rules of other players.



Fictitious play compared to Bayesian play

Fictitious play compared to Bayesian play

Fictitious play

Fictitious play compared to Bayesian play

Fictitious play

- The behaviour of every opponent is modelled by (or: projected on) **a single mixed strategy**.

Fictitious play compared to Bayesian play

Fictitious play

- The behaviour of every opponent is modelled by (or: projected on) **a single mixed strategy**.
- The next move is simply a (myopic) best reply to this mixed strategy.

Fictitious play compared to Bayesian play

Fictitious play

- The behaviour of every opponent is modelled by (or: projected on) **a single mixed strategy**.
- The next move is simply a (myopic) best reply to this mixed strategy.

Bayesian play

Fictitious play compared to Bayesian play

Fictitious play

- The behaviour of every opponent is modelled by (or: projected on) **a single mixed strategy**.
- The next move is simply a (myopic) best reply to this mixed strategy.

Bayesian play

- Opponents are modelled by **a probability distribution over reply rules** $H \rightarrow \Delta(X_j)$

Fictitious play compared to Bayesian play

Fictitious play

- The behaviour of every opponent is modelled by (or: projected on) **a single mixed strategy**.
- The next move is simply a (myopic) best reply to this mixed strategy.

Bayesian play

- Opponents are modelled by **a probability distribution over reply rules** $H \rightarrow \Delta(X_j)$:

$$v_i \in \Delta[\prod_{j \neq i} \Delta(X_j)^H] \quad (5)$$

Fictitious play compared to Bayesian play

Fictitious play

- The behaviour of every opponent is modelled by (or: projected on) **a single mixed strategy**.
- The next move is simply a (myopic) best reply to this mixed strategy.

Bayesian play

- Opponents are modelled by **a probability distribution over reply rules** $H \rightarrow \Delta(X_j)$:

$$v_i \in \Delta[\prod_{j \neq i} \Delta(X_j)^H] \quad (5)$$

(B^A means: all functions from A to B .)

Fictitious play compared to Bayesian play

Fictitious play

- The behaviour of every opponent is modelled by (or: projected on) **a single mixed strategy**.
- The next move is simply a (myopic) best reply to this mixed strategy.

Bayesian play

- Opponents are modelled by **a probability distribution over reply rules** $H \rightarrow \Delta(X_j)$:

$$\nu_i \in \Delta[\prod_{j \neq i} \Delta(X_j)^H] \quad (5)$$

(B^A means: all functions from A to B .)

- Every player maintains **behavioural strategy** that maps beliefs ν_i onto reply rules $g_i : H \rightarrow \Delta(X_i)$.

Fictitious play compared to Bayesian play

Fictitious play

- The behaviour of every opponent is modelled by (or: projected on) **a single mixed strategy**.
- The next move is simply a (myopic) best reply to this mixed strategy.

Bayesian play

- Opponents are modelled by **a probability distribution over reply rules** $H \rightarrow \Delta(X_j)$:

$$\nu_i \in \Delta[\prod_{j \neq i} \Delta(X_j)^H] \quad (5)$$

(B^A means: all functions from A to B .)

- Every player maintains **behavioural strategy** that maps beliefs ν_i onto reply rules $g_i : H \rightarrow \Delta(X_i)$.

(The reply rule g_i may for instance give a best response based on ν_i in the repeated game.)

Fictitious play compared to Bayesian play

Fictitious play

- The behaviour of every opponent is modelled by (or: projected on) **a single mixed strategy**.
- The next move is simply a (myopic) best reply to this mixed strategy.

Bayesian play

- Opponents are modelled by **a probability distribution over reply rules** $H \rightarrow \Delta(X_j)$:

$$\nu_i \in \Delta[\prod_{j \neq i} \Delta(X_j)^H] \quad (5)$$

(B^A means: all functions from A to B .)

- Every player maintains **behavioural strategy** that maps beliefs ν_i onto reply rules $g_i : H \rightarrow \Delta(X_i)$.

(The reply rule g_i may for instance give a best response based on ν_i in the repeated game.)

- Beliefs ν_i are maintained through Bayesian updating.

Part IV:

True distribution of play vs. subjective distribution of play

The true distribution of play μ on H

The reply rules together induce a true distribution of play, μ , on Ω , as follows.

The true distribution of play μ on H

The reply rules together induce a true distribution of play, μ , on Ω , as follows.

■ Let $g \in \Pi_{i=1}^n[\Delta_i^H]$ be the actual reply profile.

The true distribution of play μ on H

The reply rules together induce a true distribution of play, μ , on Ω , as follows.

- Let $g \in \Pi_{i=1}^n[\Delta_i^H]$ be the actual reply profile.
- If we establish a bijection

$$\Pi_{i=1}^n[\Delta_i^H] \sim [\Pi_{i=1}^n \Delta_i]^H$$

(exercise) and write the latter set as Δ^H , we get rid of repetitions of histories in the function argument.

The true distribution of play μ on H

The reply rules together induce a true distribution of play, μ , on Ω , as follows.

- Let $g \in \Pi_{i=1}^n[\Delta_i^H]$ be the actual reply profile.
- If we establish a bijection

$$\Pi_{i=1}^n[\Delta_i^H] \sim [\Pi_{i=1}^n \Delta_i]^H$$

(exercise) and write the latter set as Δ^H , we get rid of repetitions of histories in the function argument.

- The reply profile $g \in \Delta^H$ induces a probability distribution μ on H , inductively:

$$\mu\{hx\} = \begin{cases} 1 \cdot g(x) & \text{if } h = \emptyset, \\ \mu\{h\} \cdot g(x|h) & \text{otherwise.} \end{cases}$$

The probability measure μ on H , on its turn, induces a probability measure on (a σ -algebra of) Ω .

How to extend (any) μ on H to Ω

Definition. (Correspondence between H and Ω .)

How to extend (any) μ on H to Ω

Definition. (Correspondence between H and Ω .)

\Rightarrow Given history $h \in H$, define the set of **continuations** of h as

$$C(h) =_{Def} \{ \omega \in \Omega \mid \omega|_{|h|} = h \}$$

How to extend (any) μ on H to Ω

Definition. (Correspondence between H and Ω .)

\Rightarrow Given history $h \in H$, define the set of **continuations** of h as

$$C(h) =_{Def} \{ \omega \in \Omega \mid \omega|_{|h|} = h \}$$

Immediately $\{ C(h) \mid h \in H \} = \Omega$.

How to extend (any) μ on H to Ω

Definition. (Correspondence between H and Ω .)

\Rightarrow Given history $h \in H$, define the set of **continuations** of h as

$$C(h) =_{Def} \{ \omega \in \Omega \mid \omega|_{|h|} = h \}$$

Immediately $\{ C(h) \mid h \in H \} = \Omega$.

\Leftarrow Given a realisation $\omega \in \Omega$, define the **restriction** $\omega|_t \in H$ as the initial part of ω with length t .

How to extend (any) μ on H to Ω

Definition. (Correspondence between H and Ω .)

\Rightarrow Given history $h \in H$, define the set of **continuations** of h as

$$C(h) =_{Def} \{ \omega \in \Omega \mid \omega|_{|h|} = h \}$$

Immediately $\{ C(h) \mid h \in H \} = \Omega$.

\Leftarrow Given a realisation $\omega \in \Omega$, define the **restriction** $\omega|_t \in H$ as the initial part of ω with length t .

The measure μ on H induces a “pre-measure” μ' on $C(H) = \Omega$ through

$$\mu'\{C(h_t)\} =_{Def} \mu\{h_t\}.$$

How to extend (any) μ on H to Ω

Definition. (Correspondence between H and Ω .)

\Rightarrow Given history $h \in H$, define the set of **continuations** of h as

$$C(h) =_{\text{Def}} \{ \omega \in \Omega \mid \omega|_{|h|} = h \}$$

Immediately $\{ C(h) \mid h \in H \} = \Omega$.

\Leftarrow Given a realisation $\omega \in \Omega$, define the **restriction** $\omega|_t \in H$ as the initial part of ω with length t .

The measure μ on H induces a “pre-measure” μ' on $C(H) = \Omega$ through

$$\mu'\{C(h_t)\} =_{\text{Def}} \mu\{h_t\}.$$

Standard results in probability theory¹ now imply that μ' can be extended to a proper probability measure on (a σ -algebra of) Ω .

¹Carathéodory's extension theorem.

Player i 's subjective distribution of play μ_i on H

Player i 's subjective distribution of play μ_i on H

- The beliefs on reply rules of other players, ν_i , together with i 's own reply rule g_i , **also** forms some sort of “reply profile”.
E.g., $(\nu_2^1, g_2, \nu_2^3, \dots, \nu_2^n)$ represents the further course of play as Player 2 sees it.

Player i 's subjective distribution of play μ_i on H

- The beliefs on reply rules of other players, ν_i , together with i 's own reply rule g_i , **also** forms some sort of “reply profile”.
E.g., $(\nu_2^1, g_2, \nu_2^3, \dots, \nu_2^n)$ represents **the further course of play as Player 2 sees it**.
- We have just seen that the vector of all reply rules, g , generates a **true distribution of play**, μ , on Ω .

Player i 's subjective distribution of play μ_i on H

- The beliefs on reply rules of other players, v_i , together with i 's own reply rule g_i , **also** forms some sort of “reply profile”.
E.g., $(v_2^1, g_2, v_2^3, \dots, v_2^n)$ represents **the further course of play as Player 2 sees it**.
- We have just seen that the vector of all reply rules, g , generates a **true distribution of play**, μ , on Ω .
- Similarly, players i 's conditionalised predictive learning rule, (v_i, g_i) , generates a **subjective distribution of play**, μ_i , on Ω :

$$\begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \vdots \\ \mu_n \end{pmatrix} \text{ is generated by } (v, g) = \begin{pmatrix} g_1 & v_1^2 & v_1^3 & \dots & v_1^n \\ v_2^1 & g_2 & v_2^3 & \dots & v_2^n \\ v_3^1 & v_3^2 & g_3 & \dots & v_3^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ v_n^1 & v_n^2 & v_n^3 & \dots & g_n \end{pmatrix}.$$

Part V: Main results

The subjective view on distribution of play

Given ω , we would like to say something like:

The subjective view on distribution of play

Given ω , we would like to say something like:

First stab at the formulation of a theorem:

If $t \rightarrow \infty$, then i 's subjective view on play at t , namely,

$$\mu_i\{\cdot \mid \text{realised play until } t\},$$

approximates the true distribution of play at t , namely,

$$\mu\{\cdot \mid \text{realised play until } t\}.$$

Domination of probability measures

Definition. Let μ and τ be probability distributions. τ is said to **dominate** μ , written $\mu \ll \tau$, if $\tau\{E\} > 0$ whenever $\mu\{E\} > 0$.

Domination of probability measures

Definition. Let μ and τ be probability distributions. τ is said to **dominate** μ , written $\mu \ll \tau$, if $\tau\{E\} > 0$ whenever $\mu\{E\} > 0$.

If $\mu \ll \tau \ll \mu$, then μ and τ are said to be **equivalent**.

Domination of probability measures

Definition. Let μ and τ be probability distributions. τ is said to **dominate** μ , written $\mu \ll \tau$, if $\tau\{E\} > 0$ whenever $\mu\{E\} > 0$.

If $\mu \ll \tau \ll \mu$, then μ and τ are said to be **equivalent**.

Definition. If

$$\mu \ll \mu_i,$$

i.e., if Player i assigns positive probabilities (however small) to every realisation deemed possible by μ , then i 's beliefs are said to contain a **grain of truth** on actual play.

Domination of probability measures

Definition. Let μ and τ be probability distributions. τ is said to **dominate** μ , written $\mu \ll \tau$, if $\tau\{E\} > 0$ whenever $\mu\{E\} > 0$.

If $\mu \ll \tau \ll \mu$, then μ and τ are said to be **equivalent**.

Definition. If

$$\mu \ll \mu_i,$$

i.e., if Player i assigns positive probabilities (however small) to every realisation deemed possible by μ , then i 's beliefs are said to contain a **grain of truth** on actual play.

Used by Kalai *et al.*, 1993:

Domination of probability measures

Definition. Let μ and τ be probability distributions. τ is said to **dominate** μ , written $\mu \ll \tau$, if $\tau\{E\} > 0$ whenever $\mu\{E\} > 0$.

If $\mu \ll \tau \ll \mu$, then μ and τ are said to be **equivalent**.

Definition. If

$$\mu \ll \mu_i,$$

i.e., if Player i assigns positive probabilities (however small) to every realisation deemed possible by μ , then i 's beliefs are said to contain a **grain of truth** on actual play.

Used by Kalai *et al.*, 1993:

Theorem (Radon-Nikodym). *Given X and σ -finite probability measures μ and ν on X such that $\nu \ll \mu$*

Domination of probability measures

Definition. Let μ and τ be probability distributions. τ is said to **dominate** μ , written $\mu \ll \tau$, if $\tau\{E\} > 0$ whenever $\mu\{E\} > 0$.

If $\mu \ll \tau \ll \mu$, then μ and τ are said to be **equivalent**.

Definition. If

$$\mu \ll \mu_i,$$

i.e., if Player i assigns positive probabilities (however small) to every realisation deemed possible by μ , then i 's beliefs are said to contain a **grain of truth** on actual play.

Used by Kalai *et al.*, 1993:

Theorem (Radon-Nikodym). *Given X and σ -finite probability measures μ and ν on X such that $\nu \ll \mu$, then there is a measurable function $\psi : X \rightarrow [0, \infty)$, such that for all μ -measurable sets E ,*

$$\nu\{E\} = \int_E \psi d\mu.$$

Domination of probability measures

Definition. Let μ and τ be probability distributions. τ is said to **dominate** μ , written $\mu \ll \tau$, if $\tau\{E\} > 0$ whenever $\mu\{E\} > 0$.

If $\mu \ll \tau \ll \mu$, then μ and τ are said to be **equivalent**.

Definition. If

$$\mu \ll \mu_i,$$

i.e., if Player i assigns positive probabilities (however small) to every realisation deemed possible by μ , then i 's beliefs are said to contain a **grain of truth** on actual play.

Used by Kalai *et al.*, 1993:

Theorem (Radon-Nikodym). *Given X and σ -finite probability measures μ and ν on X such that $\nu \ll \mu$, then there is a measurable function $\psi : X \rightarrow [0, \infty)$, such that for all μ -measurable sets E ,*

$$\nu\{E\} = \int_E \psi d\mu.$$

So, if i 's beliefs contain a grain of truth on actual play, then μ_i is, roughly put, “as expressive as” μ .

Theorem of Blackwell and Dubins

Definition. Let μ on Ω represent the true distribution of play. Another measure, μ' on Ω , is said to **merge with the true distribution of play** if, for almost all play path ω

$$\lim_{t \rightarrow \infty} \| \mu\{\cdot \mid C[\omega|_t]\} - \mu'\{\cdot \mid C[\omega|_t]\} \|^2 = 0.$$

Theorem of Blackwell and Dubins

Definition. Let μ on Ω represent the true distribution of play. Another measure, μ' on Ω , is said to **merge with the true distribution of play** if, for almost all play path ω

$$\lim_{t \rightarrow \infty} \| \mu\{\cdot \mid C[\omega|_t]\} - \mu'\{\cdot \mid C[\omega|_t]\} \|^2 = 0.$$

Remarks:

Theorem of Blackwell and Dubins

Definition. Let μ on Ω represent the true distribution of play. Another measure, μ' on Ω , is said to **merge with the true distribution of play** if, for almost all play path ω

$$\lim_{t \rightarrow \infty} \| \mu\{\cdot \mid C[\omega|_t]\} - \mu'\{\cdot \mid C[\omega|_t]\} \|^2 = 0.$$

Remarks:

■ $C[\omega|_t] \in \Omega$ consists of all possible continuations of $\omega|_t \in H$.

Theorem of Blackwell and Dubins

Definition. Let μ on Ω represent the true distribution of play. Another measure, μ' on Ω , is said to **merge with the true distribution of play** if, for almost all play path ω

$$\lim_{t \rightarrow \infty} \| \mu\{\cdot \mid C[\omega|_t]\} - \mu'\{\cdot \mid C[\omega|_t]\} \|^2 = 0.$$

Remarks:

- $C[\omega|_t] \in \Omega$ consists of all possible continuations of $\omega|_t \in H$.
- It is not possible to conditionalise on histories, because $H \cap \Omega = \emptyset$.

Theorem of Blackwell and Dubins

Definition. Let μ on Ω represent the true distribution of play. Another measure, μ' on Ω , is said to **merge with the true distribution of play** if, for almost all play path ω

$$\lim_{t \rightarrow \infty} \| \mu\{\cdot \mid C[\omega|_t]\} - \mu'\{\cdot \mid C[\omega|_t]\} \|^2 = 0.$$

Remarks:

- $C[\omega|_t] \in \Omega$ consists of all possible continuations of $\omega|_t \in H$.
- It is not possible to conditionalise on histories, because $H \cap \Omega = \emptyset$.
- However, for each $\omega \in \Omega$ we have $C[\omega|_t] \subseteq \Omega$ is measurable.

Theorem of Blackwell and Dubins

Definition. Let μ on Ω represent the true distribution of play. Another measure, μ' on Ω , is said to **merge with the true distribution of play** if, for almost all play path ω

$$\lim_{t \rightarrow \infty} \| \mu\{\cdot \mid C[\omega|_t]\} - \mu'\{\cdot \mid C[\omega|_t]\} \|^2 = 0.$$

Remarks:

- $C[\omega|_t] \in \Omega$ consists of all possible continuations of $\omega|_t \in H$.
- It is not possible to conditionalise on histories, because $H \cap \Omega = \emptyset$.
- However, for each $\omega \in \Omega$ we have $C[\omega|_t] \subseteq \Omega$ is measurable.

Theorem (Blackwell and Dubins, 1962). *Let μ represent the true distribution of play, and let μ_i represent i 's view on the distribution of play. If $\mu \ll \mu_i$, then i 's beliefs merge with the true distribution of play.*

Good prediction vs. very good prediction

Let $\omega \in \Omega$ be given. At round t ,

Good prediction vs. very good prediction

Let $\omega \in \Omega$ be given. At round t ,

- Player i 's predictive learning rule yields a forecast $p^{-i} \in \Delta_{-i}$.

Good prediction vs. very good prediction

Let $\omega \in \Omega$ be given. At round t ,

- Player i 's predictive learning rule yields a forecast $p^{-i} \in \Delta_{-i}$.
- The actual reply rules g induce the distribution proper, $q^{-i} \in \Delta_{-i}$.

Good prediction vs. very good prediction

Let $\omega \in \Omega$ be given. At round t ,

- Player i 's predictive learning rule yields a forecast $p^{-i} \in \Delta_{-i}$.
- The actual reply rules g induce the distribution proper, $q^{-i} \in \Delta_{-i}$.

Notice that both p^{-i}, q^{-i} conditionalise on histories.

Good prediction vs. very good prediction

Let $\omega \in \Omega$ be given. At round t ,

- Player i 's predictive learning rule yields a forecast $p^{-i} \in \Delta_{-i}$.
- The actual reply rules g induce the distribution proper, $q^{-i} \in \Delta_{-i}$.

Notice that both p^{-i}, q^{-i} conditionalise on histories.

- Player i is said to be a **good predictor** on ω if the mean square error of prediction goes to zero a.s.:

$$\lim_{t \rightarrow \infty} \frac{1}{t} \sum_{s \leq t} \|q^{-i(s+1)} - p^{-i(s+1)}\|^2 = 0$$

Good prediction vs. very good prediction

Let $\omega \in \Omega$ be given. At round t ,

- Player i 's predictive learning rule yields a forecast $p^{-i} \in \Delta_{-i}$.
- The actual reply rules g induce the distribution proper, $q^{-i} \in \Delta_{-i}$.

Notice that both p^{-i}, q^{-i} conditionalise on histories.

- Player i is said to be a **good predictor** on ω if the mean square error of prediction goes to zero a.s.:

$$\lim_{t \rightarrow \infty} \frac{1}{t} \sum_{s \leq t} \|q^{-i(s+1)} - p^{-i(s+1)}\|^2 = 0$$

- Player i is said to be a **very good predictor** on ω if the actual square error of prediction goes to zero a.s.:

$$\lim_{t \rightarrow \infty} \|q^{-i(t+1)} - p^{-i(t+1)}\|^2 = 0$$

Good prediction vs. very good prediction

Let $\omega \in \Omega$ be given. At round t ,

- Player i 's predictive learning rule yields a forecast $p^{-i} \in \Delta_{-i}$.
- The actual reply rules g induce the distribution proper, $q^{-i} \in \Delta_{-i}$.

Notice that both p^{-i}, q^{-i} conditionalise on histories.

- Player i is said to be a **good predictor** on ω if the mean square error of prediction goes to zero a.s.:

$$\lim_{t \rightarrow \infty} \frac{1}{t} \sum_{s \leq t} \|q^{-i(s+1)} - p^{-i(s+1)}\|^2 = 0$$

- Player i is said to be a **very good predictor** on ω if the actual square error of prediction goes to zero a.s.:

$$\lim_{t \rightarrow \infty} \|q^{-i(t+1)} - p^{-i(t+1)}\|^2 = 0$$

- Very good predictor \Rightarrow good predictor.

Good prediction vs. very good prediction

Let $\omega \in \Omega$ be given. At round t ,

- Player i 's predictive learning rule yields a forecast $p^{-i} \in \Delta_{-i}$.
- The actual reply rules g induce the distribution proper, $q^{-i} \in \Delta_{-i}$.

Notice that both p^{-i}, q^{-i} conditionalise on histories.

- Player i is said to be a **good predictor** on ω if the mean square error of prediction goes to zero a.s.:

$$\lim_{t \rightarrow \infty} \frac{1}{t} \sum_{s \leq t} \|q^{-i(s+1)} - p^{-i(s+1)}\|^2 = 0$$

- Player i is said to be a **very good predictor** on ω if the actual square error of prediction goes to zero a.s.:

$$\lim_{t \rightarrow \infty} \|q^{-i(t+1)} - p^{-i(t+1)}\|^2 = 0$$

- Very good predictor \Rightarrow good predictor.
- From (Blackwell and Dubins, 1962) the following can be proven.

Corollary. If $\mu \ll \mu_i$ then i is a very good predictor a.s.

Very good prediction \nRightarrow all μ_i dominate μ

Very good prediction \nRightarrow all μ_i dominate μ


- Suppose a 2×2 game where column plays $\Pr\{L\} = p^* = 2/5$.

Very good prediction \nRightarrow all μ_i dominate μ

- Suppose a 2×2 game where column plays $\Pr\{L\} = p^* = 2/5$.
- Suppose row maintains a Beta-distribution with prior $\text{Beta}(1, 1)$ (which is the uniform distribution on $[0, 1]$).


Very good prediction \Rightarrow all μ_i dominate μ

- Suppose a 2×2 game where column plays $\Pr\{L\} = p^* = 2/5$.
- Suppose row maintains a Beta-distribution with prior Beta(1,1) (which is the uniform distribution on $[0, 1]$).
- Almost surely, the posterior at time t is Beta($2t + 1, 3t + 1$), so becomes a **very good predictor** of p^* a.s., like so:



Very good prediction \nRightarrow all μ_i dominate μ

- Suppose a 2×2 game where column plays $\Pr\{L\} = p^* = 2/5$.
- Suppose row maintains a Beta-distribution with prior Beta(1,1) (which is the uniform distribution on $[0, 1]$).
- Almost surely, the posterior at time t is Beta($2t + 1, 3t + 1$), so becomes a **very good predictor** of p^* a.s., like so:




- However by the strong law of large numbers,

$$\mu_{\text{true}}\{\text{column's empirical distr. is } p^*\} = 1$$

Very good prediction \nRightarrow all μ_i dominate μ

- Suppose a 2×2 game where column plays $\Pr\{L\} = p^* = 2/5$.
- Suppose row maintains a Beta-distribution with prior Beta(1,1) (which is the uniform distribution on $[0, 1]$).
- Almost surely, the posterior at time t is Beta($2t + 1, 3t + 1$), so becomes a **very good predictor** of p^* a.s., like so:



- However by the strong law of large numbers,

$$\mu_{\text{true}}\{\text{column's empirical distr. is } p^*\} = 1$$

(while, at any one time $\mu_{\text{row}}\{\text{column's empirical distr. is } p^*\} = 0$).

ϵ -closeness of two probability measures

ϵ -closeness of two probability measures

Definition. Let μ and τ probability distributions. μ and τ are said to be ϵ -close, written $\mu \sim_{\epsilon} \tau$, if for some $Y \subseteq X$ we have $\mu\{Y\} > 1 - \epsilon$ and $\tau\{Y\} > 1 - \epsilon$ and for all measurable $E \subseteq Y$:

$$(1 - \epsilon)\tau\{E\} \leq \mu\{E\} \leq (1 + \epsilon)\tau\{E\}.$$

ϵ -closeness of two probability measures

Definition. Let μ and τ probability distributions. μ and τ are said to be ϵ -close, written $\mu \sim_{\epsilon} \tau$, if for some $Y \subseteq X$ we have $\mu\{Y\} > 1 - \epsilon$ and $\tau\{Y\} > 1 - \epsilon$ and for all measurable $E \subseteq Y$:

$$(1 - \epsilon)\tau\{E\} \leq \mu\{E\} \leq (1 + \epsilon)\tau\{E\}.$$

- ϵ -closeness is a strong condition. It demands, for all events E , even the most improbable ones, that probabilities may differ only $(1 - \epsilon)\%$.

ϵ -closeness of two probability measures

Definition. Let μ and τ probability distributions. μ and τ are said to be ϵ -close, written $\mu \sim_{\epsilon} \tau$, if for some $Y \subseteq X$ we have $\mu\{Y\} > 1 - \epsilon$ and $\tau\{Y\} > 1 - \epsilon$ and for all measurable $E \subseteq Y$:

$$(1 - \epsilon)\tau\{E\} \leq \mu\{E\} \leq (1 + \epsilon)\tau\{E\}.$$

- ϵ -closeness is a strong condition. It demands, for all events E , **even the most improbable ones**, that probabilities may differ only $(1 - \epsilon)\%$.
- ϵ -closeness is stronger than absolute closeness. To demand that probabilities may differ only ϵ is much weaker!

ϵ -closeness of two probability measures

Definition. Let μ and τ probability distributions. μ and τ are said to be ϵ -close, written $\mu \sim_{\epsilon} \tau$, if for some $Y \subseteq X$ we have $\mu\{Y\} > 1 - \epsilon$ and $\tau\{Y\} > 1 - \epsilon$ and for all measurable $E \subseteq Y$:

$$(1 - \epsilon)\tau\{E\} \leq \mu\{E\} \leq (1 + \epsilon)\tau\{E\}.$$

- ϵ -closeness is a strong condition. It demands, for all events E , **even the most improbable ones**, that probabilities may differ only $(1 - \epsilon)\%$.
- ϵ -closeness is stronger than absolute closeness. To demand that probabilities may differ only ϵ is much weaker!

Example. Let $\epsilon = 0.01$, $\mu\{E\} = 0.001$ and $\tau\{E\} = 0.009$. Then $|\mu\{E\} - \tau\{E\}| < \epsilon$ while $\tau\{E\} = 9\mu\{E\}$.

ϵ -closeness of two probability measures

Definition. Let μ and τ probability distributions. μ and τ are said to be ϵ -close, written $\mu \sim_{\epsilon} \tau$, if for some $Y \subseteq X$ we have $\mu\{Y\} > 1 - \epsilon$ and $\tau\{Y\} > 1 - \epsilon$ and for all measurable $E \subseteq Y$:

$$(1 - \epsilon)\tau\{E\} \leq \mu\{E\} \leq (1 + \epsilon)\tau\{E\}.$$

- ϵ -closeness is a strong condition. It demands, for all events E , **even the most improbable ones**, that probabilities may differ only $(1 - \epsilon)\%$.
- ϵ -closeness is stronger than absolute closeness. To demand that probabilities may differ only ϵ is much weaker!

Example. Let $\epsilon = 0.01$, $\mu\{E\} = 0.001$ and $\tau\{E\} = 0.009$. Then $|\mu\{E\} - \tau\{E\}| < \epsilon$ while $\tau\{E\} = 9\mu\{E\}$.

- Thus, being ϵ -close means that not only does the player assess the future correctly, *it even assesses developments following unlikely histories correctly.*

Main theorem of Kalai and Lehrer

Definition. Let μ and μ' be distributions of play and $\epsilon > 0$. Then μ' is said to ϵ -play like μ if u' is ϵ -close to μ .

Main theorem of Kalai and Lehrer

Definition. Let μ and μ' be distributions of play and $\epsilon > 0$. Then μ' is said to ϵ -play like μ if μ' is ϵ -close to μ .

Theorem (Kalai and Lehrer, 1993). Let μ represent the true distribution of play, and let μ_i represent i 's view on the distribution of play. If $\mu \ll \mu_i$, then μ_i ϵ -plays like μ for any $\epsilon > 0$.

Main theorem of Kalai and Lehrer

Definition. Let μ and μ' be distributions of play and $\epsilon > 0$. Then μ' is said to ϵ -play like μ if μ' is ϵ -close to μ .

Theorem (Kalai and Lehrer, 1993). Let μ represent the true distribution of play, and let μ_i represent i 's view on the distribution of play. If $\mu \ll \mu_i$, then μ_i ϵ -plays like μ for any $\epsilon > 0$.

So after history $\omega|_t$, the probability distribution over future play is ϵ -close to whatever Player i believes the distribution is.

Main theorem of Kalai and Lehrer

Definition. Let μ and μ' be distributions of play and $\epsilon > 0$. Then μ' is said to ϵ -play like μ if u' is ϵ -close to μ .

Theorem (Kalai and Lehrer, 1993). Let μ represent the true distribution of play, and let μ_i represent i 's view on the distribution of play. If $\mu \ll \mu_i$, then μ_i ϵ -plays like μ for any $\epsilon > 0$.

So after history $\omega|_t$, the probability distribution over future play is ϵ -close to whatever Player i believes the distribution is.

■ Other than $\mu \ll \mu_i$, no other assumptions are made on μ and μ_i .

Main theorem of Kalai and Lehrer

Definition. Let μ and μ' be distributions of play and $\epsilon > 0$. Then μ' is said to ϵ -play like μ if u' is ϵ -close to μ .

Theorem (Kalai and Lehrer, 1993). Let μ represent the true distribution of play, and let μ_i represent i 's view on the distribution of play. If $\mu \ll \mu_i$, then μ_i ϵ -plays like μ for any $\epsilon > 0$.

So after history $\omega|_t$, the probability distribution over future play is ϵ -close to whatever Player i believes the distribution is.

- Other than $\mu \ll \mu_i$, no other assumptions are made on μ and μ_i .
- This theorem implies closeness of probabilities also for courses of play that are extremely unlikely.

Main theorem of Kalai and Lehrer

Definition. Let μ and μ' be distributions of play and $\epsilon > 0$. Then μ' is said to ϵ -play like μ if u' is ϵ -close to μ .

Theorem (Kalai and Lehrer, 1993). Let μ represent the true distribution of play, and let μ_i represent i 's view on the distribution of play. If $\mu \ll \mu_i$, then μ_i ϵ -plays like μ for any $\epsilon > 0$.

So after history $\omega|_t$, the probability distribution over future play is ϵ -close to whatever Player i believes the distribution is.

- Other than $\mu \ll \mu_i$, no other assumptions are made on μ and μ_i .
- This theorem implies closeness of probabilities also for courses of play that are extremely unlikely.
- The theorem does **not** state that a player learns to predict other players' **off path strategies**. (Recall Player 2's beliefs in unforgiving constellation.)

What next?

Emergence of social conventions.

What next?

Emergence of social conventions.

- It is studied how (typically large) agent populations that use different types of currency (e.g., Dollar, Euro and Yen) or standards (Red, Green and Blue) may, through **sampling**, converge to a single currency.

What next?

Emergence of social conventions.

- It is studied how (typically large) agent populations that use different types of currency (e.g., Dollar, Euro and Yen) or standards (Red, Green and Blue) may, through **sampling**, converge to a single currency.
- Emergence has more to do with **adaptation** than with learning.

What next?

Emergence of social conventions.

- It is studied how (typically large) agent populations that use different types of currency (e.g., Dollar, Euro and Yen) or standards (Red, Green and Blue) may, through **sampling**, converge to a single currency.
- Emergence has more to do with **adaptation** than with learning.
- Emergence can be described as a **discrete Markov process**.

What next?

Emergence of social conventions.

- It is studied how (typically large) agent populations that use different types of currency (e.g., Dollar, Euro and Yen) or standards (Red, Green and Blue) may, through **sampling**, converge to a single currency.
- Emergence has more to do with **adaptation** than with learning.
- Emergence can be described as a **discrete Markov process**.
 - It be analysed *qualitatively* (by proving convergence).

What next?

Emergence of social conventions.

- It is studied how (typically large) agent populations that use different types of currency (e.g., Dollar, Euro and Yen) or standards (Red, Green and Blue) may, through **sampling**, converge to a single currency.
- Emergence has more to do with **adaptation** than with learning.
- Emergence can be described as a **discrete Markov process**.
 - It be analysed *qualitatively* (by proving convergence).
 - It be analysed *quantitatively* (by indicating the rate of convergence). This is much harder.

In Chapter 22 of the *Handbook on Computational Economics* (2006), Young describes how social dynamics can drift to so-called **stochastically stable states**, provided individuals act rationally most of the time.

The following slides were not used.

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

The set of all functions from A to B
is denoted by B^A .

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

The set of all functions from A to B is denoted by B^A .

Look at the Cartesian product B^n . This is a special case. Every tuple $b \in B^n$ corresponds to a function

$$f_b : \{0, \dots, n-1\} \rightarrow B,$$

and conversely, in the following way:

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

The set of all functions from A to B is denoted by B^A .

Look at the Cartesian product B^n . This is a special case. Every tuple $b \in B^n$ corresponds to a function

$$f_b : \{0, \dots, n-1\} \rightarrow B,$$

and conversely, in the following way:

\Leftarrow Given $b \in B^n$, define f_b by $f_b(i) =_{Def} b_i$.

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

The set of all functions from A to B is denoted by B^A .

Look at the Cartesian product B^n . This is a special case. Every tuple $b \in B^n$ corresponds to a function

$$f_b : \{0, \dots, n-1\} \rightarrow B,$$

and conversely, in the following way:

\Leftarrow Given $b \in B^n$, define f_b by $f_b(i) =_{\text{Def}} b_i$.

\Rightarrow Given f_b , define $b \in B^n$ by $b = (f_b(0), \dots, f_b(n-1))$.

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

The set of all functions from A to B is denoted by B^A .

Look at the Cartesian product B^n . This is a special case. Every tuple $b \in B^n$ corresponds to a function

$$f_b : \{0, \dots, n-1\} \rightarrow B,$$

and conversely, in the following way:

\Leftarrow Given $b \in B^n$, define f_b by $f_b(i) =_{\text{Def}} b_i$.

\Rightarrow Given f_b , define $b \in B^n$ by $b = (f_b(0), \dots, f_b(n-1))$.

Clearly,

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

The set of all functions from A to B is denoted by B^A .

Look at the Cartesian product B^n . This is a special case. Every tuple $b \in B^n$ corresponds to a function

$$f_b : \{0, \dots, n-1\} \rightarrow B,$$

and conversely, in the following way:

\Leftarrow Given $b \in B^n$, define f_b by $f_b(i) =_{\text{Def}} b_i$.

\Rightarrow Given f_b , define $b \in B^n$ by $b = (f_b(0), \dots, f_b(n-1))$.

Clearly, The mappings are each

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

The set of all functions from A to B is denoted by B^A .

Look at the Cartesian product B^n . This is a special case. Every tuple $b \in B^n$ corresponds to a function

$$f_b : \{0, \dots, n-1\} \rightarrow B,$$

and conversely, in the following way:

\Leftarrow Given $b \in B^n$, define f_b by $f_b(i) =_{\text{Def}} b_i$.

\Rightarrow Given f_b , define $b \in B^n$ by $b = (f_b(0), \dots, f_b(n-1))$.

Clearly, The mappings are each

other's inverse. Both mappings are surjective.

Further, fundamental math defines 0 as \emptyset , and $n+1$ as $n \cup \{n\}$:

$$0 = \emptyset$$

$$1 = 0 \cup \{0\} = \emptyset \cup \{\emptyset\} = \{\emptyset\}$$

$$2 = 1 \cup \{1\} = \{\emptyset, \{\emptyset\}\}$$

$$3 = 2 \cup \{2\} = \{\emptyset, \{\emptyset\}, \{\{\emptyset\}\}\}$$

$$\vdots \quad \vdots$$

So

$$B^n = B^{\{0,1,\dots,n-1\}}$$

$$= \{f \mid f : \{0,1,\dots,n-1\} \rightarrow B\}.$$

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

so that

$$\begin{aligned} B^n &= \{f \mid f : \{0, \dots, n-1\} \rightarrow B\} \\ &= \{f \mid f : n \rightarrow B\}. \end{aligned}$$

We can now generalise to

$$B^A =_{Def} \{f \mid f : A \rightarrow B\}.$$

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

so that

$$\begin{aligned} B^n &= \{f \mid f : \{0, \dots, n-1\} \rightarrow B\} \\ &= \{f \mid f : n \rightarrow B\}. \end{aligned}$$

Later, we will use the following identities:

We can now generalise to

$$B^A =_{Def} \{f \mid f : A \rightarrow B\}.$$

Question: what does 2^A represent?

Notice:

$$\Delta[X] \subset [0, 1]^X.$$

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

so that

$$\begin{aligned} B^n &= \{f \mid f : \{0, \dots, n-1\} \rightarrow B\} \\ &= \{f \mid f : n \rightarrow B\}. \end{aligned}$$

We can now generalise to

$$B^A =_{Def} \{f \mid f : A \rightarrow B\}.$$

Question: what does 2^A represent?

Notice:

$$\Delta[X] \subset [0, 1]^X.$$

Later, we will use the following identities:

$$\blacksquare (Z^X)^Y \sim Z^{X \times Y}$$

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

so that

$$\begin{aligned} B^n &= \{f \mid f : \{0, \dots, n-1\} \rightarrow B\} \\ &= \{f \mid f : n \rightarrow B\}. \end{aligned}$$

We can now generalise to

$$B^A =_{Def} \{f \mid f : A \rightarrow B\}.$$

Question: what does 2^A represent?

Notice:

$$\Delta[X] \subset [0, 1]^X.$$

Later, we will use the following identities:

■ $(Z^X)^Y \sim Z^{X \times Y}$

■ $\prod_{i=1}^n (Y_i)^X \sim (\prod_{i=1}^n Y_i)^X$

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

so that

$$\begin{aligned} B^n &= \{f \mid f : \{0, \dots, n-1\} \rightarrow B\} \\ &= \{f \mid f : n \rightarrow B\}. \end{aligned}$$

We can now generalise to

$$B^A =_{Def} \{f \mid f : A \rightarrow B\}.$$

Question: what does 2^A represent?

Notice:

$$\Delta[X] \subset [0, 1]^X.$$

Later, we will use the following identities:

$$\blacksquare (Z^X)^Y \sim Z^{X \times Y}$$

$$\blacksquare \prod_{i=1}^n (Y_i)^X \sim (\prod_{i=1}^n Y_i)^X$$

Other examples (which we won't use):

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

so that

$$\begin{aligned} B^n &= \{f \mid f : \{0, \dots, n-1\} \rightarrow B\} \\ &= \{f \mid f : n \rightarrow B\}. \end{aligned}$$

We can now generalise to

$$B^A =_{Def} \{f \mid f : A \rightarrow B\}.$$

Question: what does 2^A represent?

Notice:

$$\Delta[X] \subset [0, 1]^X.$$

Later, we will use the following identities:

$$\blacksquare (Z^X)^Y \sim Z^{X \times Y}$$

$$\blacksquare \prod_{i=1}^n (Y_i)^X \sim (\prod_{i=1}^n Y_i)^X$$

Other examples (which we won't use):

$$\blacksquare (Z^X)^Y \sim (Z^Y)^X$$

Notation: $B^A = \{f \mid f : A \rightarrow B\}$

so that

$$\begin{aligned} B^n &= \{f \mid f : \{0, \dots, n-1\} \rightarrow B\} \\ &= \{f \mid f : n \rightarrow B\}. \end{aligned}$$

We can now generalise to

$$B^A =_{Def} \{f \mid f : A \rightarrow B\}.$$

Question: what does 2^A represent?

Notice:

$$\Delta[X] \subset [0, 1]^X.$$

Later, we will use the following identities:

- $(Z^X)^Y \sim Z^{X \times Y}$
- $\prod_{i=1}^n (Y_i)^X \sim (\prod_{i=1}^n Y_i)^X$

Other examples (which we won't use):

- $(Z^X)^Y \sim (Z^Y)^X$
- $X^Y \times X^Z \sim X^{(Y \cup Z)}$, provided Y and Z are disjoint

Kuhn's result

Suppose i believes that j will play strategy g_r^j with probability λ_r , $r \leq L$.

Kuhn's result

Suppose i believes that j will play strategy g_r^j with probability λ_r , $r \leq L$.

Kuhn's equivalent behavioural strategy g^j will choose the action x after history h with probability

$$g^j(x|h) =_{Def} \sum_{r=1}^L (\lambda_r | h) g_r^j(x|h)$$

The factor $\lambda_r | h$ represents the posterior probability of choosing g_r^j given h :

$$\lambda_r | h = \frac{\lambda_r \hat{g}_r^j(h)}{\sum_{w=1}^L \lambda_w \hat{g}_w^j(h)}.$$

where $\hat{g}_r^j(h)$ represents the probability of h being reached when all players other than j take the actions leading to h with probability one, and player j mixes according to g_r^j .

Kuhn's result

Suppose i believes that j will play strategy g_r^j with probability λ_r , $r \leq L$.

Kuhn's equivalent behavioural strategy g^j will choose the action x after history h with probability

$$g^j(x|h) =_{Def} \sum_{r=1}^L (\lambda_r | h) g_r^j(x|h)$$

The factor $\lambda_r | h$ represents the posterior probability of choosing g_r^j given h :

$$\lambda_r | h = \frac{\lambda_r \hat{g}_r^j(h)}{\sum_{w=1}^L \lambda_w \hat{g}_w^j(h)}.$$

where $\hat{g}_r^j(h)$ represents the probability of h being reached when all players other than j take the actions leading to h with probability one, and player j mixes according to g_r^j .

Playing against the strategies g_r^j with the probabilities λ_r and playing against an equivalently constructed behaviour strategy g^j generate identical probability distributions on future play paths.

(Kuhn, 1953; Aumann, 1964.)

Proof of main theorem (Kalai and Lehrer, 1993)

Especially the first part of the proof is rather dense. It uses two results from probability theory.

Proof of main theorem (Kalai and Lehrer, 1993)

Especially the first part of the proof is rather dense. It uses two results from probability theory.

1. The theorem on Radon-Nikodym (mentioned above), which (roughly!) says that if $\nu \ll \mu$, then ν can be expressed in terms of μ .

Proof of main theorem (Kalai and Lehrer, 1993)

Especially the first part of the proof is rather dense. It uses two results from probability theory.

1. The theorem on Radon-Nikodym (mentioned above), which (roughly!) says that if $\nu \ll \mu$, then ν can be expressed in terms of μ .
2. Lévy's theorem, which (very roughly!) is about continuity of expectation through a so-called *filter*.²

Theorem. (Lévy's Convergence Theorem). Let (Ω, \mathcal{F}, P) be a probability space, and let $\{\mathcal{F}_n\}_n$ be a non-decreasing family of σ -algebras contained in \mathcal{F} . Let \mathcal{F}_∞ be the smallest σ -algebra around $\cup\{\mathcal{F}_n\}_n$. Let X be a random variable with finite expectation. Then, both P -a.s.³ and in the L_1 -sense,⁴

$$E[X|\mathcal{F}_n] \rightarrow E[X|\mathcal{F}_\infty] \text{ as } n \rightarrow \infty.$$

²Sequence of monotone non-decreasing σ -algebras.

³ $P\{|\dots - \dots| > \epsilon\} \rightarrow 0$.

⁴ $E|\dots - \dots| \rightarrow 0$.

The Radon-Nikodym theorem

Given X and σ -finite probability measures μ and ν on X such that $\nu \ll \mu$, then there is a measurable function $f : X \rightarrow [0, \infty)$, such that for all μ -measurable sets E ,

$$\nu\{E\} = \int_E f d\mu.$$

Remarks:

The Radon-Nikodym theorem

Given X and σ -finite probability measures μ and ν on X such that $\nu \ll \mu$, then there is a measurable function $f : X \rightarrow [0, \infty)$, such that for all μ -measurable sets E ,

$$\nu\{E\} = \int_E f d\mu.$$

Remarks:

- The theorem is named after Johann Radon, who proved the theorem for the special case where the underlying space is R^n in 1913, and for Otton Nikodym who proved the general case in 1930.

The Radon-Nikodym theorem

Given X and σ -finite probability measures μ and ν on X such that $\nu \ll \mu$, then there is a measurable function $f : X \rightarrow [0, \infty)$, such that for all μ -measurable sets E ,

$$\nu\{E\} = \int_E f d\mu.$$

Remarks:

- The theorem is named after Johann Radon, who proved the theorem for the special case where the underlying space is R^n in 1913, and for Otton Nikodym who proved the general case in 1930.
- The theorem tells if and how it is possible to change from one probability measure to another.

The Radon-Nikodym theorem

Given X and σ -finite probability measures μ and ν on X such that $\nu \ll \mu$, then there is a measurable function $f : X \rightarrow [0, \infty)$, such that for all μ -measurable sets E ,

$$\nu\{E\} = \int_E f d\mu.$$

Remarks:

- The theorem is named after Johann Radon, who proved the theorem for the special case where the underlying space is R^n in 1913, and for Otton Nikodym who proved the general case in 1930.
- The theorem tells if and how it is possible to change from one probability measure to another.
- Specifically, the probability density function of a random variable is the Radon-Nikodym derivative of the induced measure with respect to some base measure (usually the Lebesgue measure for continuous random variables).

Lévy's Theorem

Let (Ω, \mathcal{F}, P) be a probability space, and let $\{\mathcal{F}_n\}_n$ be a non-decreasing family of σ -algebras contained in \mathcal{F} . Let $\mathcal{F}_\infty = \sigma(\cup\{\mathcal{F}_n\}_n)$. Let X be a random variable with finite expectation. Then, both P -a.s. and in the L_1 -sense,

$$E[X|\mathcal{F}_n] \rightarrow E[X|\mathcal{F}_\infty] \text{ as } n \rightarrow \infty.$$

Lévy's Theorem

For a sequence of random variables $\{X_n\}_{n=1}^{\infty}$ where

Lévy's Theorem

For a sequence of random variables $\{X_n\}_{n=1}^{\infty}$ where

1. $X_n \xrightarrow{a.s.} X$

Lévy's Theorem

For a sequence of random variables $\{X_n\}_{n=1}^{\infty}$ where

1. $X_n \xrightarrow{a.s.} X$
2. $|X_n| < Y$ for some random variable Y with $E[Y] < \infty$

it follows that

Lévy's Theorem

For a sequence of random variables $\{X_n\}_{n=1}^{\infty}$ where

1. $X_n \xrightarrow{a.s.} X$
2. $|X_n| < Y$ for some random variable Y with $E[Y] < \infty$

it follows that

■ $E[|X|] < \infty,$

Lévy's Theorem

For a sequence of random variables $\{X_n\}_{n=1}^{\infty}$ where

1. $X_n \xrightarrow{a.s.} X$
2. $|X_n| < Y$ for some random variable Y with $E[Y] < \infty$

it follows that

- $E[|X|] < \infty,$
- $E[X_n] \rightarrow E[X],$

Lévy's Theorem

For a sequence of random variables $\{X_n\}_{n=1}^{\infty}$ where

1. $X_n \xrightarrow{a.s.} X$
2. $|X_n| < Y$ for some random variable Y with $E[Y] < \infty$

it follows that

- $E[|X|] < \infty$,
- $E[X_n] \rightarrow E[X]$,
- $E[|X - X_n|] \rightarrow 0$.

Essentially, it is a sufficient condition for the almost sure convergence to imply L1-convergence. The condition $|X_n| < Y$, $EY < \infty$ could be relaxed. Instead, the sequence $\{X_n\}_{n=1}^{\infty}$ should be uniformly integrable.

Lévy's Theorem

For a sequence of random variables $\{X_n\}_{n=1}^{\infty}$ where

1. $X_n \xrightarrow{a.s.} X$
2. $|X_n| < Y$ for some random variable Y with $E[Y] < \infty$

it follows that

- $E[|X|] < \infty$,
- $E[X_n] \rightarrow E[X]$,
- $E[|X - X_n|] \rightarrow 0$.

Essentially, it is a sufficient condition for the almost sure convergence to imply L1-convergence. The condition $|X_n| < Y$, $EY < \infty$ could be relaxed. Instead, the sequence $\{X_n\}_{n=1}^{\infty}$ should be uniformly integrable.

The theorem is simply a special case of Lebesgue's dominated convergence theorem in measure theory.

Part VI: An Impossibility Result

Suppose players respond rationally

Definition. Given a predictive learning rule (f_i, g_i) , rule g_i is said to be **rational** given f_i if, for each $h \in H$, rule g_i maximises expected discounted payoffs over all continuations of h .

Suppose players respond rationally

Definition. Given a predictive learning rule (f_i, g_i) , rule g_i is said to be **rational** given f_i if, for each $h \in H$, rule g_i maximises expected discounted payoffs over all continuations of h .

- This is called **sequential rationality**: players optimise after each round.

Suppose players respond rationally

Definition. Given a predictive learning rule (f_i, g_i) , rule g_i is said to be **rational** given f_i if, for each $h \in H$, rule g_i maximises expected discounted payoffs over all continuations of h .

- This is called **sequential rationality**: players optimise after each round.
- Sequential rationality very much fits **CKR**.

Suppose players respond rationally

Definition. Given a predictive learning rule (f_i, g_i) , rule g_i is said to be **rational** given f_i if, for each $h \in H$, rule g_i maximises expected discounted payoffs over all continuations of h .

- This is called **sequential rationality**: players optimise after each round.
- Sequential rationality very much fits **CKR**.

Assumption. (f_i, g_i) does not depend on u_{-i} .

Suppose players respond rationally

Definition. Given a predictive learning rule (f_i, g_i) , rule g_i is said to be **rational** given f_i if, for each $h \in H$, rule g_i maximises expected discounted payoffs over all continuations of h .

- This is called **sequential rationality**: players optimise after each round.
- Sequential rationality very much fits **CKR**.

Assumption. (f_i, g_i) does not depend on u_{-i} .

- Player i 's forecast, f_i , may not depend on its own payoffs u_i .

If payoff realisations are independent across players, this is reasonable to assume, because payoffs do not convey information about the behaviour of opponents.

Suppose players respond rationally

Definition. Given a predictive learning rule (f_i, g_i) , rule g_i is said to be **rational** given f_i if, for each $h \in H$, rule g_i maximises expected discounted payoffs over all continuations of h .

- This is called **sequential rationality**: players optimise after each round.
- Sequential rationality very much fits **CKR**.

Assumption. (f_i, g_i) does not depend on u_{-i} .

- Player i 's forecast, f_i , may not depend on its own payoffs u_i .

If payoff realisations are independent across players, this is reasonable to assume, because payoffs do not convey information about the behaviour of opponents.

- Also, g_i cannot depend on u_{-i} because i 's payoffs do not give information on the payoffs of opponents.

Subjective ϵ -equilibrium

Definition. A vector of n profiles

$$(v_i, g_i) = \begin{pmatrix} g_1 & v_1^2 & v_1^3 & \dots & v_1^n \\ v_2^1 & g_2 & v_2^3 & \dots & v_2^n \\ v_3^1 & v_3^2 & g_3 & \dots & v_3^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ v_n^1 & v_n^2 & v_n^3 & \dots & g_n \end{pmatrix}$$

is said to be a **subjective**
 ϵ -equilibrium
if

Subjective ϵ -equilibrium

Definition. A vector of n profiles

$$(v_i, g_i) = \begin{pmatrix} g_1 & v_1^2 & v_1^3 & \dots & v_1^n \\ v_2^1 & g_2 & v_2^3 & \dots & v_2^n \\ v_3^1 & v_3^2 & g_3 & \dots & v_3^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ v_n^1 & v_n^2 & v_n^3 & \dots & g_n \end{pmatrix}$$

is said to be a **subjective**
 ϵ -equilibrium
if

1. All players are rational.

Subjective ϵ -equilibrium

Definition. A vector of n profiles

$$(v_i, g_i) = \begin{pmatrix} g_1 & v_1^2 & v_1^3 & \dots & v_1^n \\ v_2^1 & g_2 & v_2^3 & \dots & v_2^n \\ v_3^1 & v_3^2 & g_3 & \dots & v_3^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ v_n^1 & v_n^2 & v_n^3 & \dots & g_n \end{pmatrix}$$

is said to be a **subjective ϵ -equilibrium** if

1. All players are rational.
2. Everyone ϵ -plays like the diagonal.

Corollary. Assume the same conditions as before. Then, for every $\epsilon > 0$, and for almost every realisation of play ω (w.r.t. μ), there is a round T such that, for every $t \geq T$,

$$\{ \mu_i(\cdot \mid \omega|_t)_i \}_i$$

forms a subjective ϵ -equilibrium.

Subjective ϵ -equilibrium

Definition. A vector of n profiles

$$(v_i, g_i) = \begin{pmatrix} g_1 & v_1^2 & v_1^3 & \dots & v_1^n \\ v_2^1 & g_2 & v_2^3 & \dots & v_2^n \\ v_3^1 & v_3^2 & g_3 & \dots & v_3^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ v_n^1 & v_n^2 & v_n^3 & \dots & g_n \end{pmatrix}$$

is said to be a **subjective ϵ -equilibrium** if

1. All players are rational.
2. Everyone ϵ -plays like the diagonal.

Corollary. Assume the same conditions as before. Then, for every $\epsilon > 0$, and for almost every realisation of play ω (w.r.t. μ), there is a round T such that, for every $t \geq T$,

$$\{ \mu_i(\cdot \mid \omega|_t)_i \}_i$$

forms a subjective ϵ -equilibrium.

Proof. Immediate! □

Subjective ϵ -equilibrium

Definition. A vector of n profiles

$$(v_i, g_i) = \begin{pmatrix} g_1 & v_1^2 & v_1^3 & \dots & v_1^n \\ v_2^1 & g_2 & v_2^3 & \dots & v_2^n \\ v_3^1 & v_3^2 & g_3 & \dots & v_3^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ v_n^1 & v_n^2 & v_n^3 & \dots & g_n \end{pmatrix}$$

is said to be a **subjective ϵ -equilibrium** if

1. All players are rational.
2. Everyone ϵ -plays like the diagonal.

Corollary. Assume the same conditions as before. Then, for every $\epsilon > 0$, and for almost every realisation of play ω (w.r.t. μ), there is a round T such that, for every $t \geq T$,

$$\{ \mu_i(\cdot \mid \omega|_t)_i \}_i$$

forms a subjective ϵ -equilibrium.

Proof. Immediate! □

There's no catch: recall that players converge even if they do not play best replies but play, e.g., maxmin.

Impossibility Theorem (Foster and Young, 2001)

Theorem. *Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.*

New concepts:

Impossibility Theorem (Foster and Young, 2001)

Theorem. *Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.*

New concepts:

- An uncertain, almost-zero-sum, two person game. Multiple actions are not excluded, but the discussion assumes $|X_i| = 2$, for $i = 1, 2$.

Impossibility Theorem (Foster and Young, 2001)

Theorem. Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.

New concepts:

- An uncertain, almost-zero-sum, two person game. Multiple actions are not excluded, but the discussion assumes $|X_i| = 2$, for $i = 1, 2$.
- The range of uncertainty λ of such a game.

Impossibility Theorem (Foster and Young, 2001)

Theorem. *Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.*

New concepts:

- An uncertain, almost-zero-sum, two person game. Multiple actions are not excluded, but the discussion assumes $|X_i| = 2$, for $i = 1, 2$.
- The range of uncertainty λ of such a game.
- A player that is perfectly rational given its predictions.

Impossibility Theorem (Foster and Young, 2001)

Theorem. *Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.*

New concepts:

- An uncertain, almost-zero-sum, two person game. Multiple actions are not excluded, but the discussion assumes $|X_i| = 2$, for $i = 1, 2$.
- The range of uncertainty λ of such a game.
- A player that is perfectly rational given its predictions.
- Being a good predictor.

Impossibility Theorem (Foster and Young, 2001)

Theorem. *Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.*

New concepts:

- An uncertain, almost-zero-sum, two person game. Multiple actions are not excluded, but the discussion assumes $|X_i| = 2$, for $i = 1, 2$.
- The range of uncertainty λ of such a game.
- A player that is perfectly rational given its predictions.
- Being a good predictor.
- To have a realisation of payoffs (rather than a realisation of play).

Impossibility Theorem (Foster and Young, 2001)

Theorem. *Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.*

New concepts:

- An **uncertain, almost-zero-sum, two person game**. Multiple actions are not excluded, but the discussion assumes $|X_i| = 2$, for $i = 1, 2$.
- The **range of uncertainty λ** of such a game.
- A player that is **perfectly rational given its predictions**.
- Being a **good predictor**.
- To have a **realisation of payoffs** (rather than a realisation of play).
- Behaviour that is **asymptotically far from Nash equilibria**.

Uncertain almost-zero-sum games

Uncertain almost-zero-sum games

1. Let G be a n -player zero-sum game with payoffs $u_i : X \rightarrow R$, such that all Nash equilibria are **mixed**.

Uncertain almost-zero-sum games

1. Let G be a n -player zero-sum game with payoffs $u_i : X \rightarrow R$, such that all Nash equilibria are **mixed**.
2. Let $\lambda > 0$ be small and let $\epsilon \in \Delta[-\lambda/2, \lambda/2]$. Notice: this time, ϵ is a random variable.

Uncertain almost-zero-sum games

1. Let G be a n -player zero-sum game with payoffs $u_i : X \rightarrow R$, such that all Nash equilibria are **mixed**.
2. Let $\lambda > 0$ be small and let $\epsilon \in \Delta[-\lambda/2, \lambda/2]$. Notice: this time, ϵ is a random variable.
3. Give each player an ϵ_i , based on ϵ and i.i.d.

Uncertain almost-zero-sum games

1. Let G be a n -player zero-sum game with payoffs $u_i : X \rightarrow R$, such that all Nash equilibria are **mixed**.
2. Let $\lambda > 0$ be small and let $\epsilon \in \Delta[-\lambda/2, \lambda/2]$. Notice: this time, ϵ is a random variable.
3. Give each player an ϵ_i , based on ϵ and i.i.d.
4. Create a new game G' with payoffs $U_i(x) =_{Def} u_i(x) + \epsilon_i(x)$.

Uncertain almost-zero-sum games

1. Let G be a n -player zero-sum game with payoffs $u_i : X \rightarrow R$, such that all Nash equilibria are **mixed**.
 2. Let $\lambda > 0$ be small and let $\epsilon \in \Delta[-\lambda/2, \lambda/2]$. Notice: this time, ϵ is a random variable.
 3. Give each player an ϵ_i , based on ϵ and i.i.d.
 4. Create a new game G' with payoffs $U_i(x) =_{Def} u_i(x) + \epsilon_i(x)$.
- From here, G' 's matrix is fixed, and G' is ready to be played.

Example. “Uncertain matching pennies”. Property: when λ is sufficiently small, G' still possesses a unique (mixed) Nash equilibrium.

$$M = \begin{array}{cc} & \begin{array}{cc} \text{H} & \text{T} \end{array} \\ \begin{array}{c} \text{H} \\ \text{T} \end{array} & \left(\begin{array}{cc} 1 + \epsilon_{11}, -1 + \epsilon'_{11} & -1 + \epsilon_{12}, 1 + \epsilon'_{12} \\ -1 + \epsilon_{21}, 1 + \epsilon'_{21} & 1 + \epsilon_{22}, -1 + \epsilon'_{22} \end{array} \right) \end{array}$$

Behaviour that is asymptotically far from Nash equilibria

Behaviour that is asymptotically far from Nash equilibria

- Let $\omega \in \Omega$ be a certain realisation of play. Let $\epsilon > 0$.

Definition. Game play on ω is said to be ϵ -far from all equilibria if in at least $(1 - \epsilon)\%$ of the rounds, play at round t is at least ϵ away from every Nash equilibrium of the repeated subgame starting at t .

Behaviour that is asymptotically far from Nash equilibria

- Let $\omega \in \Omega$ be a certain realisation of play. Let $\epsilon > 0$.

Definition. Game play on ω is said to be ϵ -far from all equilibria if in at least $(1 - \epsilon)\%$ of the rounds, play at round t is at least ϵ away from every Nash equilibrium of the repeated subgame starting at t .

- Let $\mu \in \Delta[\Omega]$.

Definition. Behaviour is said to asymptotically far from equilibrium behaviour if there exists an $\epsilon > 0$ such that play is ϵ -far from all equilibria with μ -probability one.

Behaviour that is asymptotically far from Nash equilibria

- Let $\omega \in \Omega$ be a certain realisation of play. Let $\epsilon > 0$.

Definition. Game play on ω is said to be ϵ -far from all equilibria if in at least $(1 - \epsilon)\%$ of the rounds, play at round t is at least ϵ away from every Nash equilibrium of the repeated subgame starting at t .

- Let $\mu \in \Delta[\Omega]$.

Definition. Behaviour is said to asymptotically far from equilibrium behaviour if there exists an $\epsilon > 0$ such that play is ϵ -far from all equilibria with μ -probability one.

Remarks:

Behaviour that is asymptotically far from Nash equilibria

- Let $\omega \in \Omega$ be a certain realisation of play. Let $\epsilon > 0$.

Definition. Game play on ω is said to be ϵ -far from all equilibria if in at least $(1 - \epsilon)\%$ of the rounds, play at round t is at least ϵ away from every Nash equilibrium of the repeated subgame starting at t .

- Let $\mu \in \Delta[\Omega]$.

Definition. Behaviour is said to asymptotically far from equilibrium behaviour if there exists an $\epsilon > 0$ such that play is ϵ -far from all equilibria with μ -probability one.

Remarks:

- Intuition: mostly, play does not look like equilibrium play in a repeated subgame.

Behaviour that is asymptotically far from Nash equilibria

- Let $\omega \in \Omega$ be a certain realisation of play. Let $\epsilon > 0$.

Definition. Game play on ω is said to be ϵ -far from all equilibria if in at least $(1 - \epsilon)\%$ of the rounds, play at round t is at least ϵ away from every Nash equilibrium of the repeated subgame starting at t .

- Let $\mu \in \Delta[\Omega]$.

Definition. Behaviour is said to asymptotically far from equilibrium behaviour if there exists an $\epsilon > 0$ such that play is ϵ -far from all equilibria with μ -probability one.

Remarks:

- Intuition: mostly, play does not look like equilibrium play in a repeated subgame.
- This is far more stronger than the negation of asymptotic closeness, in the sense of Kalai and Lehrer.

Impossibility Theorem (Foster and Young, 2001)

Theorem. Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.

Impossibility Theorem (Foster and Young, 2001)

Theorem. *Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.*

- Assumptions within this theorem are consistent with assumptions within Kalai *et al.*'s main theorem.

Impossibility Theorem (Foster and Young, 2001)

Theorem. *Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.*

- Assumptions within this theorem are consistent with assumptions within Kalai *et al.*'s main theorem.
- Still, the two theorems seem to contradict each other.

Impossibility Theorem (Foster and Young, 2001)

Theorem. *Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.*

- Assumptions within this theorem are consistent with assumptions within Kalai *et al.*'s main theorem.
- Still, the two theorems seem to contradict each other.
- Explanation:

Impossibility Theorem (Foster and Young, 2001)

Theorem. *Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.*

- Assumptions within this theorem are consistent with assumptions within Kalai *et al.*'s main theorem.
- Still, the two theorems seem to contradict each other.
- Explanation:
 - Kalai *et al.*'s main theorem states that players will (almost) correctly predict the **on-path** portions of the other players' strategies.

Impossibility Theorem (Foster and Young, 2001)

Theorem. Let G be an uncertain, almost-zero-sum, two person game, all of whose Nash equilibria are fully mixed. Assume that both players use predictive learning rules and are perfectly rational given their predictions. If the range of uncertainty λ is sufficiently small, then for almost all realisations of the payoffs, one or both players are not good predictors and their behaviours are asymptotically far from Nash equilibria.

- Assumptions within this theorem are consistent with assumptions within Kalai *et al.*'s main theorem.
- Still, the two theorems seem to contradict each other.
- Explanation:
 - Kalai *et al.*'s main theorem states that players will (almost) correctly predict the **on-path** portions of the other players' strategies.
 - Kalai *et al.*'s main theorem does **not** state that players will (almost) learn the true strategies of their opponents.