

# Multi-agent learning

## Evolutionary game theory

*Gerard Vreeswijk*, Intelligent Systems Group, Computer Science  
Department, Faculty of Sciences, Utrecht University, The  
Netherlands.

Wednesday 15<sup>th</sup> May, 2019

# Plan for today



# Plan for today



1. Symmetric games in normal form.

# Plan for today



1. Symmetric games in normal form.
2. Evolutionary game theory.

# Plan for today



1. Symmetric games in normal form.
2. Evolutionary game theory.
  - **Static analysis:** evolutionarily stable strategies.



# Plan for today



1. Symmetric games in normal form.
2. Evolutionary game theory.
  - **Static analysis:** evolutionarily stable strategies.
  - **Dynamic analysis:** the replicator equation.

# Plan for today



1. Symmetric games in normal form.
2. Evolutionary game theory.
  - **Static analysis:** evolutionarily stable strategies.
  - **Dynamic analysis:** the replicator equation.

\* H. Peters (2008): *Game Theory: A Multi-Leveled Approach*. Springer, ISBN: 978-3-540-69290-4. Ch. 8: An Introduction to Evolutionary Games.

# Plan for today



1. Symmetric games in normal form.
2. Evolutionary game theory.
  - **Static analysis:** evolutionarily stable strategies.
  - **Dynamic analysis:** the replicator equation.

\* H. Peters (2008): *Game Theory: A Multi-Leveled Approach*. Springer, ISBN: 978-3-540-69290-4. Ch. 8: An Introduction to Evolutionary Games.

\* Shoham *et al.* (2009): *Multi-agent Systems*. Ch. 7: "Learning and Teaching".



# Plan for today



1. Symmetric games in normal form.
2. Evolutionary game theory.
  - **Static analysis:** evolutionarily stable strategies.
  - **Dynamic analysis:** the replicator equation.

\* H. Peters (2008): *Game Theory: A Multi-Leveled Approach*. Springer, ISBN: 978-3-540-69290-4. Ch. 8: An Introduction to Evolutionary Games.

\* Shoham *et al.* (2009): *Multi-agent Systems*. Ch. 7: “Learning and Teaching”.

\* Jörgen W. Weibull (1997): *Evolutionary Game Theory*, MIT Press.

# Symmetric games in normal form

# Ways to denote the payoff matrix of a symmetric game

Symmetric game: **only your actions matter, not your role.**

# Ways to denote the payoff matrix of a symmetric game

Symmetric game: **only your actions matter, not your role.** For 2-player games: it is not important whether you are row or column player:

$$\begin{array}{c} A_1 \\ A_2 \\ A_3 \end{array} \begin{pmatrix} A_1 & A_2 & A_3 \\ 1, 1 & 2, 3 & -4, 6 \\ 3, 2 & 0, 0 & 8, -7 \\ 6, -4 & -7, 8 & 5, 5 \end{pmatrix}$$



# Ways to denote the payoff matrix of a symmetric game

Symmetric game: **only your actions matter, not your role**. For 2-player games: it is not important whether you are row or column player:

$$\begin{array}{c} A_1 \\ A_2 \\ A_3 \end{array} \begin{array}{cc} A_1 & A_2 & A_3 \end{array} \begin{pmatrix} 1,1 & 2,3 & -4,6 \\ 3,2 & 0,0 & 8,-7 \\ 6,-4 & -7,8 & 5,5 \end{pmatrix} = \begin{array}{c} A_1 \\ A_2 \\ A_3 \end{array} \begin{array}{cc} A_1 & A_2 & A_3 \end{array} \begin{pmatrix} 1,1 & 2,3 & -4,6 \\ & 0,0 & 8,-7 \\ & & 5,5 \end{pmatrix}$$

■ As a bi-matrix.

# Ways to denote the payoff matrix of a symmetric game

Symmetric game: **only your actions matter, not your role**. For 2-player games: it is not important whether you are row or column player:

$$\begin{array}{c}
 \begin{array}{c} A_1 \quad A_2 \quad A_3 \\
 A_1 \left( \begin{array}{ccc} 1,1 & 2,3 & -4,6 \\
 A_2 \left( \begin{array}{ccc} 3,2 & 0,0 & 8,-7 \\
 A_3 \left( \begin{array}{ccc} 6,-4 & -7,8 & 5,5 \end{array} \right) \end{array} \right) = \begin{array}{c}
 \begin{array}{c} A_1 \quad A_2 \quad A_3 \\
 A_1 \left( \begin{array}{ccc} 1,1 & 2,3 & -4,6 \\
 A_2 \left( \begin{array}{ccc} & 0,0 & 8,-7 \\
 A_3 \left( \begin{array}{ccc} & & 5,5 \end{array} \right) \end{array} \right) \end{array} \right) \\
 \\
 \sim \begin{array}{c}
 \begin{array}{c} A_1 \quad A_2 \quad A_3 \\
 A_1 \left( \begin{array}{ccc} 1 & 2 & -4 \\
 A_2 \left( \begin{array}{ccc} 3 & 0 & 8 \\
 A_3 \left( \begin{array}{ccc} 6 & -7 & 5 \end{array} \right) \end{array} \right) \end{array} \right) .
 \end{array}
 \end{array}
 \end{array}$$

- As a bi-matrix.
- As a partially filled bi-matrix.

# Ways to denote the payoff matrix of a symmetric game

Symmetric game: **only your actions matter, not your role.** For 2-player games: it is not important whether you are row or column player:

$$\begin{array}{c} A_1 \quad A_2 \quad A_3 \\ A_1 \left( \begin{array}{ccc} 1,1 & 2,3 & -4,6 \\ 3,2 & 0,0 & 8,-7 \\ 6,-4 & -7,8 & 5,5 \end{array} \right) \\ A_2 \\ A_3 \end{array} = \begin{array}{c} A_1 \quad A_2 \quad A_3 \\ A_1 \left( \begin{array}{ccc} 1,1 & 2,3 & -4,6 \\ & 0,0 & 8,-7 \\ & & 5,5 \end{array} \right) \\ A_2 \\ A_3 \end{array} \sim \begin{array}{c} A_1 \quad A_2 \quad A_3 \\ A_1 \left( \begin{array}{ccc} 1 & 2 & -4 \\ 3 & 0 & 8 \\ 6 & -7 & 5 \end{array} \right) \\ A_2 \\ A_3 \end{array}.$$

- As a bi-matrix.
- As a partially filled bi-matrix.
- As a plain matrix.

# Symmetric normal-form games

**Example.** Hawk-dove game (share  $V$  or threaten [possibly fight:  $-C$ ]):

	H	D
H	$(V - C)/2$	$V$
D	$0$	$V/2$



# Symmetric normal-form games

**Example.** Hawk-dove game (share  $V$  or threaten [possibly fight:  $-C$ ]):

	H	D
H	$(V - C)/2$	$V$
D	$0$	$V/2$

	H	D
H	$-2, -2$	$2, 0$
D	$0, 2$	$1, 1$

$V=2, C=6$

# Symmetric normal-form games

**Example.** Hawk-dove game (share  $V$  or threaten [possibly fight:  $-C$ ]):

	H	D
H	$(V - C)/2$	$V$
D	$0$	$V/2$

	H	D
H	$-2, -2$	$2, 0$
D	$0, 2$	$1, 1$

 $V=2, C=6$ 

More examples: prisoner's dilemma, chicken (= hawk-dove), matching pennies, stag hunt.

# Symmetric normal-form games

**Example.** Hawk-dove game (share  $V$  or threaten [possibly fight:  $-C$ ]):

	H	D
H	$(V - C)/2$	$V$
D	$0$	$V/2$

	H	D
H	$-2, -2$	$2, 0$
D	$0, 2$	$1, 1$

 $V=2, C=6$ 

More examples: prisoner's dilemma, chicken (= hawk-dove), matching pennies, stag hunt.

**Definition.** A game is **symmetric** when players have equal actions and payoffs:

$$u_i(a_1, \dots, a_i, \dots, a_j, \dots, a_n) = u_j(a_1, \dots, a_j, \dots, a_i, \dots, a_n).$$

for all  $i$  and  $j$ .

# Symmetric normal-form games

**Example.** Hawk-dove game (share  $V$  or threaten [possibly fight:  $-C$ ]):

	H	D
H	$(V - C)/2$	$V$
D	$0$	$V/2$

	H	D
H	$-2, -2$	$2, 0$
D	$0, 2$	$1, 1$

$V=2, C=6$

More examples: prisoner's dilemma, chicken (= hawk-dove), matching pennies, stag hunt.

**Definition.** A game is **symmetric** when players have equal actions and payoffs:

$$u_i(a_1, \dots, a_i, \dots, a_j, \dots, a_n) = u_j(a_1, \dots, a_j, \dots, a_i, \dots, a_n).$$

for all  $i$  and  $j$ .

So a 2-player game  $G = (A, B)$  is symmetric iff  $m = n$  and  $B = A^T$ .



# Symmetric equilibrium

# Symmetric equilibrium

**Definition.** Let  $p$  be a strategy in an  $n$ -player symmetric game. If the  $n$ -vector  $(p, \dots, p)$  is a NE,  $p$  is called a **symmetric equilibrium**.

# Symmetric equilibrium

**Definition.** Let  $p$  be a strategy in an  $n$ -player symmetric game. If the  $n$ -vector  $(p, \dots, p)$  is a NE,  $p$  is called a **symmetric equilibrium**.

■ **!! Symmetric equilibria can be identified with strategies !!**

# Symmetric equilibrium

**Definition.** Let  $p$  be a strategy in an  $n$ -player symmetric game. If the  $n$ -vector  $(p, \dots, p)$  is a NE,  $p$  is called a **symmetric equilibrium**.

- **!! Symmetric equilibria can be identified with strategies !!**
- (Theorem.) Every symmetric game has at least one symmetric equilibrium.



# Symmetric equilibrium

**Definition.** Let  $p$  be a strategy in an  $n$ -player symmetric game. If the  $n$ -vector  $(p, \dots, p)$  is a NE,  $p$  is called a **symmetric equilibrium**.

- **!! Symmetric equilibria can be identified with strategies !!**
- (Theorem.) Every symmetric game has at least one symmetric equilibrium.
- (Fact.) Symmetric games can have a-symmetric equilibria.

# Symmetric equilibrium

**Definition.** Let  $p$  be a strategy in an  $n$ -player symmetric game. If the  $n$ -vector  $(p, \dots, p)$  is a NE,  $p$  is called a **symmetric equilibrium**.

- !! Symmetric equilibria can be identified with strategies !!
- (Theorem.) Every symmetric game has at least one symmetric equilibrium.
- (Fact.) Symmetric games can have a-symmetric equilibria.

For example Hawk-Dove:

	H	D
H	$-2, -2$	$2, 0$
D	$0, 2$	$1, 1$

# Symmetric equilibrium

**Definition.** Let  $p$  be a strategy in an  $n$ -player symmetric game. If the  $n$ -vector  $(p, \dots, p)$  is a NE,  $p$  is called a **symmetric equilibrium**.

- !! Symmetric equilibria can be identified with strategies !!
- (Theorem.) Every symmetric game has at least one symmetric equilibrium.
- (Fact.) Symmetric games can have asymmetric equilibria.

For example Hawk-Dove:

	H	D
H	$-2, -2$	$2, 0$
D	$0, 2$	$1, 1$

Two asymmetric equilibria and one symmetric equilibrium  $(1/3, 1/3)$ .

# Hawk vs. Dove



# Evolutionary game theory

# Evolutionary game theory: the idea



# Evolutionary game theory: the idea

There are  $n$ , say 5, species.



# Evolutionary game theory: the idea

There are  $n$ , say 5, species. An encounter between individuals of different species yields payoffs for both. For row:

$$A = \begin{matrix} & s_1 & s_2 & s_3 & s_4 & s_5 \\ \begin{matrix} s_1 \\ s_2 \\ s_3 \\ s_4 \\ s_5 \end{matrix} & \begin{pmatrix} 6 & 7 & 0 & -1 & 0 \\ -1 & 5 & -1 & 4 & 7 \\ 9 & 0 & 8 & 9 & 6 \\ 0 & -4 & -2 & 3 & -3 \\ 3 & 0 & 6 & 0 & -1 \end{pmatrix} \end{matrix}.$$

# Evolutionary game theory: the idea

There are  $n$ , say 5, species. An encounter between individuals of different species yields payoffs for both. For row:

$$A = \begin{matrix} & \begin{matrix} s_1 & s_2 & s_3 & s_4 & s_5 \end{matrix} \\ \begin{matrix} s_1 \\ s_2 \\ s_3 \\ s_4 \\ s_5 \end{matrix} & \begin{pmatrix} 6 & 7 & 0 & -1 & 0 \\ -1 & 5 & -1 & 4 & 7 \\ 9 & 0 & 8 & 9 & 6 \\ 0 & -4 & -2 & 3 & -3 \\ 3 & 0 & 6 & 0 & -1 \end{pmatrix} \end{matrix}.$$

- The population consists of a very large number of individuals, each playing a pure strategy. Individuals interact randomly.

# Evolutionary game theory: the idea

There are  $n$ , say 5, species. An encounter between individuals of different species yields payoffs for both. For row:

$$A = \begin{matrix} & \begin{matrix} s_1 & s_2 & s_3 & s_4 & s_5 \end{matrix} \\ \begin{matrix} s_1 \\ s_2 \\ s_3 \\ s_4 \\ s_5 \end{matrix} & \begin{pmatrix} 6 & 7 & 0 & -1 & 0 \\ -1 & 5 & -1 & 4 & 7 \\ 9 & 0 & 8 & 9 & 6 \\ 0 & -4 & -2 & 3 & -3 \\ 3 & 0 & 6 & 0 & -1 \end{pmatrix} \end{matrix}.$$

- The population consists of a very large number of individuals, each playing a pure strategy. Individuals interact randomly.
- We are interested in **proportions**:  $p = (p_1, \dots, p_5)^T \in \Delta^5$ .

# Evolutionary game theory: the idea

There are  $n$ , say 5, species. An encounter between individuals of different species yields payoffs for both. For row:

$$A = \begin{matrix} & \begin{matrix} s_1 & s_2 & s_3 & s_4 & s_5 \end{matrix} \\ \begin{matrix} s_1 \\ s_2 \\ s_3 \\ s_4 \\ s_5 \end{matrix} & \begin{pmatrix} 6 & 7 & 0 & -1 & 0 \\ -1 & 5 & -1 & 4 & 7 \\ 9 & 0 & 8 & 9 & 6 \\ 0 & -4 & -2 & 3 & -3 \\ 3 & 0 & 6 & 0 & -1 \end{pmatrix} \end{matrix}.$$

- The population consists of a very large number of individuals, each playing a pure strategy. Individuals interact randomly.
- We are interested in **proportions**:  $p = (p_1, \dots, p_5)^T \in \Delta^5$ .
- The **fitness** of an individual of species  $i$  is:  $f_i = \sum_{j=1}^5 p_j A_{ij} = (Ap)_i$ .

# Evolutionary game theory: the idea

There are  $n$ , say 5, species. An encounter between individuals of different species yields payoffs for both. For row:

$$A = \begin{matrix} & \begin{matrix} s_1 & s_2 & s_3 & s_4 & s_5 \end{matrix} \\ \begin{matrix} s_1 \\ s_2 \\ s_3 \\ s_4 \\ s_5 \end{matrix} & \begin{pmatrix} 6 & 7 & 0 & -1 & 0 \\ -1 & 5 & -1 & 4 & 7 \\ 9 & 0 & 8 & 9 & 6 \\ 0 & -4 & -2 & 3 & -3 \\ 3 & 0 & 6 & 0 & -1 \end{pmatrix} \end{matrix}.$$

- The population consists of a very large number of individuals, each playing a pure strategy. Individuals interact randomly.
- We are interested in **proportions**:  $p = (p_1, \dots, p_5)^T \in \Delta^5$ .
- The **fitness** of an individual of species  $i$  is:  $f_i = \sum_{j=1}^5 p_j A_{ij} = (Ap)_i$ .
- The **average fitness** is:  $\bar{f} = \sum_{i=1}^5 p_i f_i = p^T Ap$ .

# Evolutionarily stable strategies (ESSs)

# Alternative population $q$

We can think of

$$q^T A p = (q_1, \dots, q_m) \begin{pmatrix} a_{11} & \dots & a_{1m} \\ \vdots & \ddots & \vdots \\ a_{m1} & \dots & a_{mm} \end{pmatrix} \begin{pmatrix} p_1 \\ \vdots \\ p_m \end{pmatrix} = \sum_{i=1}^m q_i f_i$$

as the average fitness of a population in state  $q$  playing the game against members of a population in state  $p$ .



# Alternative population $q$

We can think of

$$q^T A p = (q_1, \dots, q_m) \begin{pmatrix} a_{11} & \dots & a_{1m} \\ \vdots & \ddots & \vdots \\ a_{m1} & \dots & a_{mm} \end{pmatrix} \begin{pmatrix} p_1 \\ \vdots \\ p_m \end{pmatrix} = \sum_{i=1}^m q_i f_i$$

as the average fitness of a population in state  $q$  playing the game against members of a population in state  $p$ .

- $p^T A q > q^T A q$  means that state  $p$  performs better against state  $q$  than  $q$  does against itself.

# Alternative population $q$

We can think of

$$q^T A p = (q_1, \dots, q_m) \begin{pmatrix} a_{11} & \dots & a_{1m} \\ \vdots & \ddots & \vdots \\ a_{m1} & \dots & a_{mm} \end{pmatrix} \begin{pmatrix} p_1 \\ \vdots \\ p_m \end{pmatrix} = \sum_{i=1}^m q_i f_i$$

as the average fitness of a population in state  $q$  playing the game against members of a population in state  $p$ .

- $p^T A q > q^T A q$  means that state  $p$  performs better against state  $q$  than  $q$  does against itself.
- $p^T A p > q^T A p$  means that state  $p$  performs better against itself than state  $q$  does against  $p$ .

# Alternative population $q$

We can think of

$$q^T A p = (q_1, \dots, q_m) \begin{pmatrix} a_{11} & \dots & a_{1m} \\ \vdots & \ddots & \vdots \\ a_{m1} & \dots & a_{mm} \end{pmatrix} \begin{pmatrix} p_1 \\ \vdots \\ p_m \end{pmatrix} = \sum_{i=1}^m q_i f_i$$

as the average fitness of a population in state  $q$  playing the game against members of a population in state  $p$ .

- $p^T A q > q^T A q$  means that state  $p$  performs better against state  $q$  than  $q$  does against itself.
- $p^T A p > q^T A p$  means that state  $p$  performs better against itself than state  $q$  does against  $p$ .
- ...
- ...

# Evolutionarily stable strategy

# Evolutionarily stable strategy

■ Idea:

# Evolutionarily stable strategy

- Idea: equilibrium  $\Rightarrow$  all species present (some may be absent!) are equally fit.

# Evolutionarily stable strategy

- Idea: equilibrium  $\Rightarrow$  all species present (some may be absent!) are equally fit.
- Additional idea:



# Evolutionarily stable strategy

- Idea: equilibrium  $\Rightarrow$  all species present (some may be absent!) are equally fit.
- Additional idea:  
  
The equilibrium  $p$  must be **stable**, that is, it must be able to resist perturbations (= small random shocks).

# Evolutionarily stable strategy

- Idea: equilibrium  $\Rightarrow$  all species present (some may be absent!) are equally fit.

- Additional idea:

The equilibrium  $p$  must be **stable**, that is, it must be able to resist perturbations (= small random shocks).

A **perturbation** of  $p$  would be  $(1 - \epsilon)p + \epsilon q$ , for some  $q$  and small  $\epsilon > 0$ .

# Evolutionarily stable strategy

- Idea: equilibrium  $\Rightarrow$  all species present (some may be absent!) are equally fit.

- Additional idea:

The equilibrium  $p$  must be **stable**, that is, it must be able to resist perturbations (= small random shocks).

A **perturbation** of  $p$  would be  $(1 - \epsilon)p + \epsilon q$ , for some  $q$  and small  $\epsilon > 0$ . Also:  $q$  **invades**  $p$ .

# Evolutionarily stable strategy

- Idea: equilibrium  $\Rightarrow$  all species present (some may be absent!) are equally fit.

- Additional idea:

The equilibrium  $p$  must be **stable**, that is, it must be able to resist perturbations (= small random shocks).

A **perturbation** of  $p$  would be  $(1 - \epsilon)p + \epsilon q$ , for some  $q$  and small  $\epsilon > 0$ . Also:  $q$  **invades**  $p$ .

**Definition.** Let  $A$  be a symmetric game. A state  $p \in \Delta^m$  is **evolutionarily stable** if for every  $q \in \Delta^m$ ,  $q \neq p$ , there is an  $0 < \epsilon_q < 1$  such that for all  $\epsilon < \epsilon_q$ :

$$p^T A[(1 - \epsilon)p + \epsilon q] > q^T A[(1 - \epsilon)p + \epsilon q].$$

# A lemma

# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .



# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.*  $(2 \Leftrightarrow 3)$  is proven through logical manipulation

# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.*  $(2 \Leftrightarrow 3)$  is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.

# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:


1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.* (2 $\Leftrightarrow$ 3) is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.



# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:


1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.* (2 $\Leftrightarrow$ 3) is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.



# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:


1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.* (2 $\Leftrightarrow$ 3) is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.



# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:


1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.* (2 $\Leftrightarrow$ 3) is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.



# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:


1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.* (2 $\Leftrightarrow$ 3) is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.





# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.*  $(2 \Leftrightarrow 3)$  is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.

Now for  $(1 \Leftrightarrow 2)$ .

# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.*  $(2 \Leftrightarrow 3)$  is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.

Now for  $(1 \Leftrightarrow 2)$ .

Evidently:

$$\lim_{\epsilon \rightarrow 0} (1 - \epsilon)A + \epsilon B = A.$$

# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.*  $(2 \Leftrightarrow 3)$  is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.

Now for  $(1 \Leftrightarrow 2)$ .

Evidently:

$$\lim_{\epsilon \rightarrow 0} (1 - \epsilon)A + \epsilon B = A.$$

■ If  $A > 0$ , then both (1) and (2) are true.

# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.*  $(2 \Leftrightarrow 3)$  is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.

Now for  $(1 \Leftrightarrow 2)$ .

Evidently:

$$\lim_{\epsilon \rightarrow 0} (1 - \epsilon)A + \epsilon B = A.$$

- If  $A > 0$ , then both (1) and (2) are true.
- If  $A < 0$ , then both (1) and (2) are false.

# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.*  $(2 \Leftrightarrow 3)$  is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.

Now for  $(1 \Leftrightarrow 2)$ .

Evidently:

$$\lim_{\epsilon \rightarrow 0} (1 - \epsilon)A + \epsilon B = A.$$

■ If  $A > 0$ , then both (1) and (2) are true.

■ If  $A < 0$ , then both (1) and (2) are false.

If  $A = 0$  then  $(1 - \epsilon)A + \epsilon B$  reduces to  $\epsilon B$ .

# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.*  $(2 \Leftrightarrow 3)$  is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.

Now for  $(1 \Leftrightarrow 2)$ .

Evidently:

$$\lim_{\epsilon \rightarrow 0} (1 - \epsilon)A + \epsilon B = A.$$

■ If  $A > 0$ , then both (1) and (2) are true.

■ If  $A < 0$ , then both (1) and (2) are false.

If  $A = 0$  then  $(1 - \epsilon)A + \epsilon B$  reduces to  $\epsilon B$ .

■ If  $B > 0$ , then both (1) and (2) are true.

# A lemma

**Lemma.** Let  $A$  and  $B$  be real numbers. The following statements are equivalent:

1. For small  $\epsilon > 0$ :

$$(1 - \epsilon)A + \epsilon B > 0.$$

2.  $A > 0$ , or  $A = 0$  and  $B > 0$ .

3.  $A \geq 0$ , and  $A = 0 \Rightarrow B > 0$ .

*Proof.*  $(2 \Leftrightarrow 3)$  is proven through logical manipulation, or by shading areas in an  $A$ - $B$  axis system.

Now for  $(1 \Leftrightarrow 2)$ .

Evidently:

$$\lim_{\epsilon \rightarrow 0} (1 - \epsilon)A + \epsilon B = A.$$

■ If  $A > 0$ , then both (1) and (2) are true.

■ If  $A < 0$ , then both (1) and (2) are false.

If  $A = 0$  then  $(1 - \epsilon)A + \epsilon B$  reduces to  $\epsilon B$ .

■ If  $B > 0$ , then both (1) and (2) are true.

■ If  $B \leq 0$ , then both (1) and (2) are false. □

# Alternative formulations of an ESS

**Theorem.** The following statements are equivalent:



# Alternative formulations of an ESS

**Theorem.** The following statements are equivalent:

1.  $p$  is an ESS

# Alternative formulations of an ESS

**Theorem.** The following statements are equivalent:

1.  $p$  is an ESS
2. for all  $q$ :  $p^T A p \geq q^T A p$

# Alternative formulations of an ESS

**Theorem.** The following statements are equivalent:

1.  $p$  is an ESS
2. for all  $q$ :  $p^T A p \geq q^T A p$ , and

# Alternative formulations of an ESS

**Theorem.** The following statements are equivalent:

1.  $p$  is an ESS
2. for all  $q$ :  $p^T A p \geq q^T A p$ , and if  $q^T A p = p^T A p$  then  $p^T A q > q^T A q$ .

# Alternative formulations of an ESS

**Theorem.** The following statements are equivalent:

1.  $p$  is an ESS
2. for all  $q$ :  $p^T A p \geq q^T A p$ , and if  $q^T A p = p^T A p$  then  $p^T A q > q^T A q$ .
3. for all  $q$ :  $p^T A p > q^T A p$

# Alternative formulations of an ESS

**Theorem.** The following statements are equivalent:

1.  $p$  is an ESS
2. for all  $q$ :  $p^T A p \geq q^T A p$ , and if  $q^T A p = p^T A p$  then  $p^T A q > q^T A q$ .
3. for all  $q$ :  $p^T A p > q^T A p$ , or

# Alternative formulations of an ESS

**Theorem.** The following statements are equivalent:

1.  $p$  is an ESS
2. for all  $q$ :  $p^T A p \geq q^T A p$ , and if  $q^T A p = p^T A p$  then  $p^T A q > q^T A q$ .
3. for all  $q$ :  $p^T A p > q^T A p$ , or  $q^T A p = p^T A p$  and  $p^T A q > q^T A q$ .

# Alternative formulations of an ESS

**Theorem.** The following statements are equivalent:

1.  $p$  is an ESS
2. for all  $q$ :  $p^T A p \geq q^T A p$ , and if  $q^T A p = p^T A p$  then  $p^T A q > q^T A q$ .
3. for all  $q$ :  $p^T A p > q^T A p$ , or  $q^T A p = p^T A p$  and  $p^T A q > q^T A q$ .

*Proof.* Suppose for every  $q \in \Delta^m$ ,  $q \neq p$ , there is an  $0 < \epsilon_q < 1$  such that for all  $\epsilon < \epsilon_q$ :

$$q^T A[(1 - \epsilon)p + \epsilon q] < p^T A[(1 - \epsilon)p + \epsilon q].$$



# Alternative formulations of an ESS

**Theorem.** The following statements are equivalent:

1.  $p$  is an ESS
2. for all  $q$ :  $p^T A p \geq q^T A p$ , and if  $q^T A p = p^T A p$  then  $p^T A q > q^T A q$ .
3. for all  $q$ :  $p^T A p > q^T A p$ , or  $q^T A p = p^T A p$  and  $p^T A q > q^T A q$ .

*Proof.* Suppose for every  $q \in \Delta^m$ ,  $q \neq p$ , there is an  $0 < \epsilon_q < 1$  such that for all  $\epsilon < \epsilon_q$ :

$$q^T A[(1 - \epsilon)p + \epsilon q] < p^T A[(1 - \epsilon)p + \epsilon q].$$

Rewriting with linear algebra yields:

$$(1 - \epsilon)(p^T A p - q^T A p) + \epsilon(p^T A q - q^T A q) > 0$$

# Alternative formulations of an ESS

**Theorem.** The following statements are equivalent:

1.  $p$  is an ESS
2. for all  $q$ :  $p^T A p \geq q^T A p$ , and if  $q^T A p = p^T A p$  then  $p^T A q > q^T A q$ .
3. for all  $q$ :  $p^T A p > q^T A p$ , or  $q^T A p = p^T A p$  and  $p^T A q > q^T A q$ .

*Proof.* Suppose for every  $q \in \Delta^m$ ,  $q \neq p$ , there is an  $0 < \epsilon_q < 1$  such that for all  $\epsilon < \epsilon_q$ :

$$q^T A[(1 - \epsilon)p + \epsilon q] < p^T A[(1 - \epsilon)p + \epsilon q].$$

Rewriting with linear algebra yields:

$$(1 - \epsilon)(p^T A p - q^T A p) + \epsilon(p^T A q - q^T A q) > 0$$

Now use the previous lemma:

For all  $A$  and  $B$ :  $(1 - \epsilon)A + \epsilon B > 0$ , for small  $\epsilon$

$$\Leftrightarrow A > 0, \text{ or } A = 0 \text{ and } B > 0 \Leftrightarrow A \geq 0, \text{ and } A = 0 \Rightarrow B > 0. \square$$

# Neutrally Stable Strategy (NSS)

# Neutrally Stable Strategy (NSS)

**Definition.** Let  $A$  be a symmetric game. A state  $p \in \Delta^m$  is **neutrally stable** if for every  $q \in \Delta^m$ ,  $q \neq p$ , there is an  $0 < \epsilon_q < 1$  such that for all  $\epsilon < \epsilon_q$ :

$$q^T A[(1 - \epsilon)p + \epsilon q] \leq p^T A[(1 - \epsilon)p + \epsilon q].$$

# Neutrally Stable Strategy (NSS)

**Definition.** Let  $A$  be a symmetric game. A state  $p \in \Delta^m$  is **neutrally stable** if for every  $q \in \Delta^m, q \neq p$ , there is an  $0 < \epsilon_q < 1$  such that for all  $\epsilon < \epsilon_q$ :

$$q^T A[(1 - \epsilon)p + \epsilon q] \leq p^T A[(1 - \epsilon)p + \epsilon q].$$

- ESS: other populations  $q \neq p$  are driven out.

# Neutrally Stable Strategy (NSS)

**Definition.** Let  $A$  be a symmetric game. A state  $p \in \Delta^m$  is **neutrally stable** if for every  $q \in \Delta^m$ ,  $q \neq p$ , there is an  $0 < \epsilon_q < 1$  such that for all  $\epsilon < \epsilon_q$ :

$$q^T A[(1 - \epsilon)p + \epsilon q] \leq p^T A[(1 - \epsilon)p + \epsilon q].$$

- ESS: other populations  $q \neq p$  are driven out.
- NSS: other populations  $q \neq p$  do not expand.

# Neutrally Stable Strategy (NSS)

**Definition.** Let  $A$  be a symmetric game. A state  $p \in \Delta^m$  is **neutrally stable** if for every  $q \in \Delta^m, q \neq p$ , there is an  $0 < \epsilon_q < 1$  such that for all  $\epsilon < \epsilon_q$ :

$$q^T A[(1 - \epsilon)p + \epsilon q] \leq p^T A[(1 - \epsilon)p + \epsilon q].$$

- ESS: other populations  $q \neq p$  are driven out.
- NSS: other populations  $q \neq p$  do not expand.

**Theorem.** The following statements are equivalent:

# Neutrally Stable Strategy (NSS)

**Definition.** Let  $A$  be a symmetric game. A state  $p \in \Delta^m$  is **neutrally stable** if for every  $q \in \Delta^m$ ,  $q \neq p$ , there is an  $0 < \epsilon_q < 1$  such that for all  $\epsilon < \epsilon_q$ :

$$q^T A[(1 - \epsilon)p + \epsilon q] \leq p^T A[(1 - \epsilon)p + \epsilon q].$$

- ESS: other populations  $q \neq p$  are driven out.
- NSS: other populations  $q \neq p$  do not expand.

**Theorem.** The following statements are equivalent:

1.  $p$  is an NSS.



# Neutrally Stable Strategy (NSS)

**Definition.** Let  $A$  be a symmetric game. A state  $p \in \Delta^m$  is **neutrally stable** if for every  $q \in \Delta^m, q \neq p$ , there is an  $0 < \epsilon_q < 1$  such that for all  $\epsilon < \epsilon_q$ :

$$q^T A[(1 - \epsilon)p + \epsilon q] \leq p^T A[(1 - \epsilon)p + \epsilon q].$$

- ESS: other populations  $q \neq p$  are driven out.
- NSS: other populations  $q \neq p$  do not expand.

**Theorem.** The following statements are equivalent:

1.  $p$  is an NSS.
2. for all  $q$ :  $p^T A p \geq q^T A p$ , and if  $q^T A p = p^T A p$  then  $p^T A q \geq q^T A q$ .

# Neutrally Stable Strategy (NSS)

**Definition.** Let  $A$  be a symmetric game. A state  $p \in \Delta^m$  is **neutrally stable** if for every  $q \in \Delta^m$ ,  $q \neq p$ , there is an  $0 < \epsilon_q < 1$  such that for all  $\epsilon < \epsilon_q$ :

$$q^T A[(1 - \epsilon)p + \epsilon q] \leq p^T A[(1 - \epsilon)p + \epsilon q].$$

- ESS: other populations  $q \neq p$  are driven out.
- NSS: other populations  $q \neq p$  do not expand.

**Theorem.** The following statements are equivalent:

1.  $p$  is an NSS.
2. for all  $q$ :  $p^T A p \geq q^T A p$ , and if  $q^T A p = p^T A p$  then  $p^T A q \geq q^T A q$ .
3. for all  $q$ :  $p^T A p > q^T A p$ , or  $q^T A p = p^T A p$  and  $p^T A q \geq q^T A q$ .

# Finding ESS

**Example.** Hawk-dove game:

	H	D
H	0,0	3,1
D	1,3	2,2

# Finding ESS

**Example.** Hawk-dove game:

	H	D
H	0,0	3,1
D	1,3	2,2

- Two pure a-symmetric NE, and one mixed symmetric NE:  
 $p = (1/2, 1/2)$ .

# Finding ESS

**Example.** Hawk-dove game:

	H	D
H	0,0	3,1
D	1,3	2,2

- Two pure a-symmetric NE, and one mixed symmetric NE:  
 $p = (1/2, 1/2)$ .
- Since  $p$  is fully mixed, every response  $q$  is a best response to  $p$ . So for all  $q$ ,  $q^T A p = p^T A p$ .

# Finding ESS

**Example.** Hawk-dove game:

	H	D
H	0,0	3,1
D	1,3	2,2

- Two pure a-symmetric NE, and one mixed symmetric NE:  
 $p = (1/2, 1/2)$ .
- Since  $p$  is fully mixed, every response  $q$  is a best response to  $p$ . So for all  $q$ ,  $q^T A p = p^T A p$ .
- We'll have to verify the second condition of an ESS:  
for all  $q \neq p$ :  $q^T A q < p^T A q$ .

# Finding ESS

**Example.** Hawk-dove game:

	H	D
H	0,0	3,1
D	1,3	2,2

- Two pure a-symmetric NE, and one mixed symmetric NE:  
 $p = (1/2, 1/2)$ .
- Since  $p$  is fully mixed, every response  $q$  is a best response to  $p$ . So for all  $q$ ,  $q^T A p = p^T A p$ .
- We'll have to verify the second condition of an ESS:  
for all  $q \neq p$ :  $q^T A q < p^T A q$ . Let  $q = (y, 1 - y)$ ,  $y \neq 1/2$ .

# Finding ESS

**Example.** Hawk-dove game:

	H	D
H	0,0	3,1
D	1,3	2,2

- Two pure a-symmetric NE, and one mixed symmetric NE:  
 $p = (1/2, 1/2)$ .
- Since  $p$  is fully mixed, every response  $q$  is a best response to  $p$ . So for all  $q$ ,  $q^T A p = p^T A p$ .
- We'll have to verify the second condition of an ESS:  
for all  $q \neq p$ :  $q^T A q < p^T A q$ . Let  $q = (y, 1 - y)$ ,  $y \neq 1/2$ . Then


$$p^T A q = \begin{pmatrix} 1/2 & 1/2 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} y \\ 1 - y \end{pmatrix} = -2y + 2\frac{1}{2},$$

$$q^T A q = \begin{pmatrix} y & 1 - y \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} y \\ 1 - y \end{pmatrix} = -2y^2 + 2.$$



# To verify that $p^T A q > q^T A q$


Equivalent  
statements:



# To verify that $p^T Aq > q^T Aq$

Equivalent  
statements:


- $p^T Aq > q^T Aq,$   
 $q \neq p$



# To verify that $p^T Aq > q^T Aq$

Equivalent  
statements:


- $p^T Aq > q^T Aq,$   
 $q \neq p$
- $-2y + 5/2 -$   
 $(-2y^2 + 2) > 0,$   
 $y \neq 1/2$



# To verify that $p^T Aq > q^T Aq$

Equivalent  
statements:


- $p^T Aq > q^T Aq,$   
 $q \neq p$
- $-2y + 5/2 -$   
 $(-2y^2 + 2) > 0,$   
 $y \neq 1/2$
- $2y^2 - 2y + 1/2 >$   
 $0,$   
 $y \neq 1/2.$



# To verify that $p^T Aq > q^T Aq$

Equivalent  
statements:

- $p^T Aq > q^T Aq,$   
 $q \neq p$
- $-2y + 5/2 -$   
 $(-2y^2 + 2) > 0,$   
 $y \neq 1/2$
- $2y^2 - 2y + 1/2 >$   
 $0,$   
 $y \neq 1/2.$
- $(1 - 2y)^2 / 2 > 0,$   
 $y \neq 1/2.$



# ESSs compared to NE

# ESSs compared to NE

## Nash equilibrium

# ESSs compared to NE

## Nash equilibrium

- Economically motivated:  
homo economicus.



# ESSs compared to NE

## Nash equilibrium

- Economically motivated:  
homo economicus.
- Assumes CKR of all players.

# ESSs compared to NE

## Nash equilibrium

- Economically motivated:  
homo economicus.
- Assumes CKR of all players.
- Traditional solution concept.

# ESSs compared to NE

## Nash equilibrium

- Economically motivated:  
homo economicus.
- Assumes CKR of all players.
- Traditional solution concept.

## Evolutionarily stable strategy

# ESSs compared to NE

## Nash equilibrium

- Economically motivated:  
homo economicus.
- Assumes CKR of all players.
- Traditional solution concept.

## Evolutionarily stable strategy

- Biologically motivated.

# ESSs compared to NE

## Nash equilibrium

- Economically motivated:  
homo economicus.
- Assumes CKR of all players.
- Traditional solution concept.

## Evolutionarily stable strategy

- Biologically motivated.
- Individuals have no control over their strategy.

# ESSs compared to NE

## Nash equilibrium

- Economically motivated:  
*homo economicus*.
- Assumes **CKR** of all players.
- Traditional solution concept.

## Evolutionarily stable strategy

- Biologically motivated.
- Individuals have no control over their strategy.
- Individuals need not be aware of the game.

# ESSs compared to NE

## Nash equilibrium

- Economically motivated:  
*homo economicus*.
- Assumes **CKR** of all players.
- Traditional solution concept.

## Evolutionarily stable strategy

- Biologically motivated.
- Individuals have no control over their strategy.
- Individuals need not be aware of the game.

- Individuals reproduce according to relative success (biological fitness).

# ESSs compared to NE

## Nash equilibrium

- Economically motivated:  
*homo economicus*.
- Assumes **CKR** of all players.
- Traditional solution concept.

## Evolutionarily stable strategy

- Biologically motivated.
- Individuals have no control over their strategy.
- Individuals need not be aware of the game.

- Individuals reproduce according to relative success (biological fitness).
- Alternatives occasionally occur (“mutation” / “invasion”).



# ESSs compared to NE

## Nash equilibrium

- Economically motivated:  
*homo economicus*.
- Assumes **CKR** of all players.
- Traditional solution concept.

## Evolutionarily stable strategy

- Biologically motivated.
- Individuals have no control over their strategy.
- Individuals need not be aware of the game.

- Individuals reproduce according to relative success (biological fitness).
- Alternatives occasionally occur (“mutation” / “invasion”).

To be an ESS, a strategy must be resistant to these alternatives.

# ESSs compared to NE

## Nash equilibrium

- Economically motivated:  
*homo economicus*.
- Assumes **CKR** of all players.
- Traditional solution concept.

## Evolutionarily stable strategy

- Biologically motivated.
- Individuals have no control over their strategy.
- Individuals need not be aware of the game.

- Individuals reproduce according to relative success (biological fitness).
- Alternatives occasionally occur (“mutation” / “invasion”).

To be an ESS, a strategy must be resistant to these alternatives.

*“Given the radically different motivating assumptions, it may come as a surprise that ESSes and Nash equilibria often coincide.”*

# ESSs compared to NE

## Nash equilibrium

- Economically motivated:  
*homo economicus*.
- Assumes **CKR** of all players.
- Traditional solution concept.

## Evolutionarily stable strategy

- Biologically motivated.
- Individuals have no control over their strategy.
- Individuals need not be aware of the game.

- Individuals reproduce according to relative success (biological fitness).
- Alternatives occasionally occur (“mutation” / “invasion”).

To be an ESS, a strategy must be resistant to these alternatives.

*“Given the radically different motivating assumptions, it may come as a surprise that ESSes and Nash equilibria often coincide.”*  
(Wikipedia “Evolutionarily stable strategy”, May 2019.)

# ESSs compared to NE

For every symmetric matrix game:

# ESSs compared to NE

For every symmetric matrix game:

Strict Nash equilibrium (NE<sub><</sub>)  
 $\Rightarrow_1$  Evolutionarily stable strategy (ESS)

# ESSs compared to NE

For every symmetric matrix game:

- Strict Nash equilibrium (NE $\subset$ )
- $\Rightarrow_1$  Evolutionarily stable strategy (ESS)
- $\Rightarrow_2$  Neutrally Stable Strategy (NSS)

# ESSs compared to NE

For every symmetric matrix game:

- Strict Nash equilibrium (NE<sub><</sub>)
- $\Rightarrow_1$  Evolutionarily stable strategy (ESS)
- $\Rightarrow_2$  Neutrally Stable Strategy (NSS)
- $\Rightarrow_3$  Nash equilibrium (NE).

# ESSs compared to NE

For every symmetric matrix game:

- Strict Nash equilibrium ( $NE<$ )
- $\Rightarrow_1$  Evolutionarily stable strategy (ESS)
- $\Rightarrow_2$  Neutrally Stable Strategy (NSS)
- $\Rightarrow_3$  Nash equilibrium (NE).

So:

$$\Delta^{NE<} \subseteq \Delta^{ESS} \subseteq \Delta^{NSS} \subseteq \Delta^{NE}.$$



# ESSs compared to NE

For every symmetric matrix game:

- Strict Nash equilibrium ( $NE_{<}$ )
- $\Rightarrow_1$  Evolutionarily stable strategy (ESS)
- $\Rightarrow_2$  Neutrally Stable Strategy (NSS)
- $\Rightarrow_3$  Nash equilibrium (NE).

So:

$$\Delta^{NE_{<}} \subseteq \Delta^{ESS} \subseteq \Delta^{NSS} \subseteq \Delta^{NE}.$$

*Proof.*

# ESSs compared to NE

For every symmetric matrix game:

- Strict Nash equilibrium ( $NE<$ )
- $\Rightarrow_1$  Evolutionarily stable strategy (ESS)
- $\Rightarrow_2$  Neutrally Stable Strategy (NSS)
- $\Rightarrow_3$  Nash equilibrium (NE).

So:

$$\Delta^{NE<} \subseteq \Delta^{ESS} \subseteq \Delta^{NSS} \subseteq \Delta^{NE}.$$

*Proof.*

$\Rightarrow_1$  from 3rd alternative formulation ESS.

# ESSs compared to NE

For every symmetric matrix game:

- Strict Nash equilibrium ( $NE_{<}$ )
- $\Rightarrow_1$  Evolutionarily stable strategy (ESS)
- $\Rightarrow_2$  Neutrally Stable Strategy (NSS)
- $\Rightarrow_3$  Nash equilibrium (NE).

So:

$$\Delta^{NE_{<}} \subseteq \Delta^{ESS} \subseteq \Delta^{NSS} \subseteq \Delta^{NE}.$$

*Proof.*

- $\Rightarrow_1$  from 3rd alternative formulation ESS.
- $\Rightarrow_2$  from  $\epsilon$ -definitions ESS and NSS.

# ESSs compared to NE

For every symmetric matrix game:

- Strict Nash equilibrium ( $NE<$ )
- $\Rightarrow_1$  Evolutionarily stable strategy (ESS)
- $\Rightarrow_2$  Neutrally Stable Strategy (NSS)
- $\Rightarrow_3$  Nash equilibrium (NE).

So:

$$\Delta^{NE<} \subseteq \Delta^{ESS} \subseteq \Delta^{NSS} \subseteq \Delta^{NE}.$$

*Proof.*

- $\Rightarrow_1$  from 3rd alternative formulation ESS.
- $\Rightarrow_2$  from  $\epsilon$ -definitions ESS and NSS.
- $\Rightarrow_3$  from 2nd alternative formulation NSS.

# ESSs compared to NE

For every symmetric matrix game:

- Strict Nash equilibrium ( $NE_{<}$ )
- $\Rightarrow_1$  Evolutionarily stable strategy (ESS)
- $\Rightarrow_2$  Neutrally Stable Strategy (NSS)
- $\Rightarrow_3$  Nash equilibrium (NE).

So:

$$\Delta^{NE_{<}} \subseteq \Delta^{ESS} \subseteq \Delta^{NSS} \subseteq \Delta^{NE}.$$

*Proof.*

- $\Rightarrow_1$  from 3rd alternative formulation ESS.
- $\Rightarrow_2$  from  $\epsilon$ -definitions ESS and NSS.
- $\Rightarrow_3$  from 2nd alternative formulation NSS.

None of the implications can be reversed, i.e., all set inclusions are strict.

# ESSs compared to NE

# ESSs compared to NE

**Claim:**  $\Delta^{\text{NE}<} \subset \Delta^{\text{ESS}} \subset \Delta^{\text{NSS}} \subset \Delta^{\text{NE}}$ , i.e., all inclusions are strict.

# ESSs compared to NE

**Claim:**  $\Delta^{\text{NE}<} \subset \Delta^{\text{ESS}} \subset \Delta^{\text{NSS}} \subset \Delta^{\text{NE}}$ , i.e., all inclusions are strict.

*Proof.*



# ESSs compared to NE

**Claim:**  $\Delta^{\text{NE}^<} \subset \Delta^{\text{ESS}} \subset \Delta^{\text{NSS}} \subset \Delta^{\text{NE}}$ , i.e., all inclusions are strict.

*Proof.* Consider

	A	B
A	$\alpha, \alpha$	$1, 0$
B	$0, 1$	$1, 1$

# ESSs compared to NE

**Claim:**  $\Delta^{\text{NE}<} \subset \Delta^{\text{ESS}} \subset \Delta^{\text{NSS}} \subset \Delta^{\text{NE}}$ , i.e., all inclusions are strict.

*Proof.* Consider

	A	B
A	$\alpha, \alpha$	$1, 0$
B	$0, 1$	$1, 1$

Expected payoff for col:

$$E_{\text{col}}(p, q) = p(q\alpha + (1 - q)) + (1 - p)(q \cdot 0 + (1 - q) \cdot 1) = p(\alpha q - 1) + 1.$$

$$\frac{\partial}{\partial q} E_{\text{col}}(p, q) = \alpha p.$$

# ESSs compared to NE

**Claim:**  $\Delta^{\text{NE}<} \subset \Delta^{\text{ESS}} \subset \Delta^{\text{NSS}} \subset \Delta^{\text{NE}}$ , i.e., all inclusions are strict.

*Proof.* Consider

	A	B
A	$\alpha, \alpha$	$1, 0$
B	$0, 1$	$1, 1$

Expected payoff for col:

$$E_{\text{col}}(p, q) = p(q\alpha + (1 - q)) + (1 - p)(q \cdot 0 + (1 - q) \cdot 1) = p(\alpha q - 1) + 1.$$

$$\frac{\partial}{\partial q} E_{\text{col}}(p, q) = \alpha p.$$

So if  $\alpha p = 0$  then  $(p, p)$  is a symmetric NE.

# ESSs compared to NE

**Claim:**  $\Delta^{\text{NE}<} \subset \Delta^{\text{ESS}} \subset \Delta^{\text{NSS}} \subset \Delta^{\text{NE}}$ , i.e., all inclusions are strict.

*Proof.* Consider

	A	B
A	$\alpha, \alpha$	$1, 0$
B	$0, 1$	$1, 1$

Expected payoff for col:

$$E_{\text{col}}(p, q) = p(q\alpha + (1 - q)) + (1 - p)(q \cdot 0 + (1 - q) \cdot 1) = p(\alpha q - 1) + 1.$$

$$\frac{\partial}{\partial q} E_{\text{col}}(p, q) = \alpha p.$$

So if  $\alpha p = 0$  then  $(p, p)$  is a symmetric NE.

■ For  $\alpha = 0$ , we have symmetric NE for all  $p$ .

# ESSs compared to NE

**Claim:**  $\Delta^{\text{NE}<} \subset \Delta^{\text{ESS}} \subset \Delta^{\text{NSS}} \subset \Delta^{\text{NE}}$ , i.e., all inclusions are strict.

*Proof.* Consider

	A	B
A	$\alpha, \alpha$	$1, 0$
B	$0, 1$	$1, 1$

Expected payoff for col:

$$E_{\text{col}}(p, q) = p(q\alpha + (1 - q)) + (1 - p)(q \cdot 0 + (1 - q) \cdot 1) = p(\alpha q - 1) + 1.$$

$$\frac{\partial}{\partial q} E_{\text{col}}(p, q) = \alpha p.$$

So if  $\alpha p = 0$  then  $(p, p)$  is a symmetric NE.

■ For  $\alpha = 0$ , we have symmetric NE for all  $p$ . None of these are strict.

# ESSs compared to NE

**Claim:**  $\Delta^{\text{NE}<} \subset \Delta^{\text{ESS}} \subset \Delta^{\text{NSS}} \subset \Delta^{\text{NE}}$ , i.e., all inclusions are strict.

*Proof.* Consider

	A	B
A	$\alpha, \alpha$	$1, 0$
B	$0, 1$	$1, 1$

Expected payoff for col:

$$E_{\text{col}}(p, q) = p(q\alpha + (1 - q)) + (1 - p)(q \cdot 0 + (1 - q) \cdot 1) = p(\alpha q - 1) + 1.$$

$$\frac{\partial}{\partial q} E_{\text{col}}(p, q) = \alpha p.$$

So if  $\alpha p = 0$  then  $(p, p)$  is a symmetric NE.

- For  $\alpha = 0$ , we have symmetric NE for all  $p$ . None of these are strict.  
Since  $q^T A p = p^T A p$  for all  $q$

# ESSs compared to NE

**Claim:**  $\Delta^{\text{NE}<} \subset \Delta^{\text{ESS}} \subset \Delta^{\text{NSS}} \subset \Delta^{\text{NE}}$ , i.e., all inclusions are strict.

*Proof.* Consider

	A	B
A	$\alpha, \alpha$	$1, 0$
B	$0, 1$	$1, 1$

Expected payoff for col:

$$E_{\text{col}}(p, q) = p(q\alpha + (1 - q)) + (1 - p)(q \cdot 0 + (1 - q) \cdot 1) = p(\alpha q - 1) + 1.$$

$$\frac{\partial}{\partial q} E_{\text{col}}(p, q) = \alpha p.$$

So if  $\alpha p = 0$  then  $(p, p)$  is a symmetric NE.

- For  $\alpha = 0$ , we have symmetric NE for all  $p$ . None of these are strict. Since  $q^T A p = p^T A p$  for all  $q$ , for ESS it must be that  $p^T A q > q^T A q$  for all  $q$

# ESSs compared to NE

**Claim:**  $\Delta^{\text{NE}<} \subset \Delta^{\text{ESS}} \subset \Delta^{\text{NSS}} \subset \Delta^{\text{NE}}$ , i.e., all inclusions are strict.

*Proof.* Consider

	A	B
A	$\alpha, \alpha$	$1, 0$
B	$0, 1$	$1, 1$

Expected payoff for col:

$$E_{\text{col}}(p, q) = p(q\alpha + (1 - q)) + (1 - p)(q \cdot 0 + (1 - q) \cdot 1) = p(\alpha q - 1) + 1.$$

$$\frac{\partial}{\partial q} E_{\text{col}}(p, q) = \alpha p.$$

So if  $\alpha p = 0$  then  $(p, p)$  is a symmetric NE.

- For  $\alpha = 0$ , we have symmetric NE for all  $p$ . None of these are strict. Since  $q^T A p = p^T A p$  for all  $q$ , for ESS it must be that  $p^T A q > q^T A q$  for all  $q$ , i.e.  $p(1 + \alpha(p - q - 1)) > 0$  for all  $q$ .



# ESSs compared to NE

**Claim:**  $\Delta^{\text{NE}<} \subset \Delta^{\text{ESS}} \subset \Delta^{\text{NSS}} \subset \Delta^{\text{NE}}$ , i.e., all inclusions are strict.

*Proof.* Consider

	A	B
A	$\alpha, \alpha$	$1, 0$
B	$0, 1$	$1, 1$

Expected payoff for col:

$$E_{\text{col}}(p, q) = p(q\alpha + (1 - q)) + (1 - p)(q \cdot 0 + (1 - q) \cdot 1) = p(\alpha q - 1) + 1.$$

$$\frac{\partial}{\partial q} E_{\text{col}}(p, q) = \alpha p.$$

So if  $\alpha p = 0$  then  $(p, p)$  is a symmetric NE.

- For  $\alpha = 0$ , we have symmetric NE for all  $p$ . None of these are strict. Since  $q^T A p = p^T A p$  for all  $q$ , for ESS it must be that  $p^T A q > q^T A q$  for all  $q$ , i.e.  $p(1 + \alpha(p - q - 1)) > 0$  for all  $q$ . For  $p = 1/2$  this is the case.

# ESSs compared to NE

**Claim:**  $\Delta^{\text{NE}<} \subset \Delta^{\text{ESS}} \subset \Delta^{\text{NSS}} \subset \Delta^{\text{NE}}$ , i.e., all inclusions are strict.

*Proof.* Consider

	A	B
A	$\alpha, \alpha$	$1, 0$
B	$0, 1$	$1, 1$

Expected payoff for col:

$$E_{\text{col}}(p, q) = p(q\alpha + (1 - q)) + (1 - p)(q \cdot 0 + (1 - q) \cdot 1) = p(\alpha q - 1) + 1.$$
$$\frac{\partial}{\partial q} E_{\text{col}}(p, q) = \alpha p.$$

So if  $\alpha p = 0$  then  $(p, p)$  is a symmetric NE.

■ For  $\alpha = 0$ , we have symmetric NE for all  $p$ . None of these are strict. Since  $q^T A p = p^T A p$  for all  $q$ , for ESS it must be that  $p^T A q > q^T A q$  for all  $q$ , i.e.  $p(1 + \alpha(p - q - 1)) > 0$  for all  $q$ . For  $p = 1/2$  this is the case.

■ Similarly,  $\alpha \in \{-1, 1\}$  makes the last two inclusions strict.  $\square$

# ESSs may not exist

Consider rock paper scissors:

	R	P	S
R	0,0	-1,1	1,-1
P	1,-1	0,0	-1,1
S	-1,1	1,-1	0,0

# ESSs may not exist

Consider rock paper scissors:    R           P           S

R	0, 0	-1, 1	1, -1
P	1, -1	0, 0	-1, 1
S	-1, 1	1, -1	0, 0

- $p = (1/3, 1/3, 1/3)$  is the only symmetric NE, so  $p$  is the only ESS candidate.

# ESSs may not exist

Consider rock paper scissors:    R           P           S

R	0, 0	-1, 1	1, -1
P	1, -1	0, 0	-1, 1
S	-1, 1	1, -1	0, 0

- $p = (1/3, 1/3, 1/3)$  is the only symmetric NE, so  $p$  is the only ESS candidate.
- Because  $\text{support}(e_1) \subseteq \text{support}(p)$ , the pure strategy  $e_1$  is also a best response. Therefore, the second condition of an ESS must hold for  $p$  as well:

$$e_1 A p = p A p \Rightarrow p A e_1 > e_1 A e_1.$$

# ESSs may not exist

Consider rock paper scissors:

	R	P	S
R	0,0	-1,1	1,-1
P	1,-1	0,0	-1,1
S	-1,1	1,-1	0,0

- $p = (1/3, 1/3, 1/3)$  is the only symmetric NE, so  $p$  is the only ESS candidate.
- Because  $\text{support}(e_1) \subseteq \text{support}(p)$ , the pure strategy  $e_1$  is also a best response. Therefore, the second condition of an ESS must hold for  $p$  as well:

$$e_1 A p = p A p \Rightarrow p A e_1 > e_1 A e_1.$$

But it does not hold:  $p A e_1 = e_1 A e_1 = 0$ .

# ESSs may not exist

Consider rock paper scissors: R P S

	R	P	S
R	0,0	-1,1	1,-1
P	1,-1	0,0	-1,1
S	-1,1	1,-1	0,0

- $p = (1/3, 1/3, 1/3)$  is the only symmetric NE, so  $p$  is the only ESS candidate.
- Because  $\text{support}(e_1) \subseteq \text{support}(p)$ , the pure strategy  $e_1$  is also a best response. Therefore, the second condition of an ESS must hold for  $p$  as well:

$$e_1 A p = p A p \Rightarrow p A e_1 > e_1 A e_1.$$

But it does not hold:  $p A e_1 = e_1 A e_1 = 0$ .

**Question to contemplate:** is it OK that ESSs may not exist?

# How many ESSs?

**Theorem.** If  $p, q \in \Delta^m$  such that  $p$  is an ESS,  $q \neq p$ , and  $\text{support}(q) \subseteq \text{support}(p)$ , then  $q \notin \Delta^{\text{NE}}$ .



# How many ESSs?

**Theorem.** If  $p, q \in \Delta^m$  such that  $p$  is an ESS,  $q \neq p$ , and  $\text{support}(q) \subseteq \text{support}(p)$ , then  $q \notin \Delta^{\text{NE}}$ .

*Proof.*

# How many ESSs?

**Theorem.** If  $p, q \in \Delta^m$  such that  $p$  is an ESS,  $q \neq p$ , and  $\text{support}(q) \subseteq \text{support}(p)$ , then  $q \notin \Delta^{\text{NE}}$ .

*Proof.* Because  $\text{support}(q) \subseteq \text{support}(p)$ , the strategy  $q$  is a best response to  $p$  as well:  $qAp = pAp$ .

# How many ESSs?

**Theorem.** If  $p, q \in \Delta^m$  such that  $p$  is an ESS,  $q \neq p$ , and  $\text{support}(q) \subseteq \text{support}(p)$ , then  $q \notin \Delta^{\text{NE}}$ .

*Proof.* Because  $\text{support}(q) \subseteq \text{support}(p)$ , the strategy  $q$  is a best response to  $p$  as well:  $qAp = pAp$ . This equation triggers the second condition of an ESS:

$$qAp = pAp \Rightarrow pAq > qAq.$$

# How many ESSs?

**Theorem.** If  $p, q \in \Delta^m$  such that  $p$  is an ESS,  $q \neq p$ , and  $\text{support}(q) \subseteq \text{support}(p)$ , then  $q \notin \Delta^{\text{NE}}$ .

*Proof.* Because  $\text{support}(q) \subseteq \text{support}(p)$ , the strategy  $q$  is a best response to  $p$  as well:  $qAp = pAp$ . This equation triggers the second condition of an ESS:

$$qAp = pAp \Rightarrow pAq > qAq.$$

The consequent of the last implication says that  $q$  cannot be a NE. □

# How many ESSs?

**Theorem.** If  $p, q \in \Delta^m$  such that  $p$  is an ESS,  $q \neq p$ , and  $\text{support}(q) \subseteq \text{support}(p)$ , then  $q \notin \Delta^{\text{NE}}$ .

*Proof.* Because  $\text{support}(q) \subseteq \text{support}(p)$ , the strategy  $q$  is a best response to  $p$  as well:  $qAp = pAp$ . This equation triggers the second condition of an ESS:

$$qAp = pAp \Rightarrow pAq > qAq.$$

The consequent of the last implication says that  $q$  cannot be a NE. □

**Corollaries.**

# How many ESSs?

**Theorem.** If  $p, q \in \Delta^m$  such that  $p$  is an ESS,  $q \neq p$ , and  $\text{support}(q) \subseteq \text{support}(p)$ , then  $q \notin \Delta^{\text{NE}}$ .

*Proof.* Because  $\text{support}(q) \subseteq \text{support}(p)$ , the strategy  $q$  is a best response to  $p$  as well:  $qAp = pAp$ . This equation triggers the second condition of an ESS:

$$qAp = pAp \Rightarrow pAq > qAq.$$

The consequent of the last implication says that  $q$  cannot be a NE. □

## Corollaries.

1. The set  $\Delta^{\text{ESS}}$  is finite and possibly empty.

# How many ESSs?

**Theorem.** If  $p, q \in \Delta^m$  such that  $p$  is an ESS,  $q \neq p$ , and  $\text{support}(q) \subseteq \text{support}(p)$ , then  $q \notin \Delta^{\text{NE}}$ .

*Proof.* Because  $\text{support}(q) \subseteq \text{support}(p)$ , the strategy  $q$  is a best response to  $p$  as well:  $qAp = pAp$ . This equation triggers the second condition of an ESS:

$$qAp = pAp \Rightarrow pAq > qAq.$$

The consequent of the last implication says that  $q$  cannot be a NE. □

## Corollaries.

1. The set  $\Delta^{\text{ESS}}$  is finite and possibly empty.
2. If an ESS is fully mixed, it is unique.

# The replicator equation



# Idea of the replicator equation

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0) A p$$

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix}$$

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix}$$

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 3(1 - x).$$

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 3(1 - x).$$

Playing Dove against  $p = (x, 1 - x)$  yields:



# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 3(1 - x).$$

Playing Dove against  $p = (x, 1 - x)$  yields:

$$(0 \ 1)Ap$$

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 3(1 - x).$$

Playing Dove against  $p = (x, 1 - x)$  yields:

$$(0 \ 1)Ap = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix}$$

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 3(1 - x).$$

Playing Dove against  $p = (x, 1 - x)$  yields:

$$(0 \ 1)Ap = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix}$$

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 3(1 - x).$$

Playing Dove against  $p = (x, 1 - x)$  yields:

$$(0 \ 1)Ap = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 2 - x.$$

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 3(1 - x).$$

Playing Dove against  $p = (x, 1 - x)$  yields:

$$(0 \ 1)Ap = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 2 - x.$$

Average fitness:

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 3(1 - x).$$

Playing Dove against  $p = (x, 1 - x)$  yields:

$$(0 \ 1)Ap = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 2 - x.$$

Average fitness:

$$pAp$$

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 3(1 - x).$$

Playing Dove against  $p = (x, 1 - x)$  yields:

$$(0 \ 1)Ap = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 2 - x.$$

Average fitness:

$$pAp = \begin{pmatrix} x & 1 - x \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix}$$

# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 3(1 - x).$$

Playing Dove against  $p = (x, 1 - x)$  yields:

$$(0 \ 1)Ap = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 2 - x.$$

Average fitness:

$$pAp = \begin{pmatrix} x & 1 - x \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} x & 1 - x \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix}$$



# Idea of the replicator equation

Consider Hawk-Dove:

	H	D
H	0,0	3,1
D	1,3	2,2

Playing Hawk against a population  $p = (x, 1 - x)$  yields:

$$(1 \ 0)Ap = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 3(1 - x).$$

Playing Dove against  $p = (x, 1 - x)$  yields:

$$(0 \ 1)Ap = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 2 - x.$$

Average fitness:

$$pAp = \begin{pmatrix} x & 1 - x \end{pmatrix} \begin{pmatrix} 0 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ 1 - x \end{pmatrix} = \begin{pmatrix} x & 1 - x \end{pmatrix} \begin{pmatrix} 3(1 - x) \\ x + 2(1 - x) \end{pmatrix} = 2 - 2x.$$

# Idea of the replicator equation

The replicator equation **stipulates** that the change in proportion of a species

# Idea of the replicator equation

The replicator equation **stipulates** that the change in proportion of a species over time

# Idea of the replicator equation

The replicator equation **stipulates** that the change in proportion of a species over time is determined by the difference between its fitness and the average fitness:

# Idea of the replicator equation

The replicator equation **stipulates** that the change in proportion of a species over time is determined by the difference between its fitness and the average fitness:

$$\dot{x} \left( = \frac{\partial x}{\partial t} \right) \stackrel{\text{Def}}{=} x[f_H - \bar{f}].$$

# Idea of the replicator equation

The replicator equation **stipulates** that the change in proportion of a species over time is determined by the difference between its fitness and the average fitness:

$$\dot{x} \left( = \frac{\partial x}{\partial t} \right) \stackrel{\text{Def}}{=} x[f_H - \bar{f}].$$

So in the Hawk-dove scenario:


$$\begin{aligned}\dot{x} &= x[f_H - \bar{f}] \\ &= x[3(1 - x) - (2 - 2x^2)] \\ &= x(x - 1)(2x - 1).\end{aligned}$$

# Idea of the replicator equation

The replicator equation **stipulates** that the change in proportion of a species over time is determined by the difference between its fitness and the average fitness:

$$\dot{x} \left( = \frac{\partial x}{\partial t} \right) \stackrel{\text{Def}}{=} x[f_H - \bar{f}].$$

So in the Hawk-dove scenario:  $\dot{x} = x[f_H - \bar{f}]$   
 $= x[3(1 - x) - (2 - 2x^2)]$   
 $= x(x - 1)(2x - 1).$



# Relation between (stable) rest points and ESSs



# Relation between (stable) rest points and ESSs

- For  $2 \times 2$  matrices:

# Relation between (stable) rest points and ESSs

- For  $2 \times 2$  matrices:
  - There is at least one ESS.

# Relation between (stable) rest points and ESSs

- For  $2 \times 2$  matrices:
  - There is at least one ESS.
  - ESSs coincide with stable rest points of the replicator dynamic.

# Relation between (stable) rest points and ESSs

- For  $2 \times 2$  matrices:
  - There is at least one ESS.
  - ESSs coincide with stable rest points of the replicator dynamic.
- For  $m \times m$  matrices,  $m \geq 2$ :

# Relation between (stable) rest points and ESSs

- For  $2 \times 2$  matrices:
  - There is at least one ESS.
  - ESSs coincide with stable rest points of the replicator dynamic.
- For  $m \times m$  matrices,  $m \geq 2$ :
  - Fully mixed rest points of the replicator dynamics coincide with fully mixed (symmetric) Nash equilibria.

# Relation between (stable) rest points and ESSs

## ■ For $2 \times 2$ matrices:

- There is at least one ESS.
- ESSs coincide with stable rest points of the replicator dynamic.

## ■ For $m \times m$ matrices, $m \geq 2$ :

- Fully mixed rest points of the replicator dynamics coincide with fully mixed (symmetric) Nash equilibria.
- There are also connections between stability of rest points and further constraints on Nash equilibria.

# Relation between (stable) rest points and ESSs

## ■ For $2 \times 2$ matrices:

- There is at least one ESS.
- ESSs coincide with stable rest points of the replicator dynamic.


## ■ For $m \times m$ matrices, $m \geq 2$ :

- Fully mixed rest points of the replicator dynamics coincide with fully mixed (symmetric) Nash equilibria.
- There are also connections between stability of rest points and further constraints on Nash equilibria.

These are further discussed in, e.g.,

\* H. Peters (2008): *Game Theory: A Multi-Leveled Approach*. Springer, ISBN: 978-3-540-69290-4. Ch. 15: Evolutionary Games.

# Implications







SN = strict Nash, ESS - evol'y stable strategy, GSS = glob'y stable state, ASS = asymp'y stable state, NSS = neutrally stable strategy, LP = limit point, LSS = Lyapunov stable state, NE = Nash eq., FP = fixed point, \* = only if fully mixed, i = isolated Nash eq.

Dotted lines are indirect implications.







# The dynamics of the replicator equation

# Possible dynamics







# Possible dynamics



## 1. Stability.


# Possible dynamics




## 1. Stability.

Convergence to a fixed proportion.



# Possible dynamics




**1. Stability.**  
Convergence to a  
fixed proportion.




**2. Periodicity.**





# Possible dynamics




**1. Stability.**  
Convergence to a  
fixed proportion.




**2. Periodicity.**  
Convergence to  
a cycle of  
proportions





# Possible dynamics




1. **Stability.**  
Convergence to a  
fixed proportion.




2. **Periodicity.**  
Convergence to  
a cycle of  
proportions,  
with some  
period  $n \geq 2$ .




# Possible dynamics




1. **Stability.**  
Convergence to a fixed proportion.



2. **Periodicity.**  
Convergence to a cycle of proportions, with some period  $n \geq 2$ .




3. **Semi-periodicity.**







# Possible dynamics




1. **Stability.**  
Convergence to a fixed proportion.




2. **Periodicity.**  
Convergence to a cycle of proportions, with some period  $n \geq 2$ .




3. **Semi-periodicity.**  
Quasi-periodic behaviour. (E.g., if the big hand of a clock would move  $\sqrt{35} \approx 6$  instead of the usual 6 degrees per tick.)




# Possible dynamics




1. **Stability.**  
Convergence to a fixed proportion.



2. **Periodicity.**  
Convergence to a cycle of proportions, with some period  $n \geq 2$ .




3. **Semi-periodicity.**  
Quasi-periodic behaviour. (E.g., if the big hand of a clock would move  $\sqrt{35} \approx 6$  instead of the usual 6 degrees per tick.)




4. **Chaos.**


# Possible dynamics




1. **Stability.**  
Convergence to a fixed proportion.



2. **Periodicity.**  
Convergence to a cycle of proportions, with some period  $n \geq 2$ .




3. **Semi-periodicity.**  
Quasi-periodic behaviour. (E.g., if the big hand of a clock would move  $\sqrt{35} \approx 6$  instead of the usual 6 degrees per tick.)




4. **Chaos.**  
Like 1, 2 and 3 deterministic, but otherwise different.  
Predictable only by execution.


# Possible dynamics




1. **Stability.**  
Convergence to a fixed proportion.



2. **Periodicity.**  
Convergence to a cycle of proportions, with some period  $n \geq 2$ .




3. **Semi-periodicity.**  
Quasi-periodic behaviour. (E.g., if the big hand of a clock would move  $\sqrt{35} \approx 6$  instead of the usual 6 degrees per tick.)



4. **Chaos.**  
Like 1, 2 and 3 deterministic, but otherwise different.  
Predictable only by execution.

Images from: *The Computational Beauty of Nature*, W.G. Flake (1998).


# The dynamics of the replicator equation



Relative score matrix  $A = \begin{pmatrix} 1 & 3 & 1 \\ 1 & 2 & 3 \\ 4 & 1 & 3 \end{pmatrix}$ , initial proportions  $p = \begin{pmatrix} 1/3 \\ 1/3 \\ 1/3 \end{pmatrix}$ .



# The dynamics of the replicator equation



# Symmetric 2x2 games

	L	R
T	$R, R$	$S, T$
B	$T, S$	$P, P$

# Symmetric 2x2 games

	L	R
T	$R, R$	$S, T$
B	$T, S$	$P, P$

- $R$ ,  $S$ ,  $T$ , and  $P$  represent payoffs named *reward*, *sucker*, *temptation* and *punishment*, respectively.



# Symmetric 2x2 games

	L	R
T	$R, R$	$S, T$
B	$T, S$	$P, P$

- $R$ ,  $S$ ,  $T$ , and  $P$  represent payoffs named *reward*, *sucker*, *temptation* and *punishment*, respectively.
- WLOG it may be assumed that  $R > P$ . If not, then swap the actions.

# Symmetric 2x2 games

	L	R
T	$R, R$	$S, T$
B	$T, S$	$P, P$

- $R$ ,  $S$ ,  $T$ , and  $P$  represent payoffs named *reward*, *sucker*, *temptation* and *punishment*, respectively.
- WLOG it may be assumed that  $R > P$ . If not, then swap the actions.
- WLOG the payoffs can be normalised so that  $R = 1$  and  $P = 0$ .

# Symmetric 2x2 games

	L	R
T	$R, R$	$S, T$
B	$T, S$	$P, P$

- $R$ ,  $S$ ,  $T$ , and  $P$  represent payoffs named *reward*, *sucker*, *temptation* and *punishment*, respectively.
- WLOG it may be assumed that  $R > P$ . If not, then swap the actions.
- WLOG the payoffs can be normalised so that  $R = 1$  and  $P = 0$ .
- So all interesting symmetric  $2 \times 2$  games can be generated by

$$(S, T) \in [-\epsilon, 1 + \epsilon] \times [-\epsilon, 1 + \epsilon], \quad R = 1, \quad P = 0, \quad \epsilon > 0.$$

# Symmetric 2x2 games


	L	R
T	$R, R$	$S, T$
B	$T, S$	$P, P$

- $R$ ,  $S$ ,  $T$ , and  $P$  represent payoffs named *reward*, *sucker*, *temptation* and *punishment*, respectively.
- WLOG it may be assumed that  $R > P$ . If not, then swap the actions.
- WLOG the payoffs can be normalised so that  $R = 1$  and  $P = 0$ .
- So all interesting symmetric  $2 \times 2$  games can be generated by

$$(S, T) \in [-\epsilon, 1 + \epsilon] \times [-\epsilon, 1 + \epsilon], \quad R = 1, \quad P = 0, \quad \epsilon > 0.$$

- For simplicity, take  $\epsilon = 1$  so that  $(S, T)$  is in  $[-1, 2] \times [-1, 2]$ .

# The (S,T) plane



Partition of the  $(S, T)$  plane which displays various symmetric  $2 \times 2$  games.

# A discrete replicator equation

# A discrete replicator equation

- Of the existing discrete replicator equations, two of them are well known. The simplest of these two is

$$x_{t+1}^i = x_t^i + x_t^i(f_t^i - \bar{f}_t), \quad (1)$$

where  $x^i$  represents the ratio (proportion) of species  $i$ ,  $f^i$  represents the fitness of species  $i$ , and  $\bar{f}$  represents the average fitness.

# A discrete replicator equation

- Of the existing discrete replicator equations, two of them are well known. The simplest of these two is

$$x_{t+1}^i = x_t^i + x_t^i(f_t^i - \bar{f}_t), \quad (1)$$

where  $x^i$  represents the ratio (proportion) of species  $i$ ,  $f^i$  represents the fitness of species  $i$ , and  $\bar{f}$  represents the average fitness.

- Eq. (1) can perhaps best be understood by assuming that the growth of  $x_i$  in one time unit is entirely determined by the difference in fitness and average fitness:

$$\frac{x_{t+1}^i - x_t^i}{1} = x_t^i(f_t^i - \bar{f}_t). \quad (2)$$



# A discrete replicator equation

- Of the existing discrete replicator equations, two of them are well known. The simplest of these two is

$$x_{t+1}^i = x_t^i + x_t^i(f_t^i - \bar{f}_t), \quad (1)$$

where  $x^i$  represents the ratio (proportion) of species  $i$ ,  $f^i$  represents the fitness of species  $i$ , and  $\bar{f}$  represents the average fitness.

- Eq. (1) can perhaps best be understood by assuming that the growth of  $x_i$  in one time unit is entirely determined by the difference in fitness and average fitness:

$$\frac{x_{t+1}^i - x_t^i}{1} = x_t^i(f_t^i - \bar{f}_t). \quad (2)$$

In this way, Eq. (2) very much resembles the continuous replicator equation  $\dot{x}^i = x_t^i(f_t^i - \bar{f}_t)$ .

# A discrete replicator equation

# A discrete replicator equation

- Suppose the RSTP-game is played,  $x_t$  represents the proportion with which the first action is played,  $1 - x_t$  represents the proportion with which the second action is played, and the payoffs are plugged into the discrete replicator equation (1).

# A discrete replicator equation

- Suppose the RSTP-game is played,  $x_t$  represents the proportion with which the first action is played,  $1 - x_t$  represents the proportion with which the second action is played, and the payoffs are plugged into the discrete replicator equation (1).
- We obtain

$$f_t^1 = x_t R + (1 - x_t) S, \quad f_t^2 = x_t T + (1 - x_t) P, \quad \bar{f}_t = x_t f_t^1 + (1 - x_t) f_t^2, \\ x_{t+1} = x_t + x_t (f_t^1 - \bar{f}_t) \tag{3}$$

$$= x_t + x_t (1 - x_t) [S - P + (P - S + R - T)x_t]. \tag{4}$$

The step from (3) to (4) follows with some algebra.

# A discrete replicator equation

- Suppose the RSTP-game is played,  $x_t$  represents the proportion with which the first action is played,  $1 - x_t$  represents the proportion with which the second action is played, and the payoffs are plugged into the discrete replicator equation (1).
- We obtain


$$f_t^1 = x_t R + (1 - x_t) S, \quad f_t^2 = x_t T + (1 - x_t) P, \quad \bar{f}_t = x_t f_t^1 + (1 - x_t) f_t^2, \\ x_{t+1} = x_t + x_t (f_t^1 - \bar{f}_t) \tag{3}$$

$$= x_t + x_t (1 - x_t) [S - P + (P - S + R - T)x_t]. \tag{4}$$


The step from (3) to (4) follows with some algebra.

- If (4) with  $R = 1$  and  $P = 0$  is iterated 150 times on the  $(S, T)$  square  $[-8.5, 21.5] \times [-8.5, 21.5]$  with random start values picked from  $[0, 1]$  and the first 50 iterations are thrown away (the so-called *transient phase*), we obtain [next page]

# Convergence / divergence of the discrete replicator




$[-8.5, 21.5] \times [-8.5, 21.5]$




$[-5.5, 6.5] \times [-4.5, 7.5]$

**End values (if any)** in the  $(S, T)$ -plane. Red: fixed point. Dark red: low value; light red: high value. Green: divergent but bounded. Dark green: small amplitude; light green: large amplitude. Blue: divergent.

# Bifurcation plot of the discrete replicator



$[-8.5, 21.5] \times [-8.5, 21.5]$




$[2.5, 12.5] \times [1.5, 11.5]$


**Periods (if any)** in the  $(S, T)$ -plane. Grey: 0 (chaotic); red: period 1 (fixed point); orange: period 2; brown: period 3; yellow: 4; green: 5; lime: 6; turquoise: 7; cyan: 8; sky: 9; blue: 10; violet: 11; magenta: 12; pink: 13; white: 16; black: greater than 13 but  $\neq 16$ .



# Bifurcation plot of the discrete replicator



$[5, 8] \times [5.5, 8.5]$



$[6.4, 6.6] \times [6.9, 7.1]$

**Periods (if any)** in the  $(S, T)$ -plane. Grey: 0 (chaotic); red: period 1 (fixed point); orange: period 2; brown: period 3; yellow: 4; green: 5; lime: 6; turquoise: 7; cyan: 8; sky: 9; blue: 10; violet: 11; magenta: 12; pink: 13; white: 16; black: greater than 13 but  $\neq 16$ .




# Replicator dynamic for two competing populations


# One population




# One population




# One population



# One population



# One population



One population: intra-population interaction.

# Two competing populations




# Two competing populations







# Two competing populations



# Two competing populations



# Two competing populations



Two populations: inter-population interaction.

# Replicator dynamic for two competing populations

Consider interaction between two competing populations, “the row players” and “the column players”:

	L	R
T	0,0	2,2
B	1,5	1,5

# Replicator dynamic for two competing populations

Consider interaction between two competing populations, “the row players” and “the column players”:

	L	R
T	0,0	2,2
B	1,5	1,5

- Assume proportion of row plays  $T$  is  $x$ ; prop. of column plays  $L$  is  $y$ .

# Replicator dynamic for two competing populations

Consider interaction between two competing populations, “the row players” and “the column players”:

	L	R
T	0,0	2,2
B	1,5	1,5

- Assume proportion of row plays  $T$  is  $x$ ; prop. of column plays  $L$  is  $y$ .
- Expected payoff for  $T$ -players :

$$f_T = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 2 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} y \\ 1-y \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 2-2y \\ 1 \end{pmatrix} = 2-2y.$$

# Replicator dynamic for two competing populations

Consider interaction between two competing populations, “the row players” and “the column players”:

	L	R
T	0,0	2,2
B	1,5	1,5

- Assume proportion of row plays  $T$  is  $x$ ; prop. of column plays  $L$  is  $y$ .
- Expected payoff for  $T$ -players :

$$f_T = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 2 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} y \\ 1-y \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 2-2y \\ 1 \end{pmatrix} = 2-2y.$$

- Similarly, the expected payoff for  $B$ -players is 1.

# Replicator dynamic for two competing populations

Consider interaction between two competing populations, “the row players” and “the column players”:

	L	R
T	0,0	2,2
B	1,5	1,5

- Assume proportion of row plays  $T$  is  $x$ ; prop. of column plays  $L$  is  $y$ .
- Expected payoff for  $T$ -players :

$$f_T = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 2 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} y \\ 1-y \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 2-2y \\ 1 \end{pmatrix} = 2-2y.$$

- Similarly, the expected payoff for  $B$ -players is 1.
- Expected payoff for the row population:  $\bar{f} = x(2-2y) + (1-x)1$ .



# Replicator dynamic for two competing populations

Consider interaction between two competing populations, “the row players” and “the column players”:

	L	R
T	0,0	2,2
B	1,5	1,5

- Assume proportion of row plays  $T$  is  $x$ ; prop. of column plays  $L$  is  $y$ .
- Expected payoff for  $T$ -players :

$$f_T = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 2 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} y \\ 1-y \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 2-2y \\ 1 \end{pmatrix} = 2-2y.$$

- Similarly, the expected payoff for  $B$ -players is 1.
- Expected payoff for the row population:  $\bar{f} = x(2-2y) + (1-x)1$ .
- Replicator equation for the row players:  $\dot{x} = x[f_T - \bar{f}] = x[(2-2y) - (x(2-2y) + (1-x)1)] = x(1-x)(1-2y)$ .

# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).  $y = 1/2$
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).

1

0

$x$


1

# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).






# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).






# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).




# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

- Determine  $(x, y)$  where  $\dot{x} = 0$  (blue).
- Determine  $(x, y)$  where  $\dot{y} = 0$  (red).
- Determine dynamics elsewhere for  $x$  (red) and  $y$  (blue).



# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

Rest points at  $x = 0$ ,


$$(x, y) = (1, 0),$$

$$(x, y) = (1, 1).$$

Stable rest point at

$$(x, y) = (1, 0).$$

Corresponding vector density plot:



# Replicator dynamic for two competing populations

Similarly for the share  $y$  of column players that play  $L$ . We obtain a system of differential equations:

$$\begin{cases} \dot{x} = x(x-1)(2y-1) \\ \dot{y} = 2xy(y-1) \end{cases}$$

Rest points at

$$\begin{aligned} x &= 0, \\ (x, y) &= (1, 0), \\ (x, y) &= (1, 1). \end{aligned}$$

Stable rest point at

$$(x, y) = (1, 0).$$

Corresponding stream density plot:

