

# Multi-agent learning

## Gradient ascent

*Gerard Vreeswijk*, Intelligent Software Systems, Computer Science  
Department, Faculty of Sciences, Utrecht University, The  
Netherlands.

Saturday 16<sup>th</sup> May, 2020

# Gradient ascent: idea

## Idea

# Gradient ascent: idea

## Idea

- Every opponent is identified with a (possibly mixed) strategy.

# Gradient ascent: idea

## Idea

- Every opponent is identified with a (possibly mixed) strategy.
- Players can observe the (possibly mixed) strategy of their opponent.

# Gradient ascent: idea

## Idea

- Every opponent is identified with a (possibly mixed) strategy.
- Players can observe the (possibly mixed) strategy of their opponent.
- After observation, every player changes its strategy a tiny bit in the right direction.

# Gradient ascent: idea

## Idea

- Every opponent is identified with a (possibly mixed) strategy.
- Players can observe the (possibly mixed) strategy of their opponent.
- After observation, every player changes its strategy a tiny bit in the right direction.
- Comparison with fictitious play.

# Gradient ascent: idea

## Idea

- Every opponent is identified with a (possibly mixed) strategy.
- Players can observe the (possibly mixed) strategy of their opponent.
- After observation, every player changes its strategy a tiny bit in the right direction.
- Comparison with fictitious play.
  - Like in fictitious play, opponents are modelled through a mixed strategy.

# Gradient ascent: idea

## Idea

- Every opponent is identified with a (possibly mixed) strategy.
- Players can observe the (possibly mixed) strategy of their opponent.
- After observation, every player changes its strategy a tiny bit in the right direction.
- Comparison with fictitious play.
  - Like in fictitious play, opponents are modelled through a mixed strategy.
  - In fictitious play, players learn projected opponent strategies, and play a best response to it.



# Gradient ascent: idea

## Idea

- Every opponent is identified with a (possibly mixed) strategy.
- Players can observe the (possibly mixed) strategy of their opponent.
- After observation, every player changes its strategy a tiny bit in the right direction.
- Comparison with fictitious play.
  - Like in fictitious play, opponents are modelled through a mixed strategy.
  - In fictitious play, players learn projected opponent strategies, and play a best response to it.
  - In gradient ascent, players do not project a mixed strategy, and do not play a best response.

# Plan for today

# Plan for today

1. Two-player, two-action, general sum games with real payoffs.

# Plan for today

1. Two-player, two-action, general sum games with real payoffs.
2. Dynamics of (mixed) strategies in such games.

# Plan for today

1. Two-player, two-action, general sum games with real payoffs.
2. Dynamics of (mixed) strategies in such games.

Examples:

# Plan for today

1. Two-player, two-action, general sum games with real payoffs.
2. Dynamics of (mixed) strategies in such games.

Examples:

(a) Coordination game

# Plan for today

1. Two-player, two-action, general sum games with real payoffs.
2. Dynamics of (mixed) strategies in such games.

Examples:

- (a) Coordination game
- (b) Prisoners' dilemma

# Plan for today

1. Two-player, two-action, general sum games with real payoffs.
2. Dynamics of (mixed) strategies in such games.

Examples:

(a) Coordination game

(c) Stag hunt

(b) Prisoners' dilemma



# Plan for today

1. Two-player, two-action, general sum games with real payoffs.
2. Dynamics of (mixed) strategies in such games.

Examples:

- |                        |                    |
|------------------------|--------------------|
| (a) Coordination game  | (c) Stag hunt      |
| (b) Prisoners' dilemma | (d) Other examples |

# Plan for today

1. Two-player, two-action, general sum games with real payoffs.
2. Dynamics of (mixed) strategies in such games.

Examples:

- |                        |                    |
|------------------------|--------------------|
| (a) Coordination game  | (c) Stag hunt      |
| (b) Prisoners' dilemma | (d) Other examples |

3. **IGA: Infinitesimal Gradient Ascent.** Singh, Kearns and Mansour (2000).

# Plan for today

1. Two-player, two-action, general sum games with real payoffs.
2. Dynamics of (mixed) strategies in such games.

Examples:

- |                        |                    |
|------------------------|--------------------|
| (a) Coordination game  | (c) Stag hunt      |
| (b) Prisoners' dilemma | (d) Other examples |

3. **IGA: Infinitesimal Gradient Ascent.** Singh, Kearns and Mansour (2000).

■ Convergence of IGA.

# Plan for today

1. Two-player, two-action, general sum games with real payoffs.
2. Dynamics of (mixed) strategies in such games.

Examples:

- |                        |                    |
|------------------------|--------------------|
| (a) Coordination game  | (c) Stag hunt      |
| (b) Prisoners' dilemma | (d) Other examples |

3. **IGA: Infinitesimal Gradient Ascent.** Singh, Kearns and Mansour (2000).

■ Convergence of IGA.

4. **IGA-WoLF: Win or Learn Fast.** Bowling and Veloso (2001, 2002).

# Plan for today

1. Two-player, two-action, general sum games with real payoffs.
2. Dynamics of (mixed) strategies in such games.

Examples:

- |                        |                    |
|------------------------|--------------------|
| (a) Coordination game  | (c) Stag hunt      |
| (b) Prisoners' dilemma | (d) Other examples |

3. **IGA: Infinitesimal Gradient Ascent.** Singh, Kearns and Mansour (2000).

- Convergence of IGA.

4. **IGA-WoLF: Win or Learn Fast.** Bowling and Veloso (2001, 2002).

- Convergence of IGA-WoLF

# Plan for today

1. Two-player, two-action, general sum games with real payoffs.
2. Dynamics of (mixed) strategies in such games.

Examples:

- |                        |                    |
|------------------------|--------------------|
| (a) Coordination game  | (c) Stag hunt      |
| (b) Prisoners' dilemma | (d) Other examples |

3. **IGA: Infinitesimal Gradient Ascent**. Singh, Kearns and Mansour (2000).

- Convergence of IGA.

4. **IGA-WoLF: Win or Learn Fast**. Bowling and Veloso (2001, 2002).

- Convergence of IGA-WoLF + analysis of the proof of convergence.

# **Part 1:**

## **Payoffs of general 2x2 games in normal form**

# Two-player, two-action, general sum games

In its most general form, a two-player, two-action game in normal form with real-valued payoffs can be represented by

$$M = \begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} r_{11}, c_{11} & r_{12}, c_{12} \\ r_{21}, c_{21} & r_{22}, c_{22} \end{array} \right) \end{array}$$



# Two-player, two-action, general sum games

In its most general form, a two-player, two-action game in normal form with real-valued payoffs can be represented by

$$M = \begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} r_{11}, c_{11} & r_{12}, c_{12} \\ r_{21}, c_{21} & r_{22}, c_{22} \end{array} \right) \end{array}$$

Row plays mixed  $(\alpha, 1 - \alpha)$ . Column plays mixed  $(\beta, 1 - \beta)$ .

# Two-player, two-action, general sum games

In its most general form, a two-player, two-action game in normal form with real-valued payoffs can be represented by

$$M = \begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} r_{11}, c_{11} & r_{12}, c_{12} \\ r_{21}, c_{21} & r_{22}, c_{22} \end{array} \right) \end{array}$$

Row plays mixed  $(\alpha, 1 - \alpha)$ . Column plays mixed  $(\beta, 1 - \beta)$ .

Expected payoff:

# Two-player, two-action, general sum games

In its most general form, a two-player, two-action game in normal form with real-valued payoffs can be represented by

$$M = \begin{array}{c} \text{T} \\ \text{B} \end{array} \begin{array}{cc} \text{L} & \text{R} \\ \left( \begin{array}{cc} r_{11}, c_{11} & r_{12}, c_{12} \\ r_{21}, c_{21} & r_{22}, c_{22} \end{array} \right) \end{array}$$

Row plays mixed  $(\alpha, 1 - \alpha)$ . Column plays mixed  $(\beta, 1 - \beta)$ .

Expected payoff:

$$u_1(\alpha, \beta)$$

# Two-player, two-action, general sum games

In its most general form, a two-player, two-action game in normal form with real-valued payoffs can be represented by

$$M = \begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} r_{11}, c_{11} & r_{12}, c_{12} \\ r_{21}, c_{21} & r_{22}, c_{22} \end{array} \right) \end{array}$$

Row plays mixed  $(\alpha, 1 - \alpha)$ . Column plays mixed  $(\beta, 1 - \beta)$ .

Expected payoff:

$$u_1(\alpha, \beta) = \alpha[\beta r_{11} + (1 - \beta)r_{12}] + (1 - \alpha)[\beta r_{21} + (1 - \beta)r_{22}]$$

# Two-player, two-action, general sum games

In its most general form, a two-player, two-action game in normal form with real-valued payoffs can be represented by

$$M = \begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} r_{11}, c_{11} & r_{12}, c_{12} \\ r_{21}, c_{21} & r_{22}, c_{22} \end{array} \right) \end{array}$$

Row plays mixed  $(\alpha, 1 - \alpha)$ . Column plays mixed  $(\beta, 1 - \beta)$ .

Expected payoff:

$$\begin{aligned} u_1(\alpha, \beta) &= \alpha[\beta r_{11} + (1 - \beta)r_{12}] + (1 - \alpha)[\beta r_{21} + (1 - \beta)r_{22}] \\ &= \alpha\beta + \alpha(r_{12} - r_{11}) + \beta(r_{21} - r_{11}) + r_{11}. \end{aligned}$$

# Two-player, two-action, general sum games

In its most general form, a two-player, two-action game in normal form with real-valued payoffs can be represented by

$$M = \begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} r_{11}, c_{11} & r_{12}, c_{12} \\ r_{21}, c_{21} & r_{22}, c_{22} \end{array} \right) \end{array}$$

Row plays mixed  $(\alpha, 1 - \alpha)$ . Column plays mixed  $(\beta, 1 - \beta)$ .

Expected payoff:

$$\begin{aligned} u_1(\alpha, \beta) &= \alpha[\beta r_{11} + (1 - \beta)r_{12}] + (1 - \alpha)[\beta r_{21} + (1 - \beta)r_{22}] \\ &= \alpha\beta + \alpha(r_{12} - r_{11}) + \beta(r_{21} - r_{11}) + r_{11}. \end{aligned}$$

$$u_2(\alpha, \beta)$$

# Two-player, two-action, general sum games

In its most general form, a two-player, two-action game in normal form with real-valued payoffs can be represented by

$$M = \begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} r_{11}, c_{11} & r_{12}, c_{12} \\ r_{21}, c_{21} & r_{22}, c_{22} \end{array} \right) \end{array}$$

Row plays mixed  $(\alpha, 1 - \alpha)$ . Column plays mixed  $(\beta, 1 - \beta)$ .

Expected payoff:

$$\begin{aligned} u_1(\alpha, \beta) &= \alpha[\beta r_{11} + (1 - \beta)r_{12}] + (1 - \alpha)[\beta r_{21} + (1 - \beta)r_{22}] \\ &= \alpha\beta r_{11} + \alpha(1 - \beta)r_{12} + (1 - \alpha)\beta r_{21} + (1 - \alpha)(1 - \beta)r_{22} \\ &= \alpha\beta(r_{11} - r_{12} - r_{21} + r_{22}) + \alpha(1 - \beta)r_{12} + \beta(1 - \alpha)r_{21} + (1 - \alpha)(1 - \beta)r_{22} \\ u_2(\alpha, \beta) &= \beta[\alpha c_{11} + (1 - \alpha)c_{21}] + (1 - \beta)[\alpha c_{12} + (1 - \alpha)c_{22}] \\ &= \alpha\beta c_{11} + \beta(1 - \alpha)c_{21} + (1 - \beta)\alpha c_{12} + (1 - \beta)(1 - \alpha)c_{22} \\ &= \alpha\beta(c_{11} - c_{12} - c_{21} + c_{22}) + \beta(1 - \alpha)c_{21} + \alpha(1 - \beta)c_{12} + (1 - \alpha)(1 - \beta)c_{22} \end{aligned}$$

# Two-player, two-action, general sum games

In its most general form, a two-player, two-action game in normal form with real-valued payoffs can be represented by

$$M = \begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} r_{11}, c_{11} & r_{12}, c_{12} \\ r_{21}, c_{21} & r_{22}, c_{22} \end{array} \right) \end{array}$$

Row plays mixed  $(\alpha, 1 - \alpha)$ . Column plays mixed  $(\beta, 1 - \beta)$ .

Expected payoff:

$$u_1(\alpha, \beta) = \alpha[\beta r_{11} + (1 - \beta)r_{12}] + (1 - \alpha)[\beta r_{21} + (1 - \beta)r_{22}]$$

$$= u \alpha \beta + \alpha(r_{12} - r_{22}) + \beta(r_{21} - r_{22}) + r_{22}.$$

$$u_2(\alpha, \beta) = \beta[\alpha c_{11} + (1 - \alpha)c_{21}] + (1 - \beta)[\alpha c_{12} + (1 - \alpha)c_{22}]$$

$$= u' \alpha \beta + \alpha(c_{21} - c_{22}) + \beta(c_{12} - c_{22}) + c_{22}.$$

where  $u = (r_{11} - r_{12}) - (r_{21} - r_{22})$  and  $u' = (c_{11} - c_{21}) - (c_{12} - c_{22})$ .



# Gradient of expected payoff

Gradient:

$$\frac{\partial u_1(\alpha, \beta)}{\partial \alpha} = \beta u + (r_{12} - r_{22})$$

$$\frac{\partial u_2(\alpha, \beta)}{\partial \beta} = \alpha u' + (c_{21} - c_{22})$$

# Gradient of expected payoff

Gradient:

$$\frac{\partial u_1(\alpha, \beta)}{\partial \alpha} = \beta u + (r_{12} - r_{22})$$
$$\frac{\partial u_2(\alpha, \beta)}{\partial \beta} = \alpha u' + (c_{21} - c_{22})$$

As an affine map:

$$\begin{aligned} \begin{bmatrix} \partial u_1 / \partial \alpha \\ \partial u_2 / \partial \beta \end{bmatrix} &= \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \\ &\quad + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \\ &= U \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + C \end{aligned}$$

# Gradient of expected payoff

Gradient:

$$\begin{aligned}\frac{\partial u_1(\alpha, \beta)}{\partial \alpha} &= \beta u + (r_{12} - r_{22}) \\ \frac{\partial u_2(\alpha, \beta)}{\partial \beta} &= \alpha u' + (c_{21} - c_{22})\end{aligned}$$

Stationary point:

$$(\alpha^*, \beta^*) = \left( \frac{c_{22} - c_{21}}{u'}, \frac{r_{22} - r_{12}}{u} \right)$$

As an affine map:

$$\begin{aligned}\begin{bmatrix} \partial u_1 / \partial \alpha \\ \partial u_2 / \partial \beta \end{bmatrix} &= \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \\ &\quad + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \\ &= U \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + C\end{aligned}$$

# Gradient of expected payoff

Gradient:

$$\begin{aligned}\frac{\partial u_1(\alpha, \beta)}{\partial \alpha} &= \beta u + (r_{12} - r_{22}) \\ \frac{\partial u_2(\alpha, \beta)}{\partial \beta} &= \alpha u' + (c_{21} - c_{22})\end{aligned}$$

Stationary point:

$$(\alpha^*, \beta^*) = \left( \frac{c_{22} - c_{21}}{u'}, \frac{r_{22} - r_{12}}{u} \right)$$

Remarks:

As an affine map:

$$\begin{aligned}\begin{bmatrix} \partial u_1 / \partial \alpha \\ \partial u_2 / \partial \beta \end{bmatrix} &= \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \\ &\quad + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \\ &= U \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + C\end{aligned}$$

# Gradient of expected payoff

Gradient:

$$\begin{aligned}\frac{\partial u_1(\alpha, \beta)}{\partial \alpha} &= \beta u + (r_{12} - r_{22}) \\ \frac{\partial u_2(\alpha, \beta)}{\partial \beta} &= \alpha u' + (c_{21} - c_{22})\end{aligned}$$

As an affine map:

$$\begin{aligned}\begin{bmatrix} \partial u_1 / \partial \alpha \\ \partial u_2 / \partial \beta \end{bmatrix} &= \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \\ &\quad + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \\ &= U \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + C\end{aligned}$$

Stationary point:

$$(\alpha^*, \beta^*) = \left( \frac{c_{22} - c_{21}}{u'}, \frac{r_{22} - r_{12}}{u} \right)$$

Remarks:

- There is at most one stationary point.

# Gradient of expected payoff

Gradient:

$$\begin{aligned}\frac{\partial u_1(\alpha, \beta)}{\partial \alpha} &= \beta u + (r_{12} - r_{22}) \\ \frac{\partial u_2(\alpha, \beta)}{\partial \beta} &= \alpha u' + (c_{21} - c_{22})\end{aligned}$$

As an affine map:

$$\begin{aligned}\begin{bmatrix} \partial u_1 / \partial \alpha \\ \partial u_2 / \partial \beta \end{bmatrix} &= \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \\ &\quad + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \\ &= U \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + C\end{aligned}$$

Stationary point:

$$(\alpha^*, \beta^*) = \left( \frac{c_{22} - c_{21}}{u'}, \frac{r_{22} - r_{12}}{u} \right)$$

Remarks:

- There is at most one stationary point.
- If a stationary point exists, it may lie outside  $[0, 1]^2$ .

# Gradient of expected payoff

Gradient:

$$\begin{aligned}\frac{\partial u_1(\alpha, \beta)}{\partial \alpha} &= \beta u + (r_{12} - r_{22}) \\ \frac{\partial u_2(\alpha, \beta)}{\partial \beta} &= \alpha u' + (c_{21} - c_{22})\end{aligned}$$

As an affine map:

$$\begin{aligned}\begin{bmatrix} \partial u_1 / \partial \alpha \\ \partial u_2 / \partial \beta \end{bmatrix} &= \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \\ &\quad + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \\ &= U \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + C\end{aligned}$$

Stationary point:

$$(\alpha^*, \beta^*) = \left( \frac{c_{22} - c_{21}}{u'}, \frac{r_{22} - r_{12}}{u} \right)$$


Remarks:

- There is at most one stationary point.
- If a stationary point exists, it may lie outside  $[0, 1]^2$ .
- If there is a stationary point inside  $[0, 1]^2$ , it is a weak (i.e., non-strict) Nash equilibrium.

# Example: payoffs in Stag Hunt ( $r=4$ , $t=3$ , $s=1$ , $p=3$ )



# Example: payoffs in Stag Hunt ( $r=4$ , $t=3$ , $s=1$ , $p=3$ )



Player 1  
may  
only  
move  
“back  
–  
front”;  
Player  
2 may  
only  
move  
“left –  
right”.

# Part 2: IGA

# Gradient ascent

Affine differential map:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \begin{bmatrix} \partial u_1 / \partial \alpha \\ \partial u_2 / \partial \beta \end{bmatrix}_t$$

# Gradient ascent

Affine differential map:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \begin{bmatrix} \partial u_1 / \partial \alpha \\ \partial u_2 / \partial \beta \end{bmatrix}_t$$

- Because  $\alpha, \beta \in [0, 1]$ , the dynamics must be confined to  $[0, 1]^2$ .

# Gradient ascent

Affine differential map:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \begin{bmatrix} \partial u_1 / \partial \alpha \\ \partial u_2 / \partial \beta \end{bmatrix}_t$$

- Because  $\alpha, \beta \in [0, 1]$ , the dynamics must be confined to  $[0, 1]^2$ .
- Suppose the state  $(\alpha, \beta)$  is on the boundary of the probability space  $[0, 1]^2$ , and the gradient vector points outwards.

Intuition: one of the players has an incentive to improve, but cannot improve further.

# Gradient ascent

Affine differential map:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \begin{bmatrix} \partial u_1 / \partial \alpha \\ \partial u_2 / \partial \beta \end{bmatrix}_t$$

- Because  $\alpha, \beta \in [0, 1]$ , the dynamics must be confined to  $[0, 1]^2$ .

- Suppose the state  $(\alpha, \beta)$  is on the boundary of the probability space  $[0, 1]^2$ , and the gradient vector points outwards.

Intuition: one of the players has an incentive to improve, but cannot improve further.

- To maintain dynamics within  $[0, 1]^2$ , the gradient is projected back on to  $[0, 1]^2$ .

Intuition: if one of the players has an incentive to improve, but *cannot* improve, then he *will not* improve.

# Gradient ascent

Affine differential map:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \begin{bmatrix} \partial u_1 / \partial \alpha \\ \partial u_2 / \partial \beta \end{bmatrix}_t$$

- Because  $\alpha, \beta \in [0, 1]$ , the dynamics must be confined to  $[0, 1]^2$ .

- Suppose the state  $(\alpha, \beta)$  is on the boundary of the probability space  $[0, 1]^2$ , and the gradient vector points outwards.

Intuition: one of the players has an incentive to improve, but cannot improve further.

- To maintain dynamics within  $[0, 1]^2$ , the gradient is projected back on to  $[0, 1]^2$ .

Intuition: if one of the players has an incentive to improve, but *cannot* improve, then he *will not* improve.

- If nonzero, the projected gradient is parallel to the (closest) boundary of  $[0, 1]^2$ .

# Infinitesimal Gradient Ascent : IGA (Singh *et al.*, 2000)

Affine differential map:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \left( \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \right)$$



# Infinitesimal Gradient Ascent : IGA (Singh *et al.*, 2000)

Affine differential map:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \left( \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \right)$$

**Theorem** (Singh, Kearns and Mansour, 2000) *If players follow IGA, where  $\eta \rightarrow 0$ , their average payoffs will converge to the (expected) payoffs of a NE. If their strategies converge, they will converge to that same NE.*

# Infinitesimal Gradient Ascent : IGA (Singh *et al.*, 2000)

Affine differential map:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \left( \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \right)$$

**Theorem** (Singh, Kearns and Mansour, 2000) *If players follow IGA, where  $\eta \rightarrow 0$ , their average payoffs will converge to the (expected) payoffs of a NE. If their strategies converge, they will converge to that same NE.*

The proof is based on a qualitative result in the theory of differential equations, which says that the behaviour of an affine differential map is determined by the multiplicative matrix  $U$ :

# Infinitesimal Gradient Ascent : IGA (Singh *et al.*, 2000)

Affine differential map:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \left( \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \right)$$

**Theorem** (Singh, Kearns and Mansour, 2000) *If players follow IGA, where  $\eta \rightarrow 0$ , their average payoffs will converge to the (expected) payoffs of a NE. If their strategies converge, they will converge to that same NE.*

The proof is based on a qualitative result in the theory of differential equations, which says that the behaviour of an affine differential map is determined by the multiplicative matrix  $U$ :

1. If  $U$  is invertible, and its eigenvalue  $\lambda$  (solution of  $Ux = \lambda x \Leftrightarrow$  solution of  $\text{Det}[U - \lambda I] = 0$ ) is real,  $\exists$  stationary point

# Infinitesimal Gradient Ascent : IGA (Singh *et al.*, 2000)

Affine differential map:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \left( \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \right)$$

**Theorem** (Singh, Kearns and Mansour, 2000) *If players follow IGA, where  $\eta \rightarrow 0$ , their average payoffs will converge to the (expected) payoffs of a NE. If their strategies converge, they will converge to that same NE.*

The proof is based on a qualitative result in the theory of differential equations, which says that the behaviour of an affine differential map is determined by the multiplicative matrix  $U$ :

1. If  $U$  is invertible, and its eigenvalue  $\lambda$  (solution of  $Ux = \lambda x \Leftrightarrow$  solution of  $\text{Det}[U - \lambda I] = 0$ ) is real,  $\exists$  stationary point, w.i. is a saddle point.

# Infinitesimal Gradient Ascent : IGA (Singh *et al.*, 2000)

Affine differential map:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \left( \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \right)$$

**Theorem** (Singh, Kearns and Mansour, 2000) *If players follow IGA, where  $\eta \rightarrow 0$ , their average payoffs will converge to the (expected) payoffs of a NE. If their strategies converge, they will converge to that same NE.*

The proof is based on a qualitative result in the theory of differential equations, which says that the behaviour of an affine differential map is determined by the multiplicative matrix  $U$ :

1. If  $U$  is invertible, and its eigenvalue  $\lambda$  (solution of  $Ux = \lambda x \Leftrightarrow$  solution of  $\text{Det}[U - \lambda I] = 0$ ) is real,  $\exists$  stationary point, w.i. is a saddle point.
2. If  $U$  is invertible, and its eigenvalue  $\lambda$  is imaginary, there is a stationary point

# Infinitesimal Gradient Ascent : IGA (Singh *et al.*, 2000)

Affine differential map:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \left( \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \right)$$

**Theorem** (Singh, Kearns and Mansour, 2000) *If players follow IGA, where  $\eta \rightarrow 0$ , their average payoffs will converge to the (expected) payoffs of a NE. If their strategies converge, they will converge to that same NE.*

The proof is based on a qualitative result in the the theory of differential equations, which says that the behaviour of an affine differential map is determined by the multiplicative matrix  $U$ :

1. If  $U$  is invertible, and its eigenvalue  $\lambda$  (solution of  $Ux = \lambda x \Leftrightarrow$  solution of  $\text{Det}[U - \lambda I] = 0$ ) is real,  $\exists$  stationary point, w.i. is a saddle point.
2. If  $U$  is invertible, and its eigenvalue  $\lambda$  is imaginary, there is a stationary point, which, in particular, is a centric point.

# Infinitesimal Gradient Ascent : IGA (Singh *et al.*, 2000)

Affine differential map:


$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \left( \begin{bmatrix} 0 & u \\ u' & 0 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + \begin{bmatrix} r_{12} - r_{22} \\ c_{21} - c_{22} \end{bmatrix} \right)$$

**Theorem** (Singh, Kearns and Mansour, 2000) *If players follow IGA, where  $\eta \rightarrow 0$ , their average payoffs will converge to the (expected) payoffs of a NE. If their strategies converge, they will converge to that same NE.*

The proof is based on a qualitative result in the theory of differential equations, which says that the behaviour of an affine differential map is determined by the multiplicative matrix  $U$ :

1. If  $U$  is invertible, and its eigenvalue  $\lambda$  (solution of  $Ux = \lambda x \Leftrightarrow$  solution of  $\text{Det}[U - \lambda I] = 0$ ) is real,  $\exists$  stationary point, w.i. is a saddle point.
2. If  $U$  is invertible, and its eigenvalue  $\lambda$  is imaginary, there is a stationary point, which, in particular, is a centric point.
3. If  $U$  is not invertible (iff  $u = 0$  or  $u' = 0$ ), there is no stationary point.

# Saddle point





# Gradient ascent: Coordination game

# Gradient ascent: Coordination game

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} 1,1 & 0,0 \\ 0,0 & 1,1 \end{array} \right) \end{array}$$

# Gradient ascent: Coordination game

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} 1,1 & 0,0 \\ 0,0 & 1,1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 2 \cdot \beta - 1 \\ 2 \cdot \alpha - 1 \end{bmatrix}$$

# Gradient ascent: Coordination game

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} 1,1 & 0,0 \\ 0,0 & 1,1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 2 \cdot \beta - 1 \\ 2 \cdot \alpha - 1 \end{bmatrix}$$

- Stationary at  $(1/2, 1/2)$ .

# Gradient ascent: Coordination game

- Symmetric, but not zero sum:

$$\begin{array}{c} \text{L} \quad \text{R} \\ \text{T} \left( \begin{array}{cc} 1,1 & 0,0 \end{array} \right) \\ \text{B} \left( \begin{array}{cc} 0,0 & 1,1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 2 \cdot \beta - 1 \\ 2 \cdot \alpha - 1 \end{bmatrix}$$

- Stationary at  $(1/2, 1/2)$ .

- Matrix

$$U = \begin{bmatrix} 0 & 2 \\ 2 & 0 \end{bmatrix}$$

has real eigenvalues:  $\lambda^2 - 4 = 0$ .

# Gradient ascent: Coordination game

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} 1,1 & 0,0 \\ 0,0 & 1,1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 2 \cdot \beta - 1 \\ 2 \cdot \alpha - 1 \end{bmatrix}$$

- Stationary at  $(1/2, 1/2)$ .

- Matrix

$$U = \begin{bmatrix} 0 & 2 \\ 2 & 0 \end{bmatrix}$$

has real eigenvalues:  $\lambda^2 - 4 = 0$ .

Saddle point inside  $[0, 1]^2$ .

# Gradient ascent: Coordination game

- Symmetric, but not zero sum:

$$\begin{array}{c} \text{T} \\ \text{B} \end{array} \begin{array}{cc} \text{L} & \text{R} \\ \left( \begin{array}{cc} 1,1 & 0,0 \\ 0,0 & 1,1 \end{array} \right) \end{array}$$

- Gradient:


$$\begin{bmatrix} 2 \cdot \beta - 1 \\ 2 \cdot \alpha - 1 \end{bmatrix}$$

- Stationary at  $(1/2, 1/2)$ .

- Matrix

$$U = \begin{bmatrix} 0 & 2 \\ 2 & 0 \end{bmatrix}$$

has real eigenvalues:  $\lambda^2 - 4 = 0$ .  
Saddle point inside  $[0, 1]^2$ .



# Gradient ascent: Prisoners' Dilemma



# Gradient ascent: Prisoners' Dilemma

- Symmetric, but not zero sum:

	L	R
T	3,3	0,5
B	5,0	1,1

# Gradient ascent: Prisoners' Dilemma

- Symmetric, but not zero sum:

	L	R
T	3,3	0,5
B	5,0	1,1

- Gradient:

$$\begin{bmatrix} -1 \cdot \beta - 1 \\ -1 \cdot \alpha - 1 \end{bmatrix}$$

# Gradient ascent: Prisoners' Dilemma

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} 3,3 & 0,5 \\ 5,0 & 1,1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -1 \cdot \beta - 1 \\ -1 \cdot \alpha - 1 \end{bmatrix}$$

- Stationary at  $(-1, -1)$ .

# Gradient ascent: Prisoners' Dilemma

- Symmetric, but not zero sum:

$$\begin{array}{c} \text{T} \\ \text{B} \end{array} \begin{array}{cc} \text{L} & \text{R} \\ \left( \begin{array}{cc} 3,3 & 0,5 \\ 5,0 & 1,1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -1 \cdot \beta - 1 \\ -1 \cdot \alpha - 1 \end{bmatrix}$$

- Stationary at  $(-1, -1)$ .

- Matrix

$$U = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

has real eigenvalues:  $\lambda^2 - 1 = 0$ .

# Gradient ascent: Prisoners' Dilemma

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} 3,3 & 0,5 \\ 5,0 & 1,1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -1 \cdot \beta - 1 \\ -1 \cdot \alpha - 1 \end{bmatrix}$$

- Stationary at  $(-1, -1)$ .

- Matrix

$$U = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

has real eigenvalues:  $\lambda^2 - 1 = 0$ .

Saddle point outside  $[0, 1]^2$ .

# Gradient ascent: Prisoners' Dilemma

- Symmetric, but not zero sum:

$$\begin{array}{c} \text{T} \\ \text{B} \end{array} \begin{array}{cc} \text{L} & \text{R} \\ \left( \begin{array}{cc} 3,3 & 0,5 \\ 5,0 & 1,1 \end{array} \right) \end{array}$$

- Gradient:


$$\begin{bmatrix} -1 \cdot \beta - 1 \\ -1 \cdot \alpha - 1 \end{bmatrix}$$

- Stationary at  $(-1, -1)$ .

- Matrix

$$U = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

has real eigenvalues:  $\lambda^2 - 1 = 0$ .  
Saddle point outside  $[0, 1]^2$ .



# Gradient ascent: Stag hunt

# Gradient ascent: Stag hunt

- Symmetric, but not zero sum:

	L	R
T	5,5	0,3
B	3,0	2,2



# Gradient ascent: Stag hunt

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} 5,5 & 0,3 \\ 3,0 & 2,2 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 4 \cdot \beta - 2 \\ 4 \cdot \alpha - 2 \end{bmatrix}$$

# Gradient ascent: Stag hunt

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} 5,5 & 0,3 \\ 3,0 & 2,2 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 4 \cdot \beta - 2 \\ 4 \cdot \alpha - 2 \end{bmatrix}$$

- Stationary at  $(1/2, 1/2)$ .

# Gradient ascent: Stag hunt

- Symmetric, but not zero sum:

$$\begin{array}{c} \text{L} \quad \text{R} \\ \text{T} \left( \begin{array}{cc} 5,5 & 0,3 \end{array} \right) \\ \text{B} \left( \begin{array}{cc} 3,0 & 2,2 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 4 \cdot \beta - 2 \\ 4 \cdot \alpha - 2 \end{bmatrix}$$

- Stationary at  $(1/2, 1/2)$ .

- Matrix

$$U = \begin{bmatrix} 0 & 4 \\ 4 & 0 \end{bmatrix}$$

has real eigenvalues:

$$\lambda^2 - 16 = 0.$$

# Gradient ascent: Stag hunt

- Symmetric, but not zero sum:

$$\begin{array}{c} \text{L} \quad \text{R} \\ \text{T} \left( \begin{array}{cc} 5,5 & 0,3 \end{array} \right) \\ \text{B} \left( \begin{array}{cc} 3,0 & 2,2 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 4 \cdot \beta - 2 \\ 4 \cdot \alpha - 2 \end{bmatrix}$$

- Stationary at  $(1/2, 1/2)$ .

- Matrix

$$U = \begin{bmatrix} 0 & 4 \\ 4 & 0 \end{bmatrix}$$

has real eigenvalues:

$\lambda^2 - 16 = 0$ . Saddle point  
inside  $[0, 1]^2$ .

# Gradient ascent: Stag hunt

- Symmetric, but not zero sum:

$$\begin{array}{c} \text{L} \quad \text{R} \\ \text{T} \left( \begin{array}{cc} 5,5 & 0,3 \\ 3,0 & 2,2 \end{array} \right) \\ \text{B} \end{array}$$

- Gradient:

$$\begin{bmatrix} 4 \cdot \beta - 2 \\ 4 \cdot \alpha - 2 \end{bmatrix}$$


- Stationary at  $(1/2, 1/2)$ .

- Matrix

$$U = \begin{bmatrix} 0 & 4 \\ 4 & 0 \end{bmatrix}$$

has real eigenvalues:

$\lambda^2 - 16 = 0$ . Saddle point  
inside  $[0, 1]^2$ .



# Gradient ascent: Game of Chicken

# Gradient ascent: Game of Chicken

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} 0,0 & -1,1 \\ 1,-1 & -3,-3 \end{array} \right) \end{array}$$

# Gradient ascent: Game of Chicken

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} 0,0 & -1,1 \\ 1,-1 & -3,-3 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -3 \cdot \beta + 2 \\ -3 \cdot \alpha + 2 \end{bmatrix}$$



# Gradient ascent: Game of Chicken

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} 0, 0 & -1, 1 \\ 1, -1 & -3, -3 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -3 \cdot \beta + 2 \\ -3 \cdot \alpha + 2 \end{bmatrix}$$

- Stationary at  $(2/3, 2/3)$ .

# Gradient ascent: Game of Chicken

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} 0, 0 & -1, 1 \\ 1, -1 & -3, -3 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -3 \cdot \beta + 2 \\ -3 \cdot \alpha + 2 \end{bmatrix}$$

- Stationary at  $(2/3, 2/3)$ .

- Matrix

$$U = \begin{bmatrix} 0 & -3 \\ -3 & 0 \end{bmatrix}$$

has real eigenvalues:  $\lambda^2 - 9 = 0$ .

# Gradient ascent: Game of Chicken

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} 0,0 & -1,1 \\ 1,-1 & -3,-3 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -3 \cdot \beta + 2 \\ -3 \cdot \alpha + 2 \end{bmatrix}$$

- Stationary at  $(2/3, 2/3)$ .

- Matrix

$$U = \begin{bmatrix} 0 & -3 \\ -3 & 0 \end{bmatrix}$$

has real eigenvalues:  $\lambda^2 - 9 = 0$ .

Saddle point inside  $[0, 1]^2$ .

# Gradient ascent: Game of Chicken

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} 0,0 & -1,1 \\ 1,-1 & -3,-3 \end{array} \right) \end{array}$$

- Gradient:


$$\begin{bmatrix} -3 \cdot \beta + 2 \\ -3 \cdot \alpha + 2 \end{bmatrix}$$

- Stationary at  $(2/3, 2/3)$ .

- Matrix

$$U = \begin{bmatrix} 0 & -3 \\ -3 & 0 \end{bmatrix}$$

has real eigenvalues:  $\lambda^2 - 9 = 0$ .  
Saddle point inside  $[0,1]^2$ .



# Gradient ascent: Battle of the Sexes

# Gradient ascent: Battle of the Sexes

- Symmetric, but not zero sum:

	L	R
T	0,0	2,3
B	3,2	1,1

# Gradient ascent: Battle of the Sexes

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} 0,0 & 2,3 \\ 3,2 & 1,1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -4 \cdot \beta + 1 \\ -4 \cdot \alpha + 1 \end{bmatrix}$$

# Gradient ascent: Battle of the Sexes

- Symmetric, but not zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} 0,0 & 2,3 \\ 3,2 & 1,1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -4 \cdot \beta + 1 \\ -4 \cdot \alpha + 1 \end{bmatrix}$$

- Stationary at  $(1/4, 1/4)$ .



# Gradient ascent: Battle of the Sexes

- Symmetric, but not zero sum:

$$\begin{array}{c} \text{L} \quad \text{R} \\ \text{T} \left( \begin{array}{cc} 0,0 & 2,3 \\ 3,2 & 1,1 \end{array} \right) \\ \text{B} \end{array}$$

- Gradient:

$$\begin{bmatrix} -4 \cdot \beta + 1 \\ -4 \cdot \alpha + 1 \end{bmatrix}$$

- Stationary at  $(1/4, 1/4)$ .

- Matrix

$$U = \begin{bmatrix} 0 & -4 \\ -4 & 0 \end{bmatrix}$$

has real eigenvalues:

$$\lambda^2 - 16 = 0.$$

# Gradient ascent: Battle of the Sexes

- Symmetric, but not zero sum:

$$\begin{array}{c} \text{L} \quad \text{R} \\ \text{T} \left( \begin{array}{cc} 0,0 & 2,3 \\ 3,2 & 1,1 \end{array} \right) \\ \text{B} \end{array}$$

- Gradient:

$$\begin{bmatrix} -4 \cdot \beta + 1 \\ -4 \cdot \alpha + 1 \end{bmatrix}$$

- Stationary at  $(1/4, 1/4)$ .

- Matrix

$$U = \begin{bmatrix} 0 & -4 \\ -4 & 0 \end{bmatrix}$$

has real eigenvalues:

$\lambda^2 - 16 = 0$ . Saddle point  
inside  $[0, 1]^2$ .

# Gradient ascent: Battle of the Sexes

- Symmetric, but not zero sum:

$$\begin{array}{c} \text{L} \quad \text{R} \\ \text{T} \left( \begin{array}{cc} 0,0 & 2,3 \end{array} \right) \\ \text{B} \left( \begin{array}{cc} 3,2 & 1,1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -4 \cdot \beta + 1 \\ -4 \cdot \alpha + 1 \end{bmatrix}$$


- Stationary at  $(1/4, 1/4)$ .

- Matrix

$$U = \begin{bmatrix} 0 & -4 \\ -4 & 0 \end{bmatrix}$$

has real eigenvalues:

$\lambda^2 - 16 = 0$ . Saddle point  
inside  $[0, 1]^2$ .



# Gradient ascent: Matching pennies

# Gradient ascent: Matching pennies

- Symmetric, zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} 1, -1 & -1, 1 \\ -1, 1 & 1, -1 \end{array} \right) \end{array}$$

# Gradient ascent: Matching pennies

- Symmetric, zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} 1, -1 & -1, 1 \\ -1, 1 & 1, -1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 4 \cdot \beta - 2 \\ -4 \cdot \alpha + 2 \end{bmatrix}$$

# Gradient ascent: Matching pennies

- Symmetric, zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} 1, -1 & -1, 1 \\ -1, 1 & 1, -1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 4 \cdot \beta - 2 \\ -4 \cdot \alpha + 2 \end{bmatrix}$$

- Stationary at  $(1/2, 1/2)$ .

# Gradient ascent: Matching pennies

- Symmetric, zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} 1, -1 & -1, 1 \\ -1, 1 & 1, -1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 4 \cdot \beta - 2 \\ -4 \cdot \alpha + 2 \end{bmatrix}$$

- Stationary at  $(1/2, 1/2)$ .

- Matrix

$$U = \begin{bmatrix} 0 & 4 \\ -4 & 0 \end{bmatrix}$$

has imaginary eigenvalues:

$$\lambda^2 + 16 = 0.$$



# Gradient ascent: Matching pennies

- Symmetric, zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} 1, -1 & -1, 1 \\ -1, 1 & 1, -1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 4 \cdot \beta - 2 \\ -4 \cdot \alpha + 2 \end{bmatrix}$$

- Stationary at  $(1/2, 1/2)$ .

- Matrix

$$U = \begin{bmatrix} 0 & 4 \\ -4 & 0 \end{bmatrix}$$

has imaginary eigenvalues:

$\lambda^2 + 16 = 0$ . Centric point  
inside  $[0, 1]^2$ .

# Gradient ascent: Matching pennies

- Symmetric, zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} 1, -1 & -1, 1 \\ -1, 1 & 1, -1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} 4 \cdot \beta - 2 \\ -4 \cdot \alpha + 2 \end{bmatrix}$$


- Stationary at  $(1/2, 1/2)$ .

- Matrix

$$U = \begin{bmatrix} 0 & 4 \\ -4 & 0 \end{bmatrix}$$

has imaginary eigenvalues:

$\lambda^2 + 16 = 0$ . Centric point  
inside  $[0, 1]^2$ .



# Gradient ascent: other game with centric

# Gradient ascent: other game with centric

- Symmetric, zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} -2, 2 & 1, 1 \\ 3, -3 & -2, 1 \end{array} \right) \end{array}$$

# Gradient ascent: other game with centric

- Symmetric, zero sum:

$$\begin{array}{cc} & \begin{array}{cc} L & R \end{array} \\ \begin{array}{c} T \\ B \end{array} & \left( \begin{array}{cc} -2, 2 & 1, 1 \\ 3, -3 & -2, 1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -8 \cdot \beta + 3 \\ 5 \cdot \alpha - 4 \end{bmatrix}$$

# Gradient ascent: other game with centric

- Symmetric, zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} -2, 2 & 1, 1 \\ 3, -3 & -2, 1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -8 \cdot \beta + 3 \\ 5 \cdot \alpha - 4 \end{bmatrix}$$

- Stationary at  $(4/5, 3/8)$ .

# Gradient ascent: other game with centric

- Symmetric, zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} -2, 2 & 1, 1 \\ 3, -3 & -2, 1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -8 \cdot \beta + 3 \\ 5 \cdot \alpha - 4 \end{bmatrix}$$

- Stationary at  $(4/5, 3/8)$ .

- Matrix

$$U = \begin{bmatrix} 0 & -8 \\ 5 & 0 \end{bmatrix}$$

has imaginary eigenvalues:

$$\lambda^2 + 40 = 0.$$

# Gradient ascent: other game with centric

- Symmetric, zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} -2, 2 & 1, 1 \\ 3, -3 & -2, 1 \end{array} \right) \end{array}$$

- Gradient:

$$\begin{bmatrix} -8 \cdot \beta + 3 \\ 5 \cdot \alpha - 4 \end{bmatrix}$$

- Stationary at  $(4/5, 3/8)$ .

- Matrix

$$U = \begin{bmatrix} 0 & -8 \\ 5 & 0 \end{bmatrix}$$

has imaginary eigenvalues:

$\lambda^2 + 40 = 0$ . Centric point  
inside  $[0, 1]^2$ .



# Gradient ascent: other game with centric

- Symmetric, zero sum:

$$\begin{array}{cc} & \begin{array}{cc} \text{L} & \text{R} \end{array} \\ \begin{array}{c} \text{T} \\ \text{B} \end{array} & \left( \begin{array}{cc} -2, 2 & 1, 1 \\ 3, -3 & -2, 1 \end{array} \right) \end{array}$$

- Gradient:


$$\begin{bmatrix} -8 \cdot \beta + 3 \\ 5 \cdot \alpha - 4 \end{bmatrix}$$

- Stationary at  $(4/5, 3/8)$ .

- Matrix

$$U = \begin{bmatrix} 0 & -8 \\ 5 & 0 \end{bmatrix}$$

has imaginary eigenvalues:  
 $\lambda^2 + 40 = 0$ . Centric point  
inside  $[0, 1]^2$ .



# Convergence of IGA (Singh *et al.*, 2000)

**Proof outline.** There are two main cases:

# Convergence of IGA (Singh *et al.*, 2000)

**Proof outline.** There are two main cases:

1. **There is no stationary point, or the stationary point lies outside  $[0, 1]^2$ .** *Then there is movement everywhere in  $[0, 1]^2$ .*

Since movement is caused by an affine differential map the flow is in one direction, hence gets stuck somewhere at the boundary.

# Convergence of IGA (Singh *et al.*, 2000)

**Proof outline.** There are two main cases:

1. **There is no stationary point, or the stationary point lies outside  $[0, 1]^2$ .** *Then there is movement everywhere in  $[0, 1]^2$ .*

Since movement is caused by an affine differential map the flow is in one direction, hence gets stuck somewhere at the boundary.

2. **There is a stationary point inside  $[0, 1]^2$ .**

# Convergence of IGA (Singh *et al.*, 2000)

**Proof outline.** There are two main cases:

1. **There is no stationary point, or the stationary point lies outside  $[0, 1]^2$ .** *Then there is movement everywhere in  $[0, 1]^2$ .*

Since movement is caused by an affine differential map the flow is in one direction, hence gets stuck somewhere at the boundary.

2. **There is a stationary point inside  $[0, 1]^2$ .**

(a) The stationary point is an

**attractor.** Then it attracts movement which then becomes stationary.

# Convergence of IGA (Singh *et al.*, 2000)

**Proof outline.** There are two main cases:

1. **There is no stationary point, or the stationary point lies outside  $[0, 1]^2$ .** *Then there is movement everywhere in  $[0, 1]^2$ .*

Since movement is caused by an affine differential map the flow is in one direction, hence gets stuck somewhere at the boundary.

2. **There is a stationary point inside  $[0, 1]^2$ .**

(a) The stationary point is an

**attractor**. Then it attracts movement which then becomes stationary.

- (b) The stationary point is a **repellor**. Then it repels movement towards the boundary.

# Convergence of IGA (Singh *et al.*, 2000)

**Proof outline.** There are two main cases:

1. **There is no stationary point, or the stationary point lies outside  $[0, 1]^2$ .** *Then there is movement everywhere in  $[0, 1]^2$ .*

Since movement is caused by an affine differential map the flow is in one direction, hence gets stuck somewhere at the boundary.

2. **There is a stationary point inside  $[0, 1]^2$ .**

(a) The stationary point is an

**attractor**. Then it attracts movement which then becomes stationary.

(b) The stationary point is a **repellor**. Then it repels movement towards the boundary.

(c) Both (2a) and (2b): saddle point.

# Convergence of IGA (Singh *et al.*, 2000)

**Proof outline.** There are two main cases:

1. **There is no stationary point, or the stationary point lies outside  $[0, 1]^2$ .** *Then there is movement everywhere in  $[0, 1]^2$ .*

Since movement is caused by an affine differential map the flow is in one direction, hence gets stuck somewhere at the boundary.

2. **There is a stationary point inside  $[0, 1]^2$ .**

(a) The stationary point is an

**attractor**. Then it attracts movement which then becomes stationary.

- (b) The stationary point is a **repellor**. Then it repels movement towards the boundary.
- (c) Both (2a) and (2b): saddle point.
- (d) **None of the above. Then strategies do not converge.**



# Convergence of IGA (Singh *et al.*, 2000)

**Proof outline.** There are two main cases:

1. **There is no stationary point, or the stationary point lies outside  $[0, 1]^2$ .** *Then there is movement everywhere in  $[0, 1]^2$ .*

Since movement is caused by an affine differential map the flow is in one direction, hence gets stuck somewhere at the boundary.

2. **There is a stationary point inside  $[0, 1]^2$ .**

- (a) The stationary point is an

**attractor.** Then it attracts movement which then becomes stationary.

- (b) The stationary point is a **repellor.** Then it repels movement towards the boundary.
- (c) Both (2a) and (2b): saddle point.
- (d) **None of the above. Then strategies do not converge.**

In three out of four cases, the dynamics ends, hence ends in Nash.

# Convergence of IGA (Singh *et al.*, 2000)

**Proof outline.** There are two main cases:

1. **There is no stationary point, or the stationary point lies outside  $[0, 1]^2$ .** *Then there is movement everywhere in  $[0, 1]^2$ .*

Since movement is caused by an affine differential map the flow is in one direction, hence gets stuck somewhere at the boundary.

2. **There is a stationary point inside  $[0, 1]^2$ .**

- (a) The stationary point is an

**attractor.** Then it attracts movement which then becomes stationary.

- (b) The stationary point is a **repellor.** Then it repels movement towards the boundary.
- (c) Both (2a) and (2b): saddle point.
- (d) **None of the above. Then strategies do not converge.**

In three out of four cases, the dynamics ends, hence ends in Nash.

# Part 3: IGA-WoLF

# IGA-WoLF (Bowling *et al.*, 2001)

Bowling and Veloso modify IGA so as to ensure convergence in Case 2d.  
Idea: Win or Learn Fast

# IGA-WoLF (Bowling *et al.*, 2001)

Bowling and Veloso modify IGA so as to ensure convergence in Case 2d.  
Idea: Win or Learn Fast (**WoLF**).

# IGA-WoLF (Bowling *et al.*, 2001)

Bowling and Veloso modify IGA so as to ensure convergence in Case 2d.  
Idea: Win or Learn Fast (**WoLF**).

To this end, IGA-WoLF uses a variable learning rate:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \begin{bmatrix} l_t^1 \cdot \partial u_1 / \partial \alpha \\ l_t^2 \cdot \partial u_2 / \partial \beta \end{bmatrix}_t$$

# IGA-WoLF (Bowling *et al.*, 2001)

Bowling and Veloso modify IGA so as to ensure convergence in Case 2d.  
Idea: Win or Learn Fast (**WoLF**).

To this end, IGA-WoLF uses a variable learning rate:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \begin{bmatrix} l_t^1 \cdot \partial u_1 / \partial \alpha \\ l_t^2 \cdot \partial u_2 / \partial \beta \end{bmatrix}_t$$

where  $l_t^{\{1,2\}} \in \{l_{\min}, l_{\max}\}$  all positive.

# IGA-WoLF (Bowling *et al.*, 2001)

Bowling and Veloso modify IGA so as to ensure convergence in Case 2d.  
Idea: Win or Learn Fast (**WoLF**).

To this end, IGA-WoLF uses a variable learning rate:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \begin{bmatrix} l_t^1 \cdot \partial u_1 / \partial \alpha \\ l_t^2 \cdot \partial u_2 / \partial \beta \end{bmatrix}_t$$

where  $l_t^{\{1,2\}} \in \{l_{\min}, l_{\max}\}$  all positive.

$$l_t^1 =_{\text{Def}} \begin{cases} l_{\min} & \text{if } u_1(\alpha_t, \beta_t) > u_1(\alpha^e, \beta_t) \\ l_{\max} & \text{otherwise} \end{cases} \quad \begin{array}{l} \text{Winning} \\ \text{Losing} \end{array}$$



# IGA-WoLF (Bowling *et al.*, 2001)

Bowling and Veloso modify IGA so as to ensure convergence in Case 2d.  
Idea: Win or Learn Fast (**WoLF**).

To this end, IGA-WoLF uses a variable learning rate:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \begin{bmatrix} l_t^1 \cdot \partial u_1 / \partial \alpha \\ l_t^2 \cdot \partial u_2 / \partial \beta \end{bmatrix}_t$$

where  $l_t^{\{1,2\}} \in \{l_{\min}, l_{\max}\}$  all positive.

$$l_t^1 =_{\text{Def}} \begin{cases} l_{\min} & \text{if } u_1(\alpha_t, \beta_t) > u_1(\alpha^e, \beta_t) \\ l_{\max} & \text{otherwise} \end{cases} \quad \begin{array}{l} \text{Winning} \\ \text{Losing} \end{array}$$
$$l_t^2 =_{\text{Def}} \begin{cases} l_{\min} & \text{if } u_2(\alpha_t, \beta_t) > u_2(\alpha_t, \beta^e) \\ l_{\max} & \text{otherwise} \end{cases} \quad \begin{array}{l} \text{Winning} \\ \text{Losing} \end{array}$$

# IGA-WoLF (Bowling *et al.*, 2001)

Bowling and Veloso modify IGA so as to ensure convergence in Case 2d.  
Idea: Win or Learn Fast (**WoLF**).

To this end, IGA-WoLF uses a variable learning rate:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \begin{bmatrix} l_t^1 \cdot \partial u_1 / \partial \alpha \\ l_t^2 \cdot \partial u_2 / \partial \beta \end{bmatrix}_t$$

where  $l_t^{\{1,2\}} \in \{l_{\min}, l_{\max}\}$  all positive.

$$l_t^1 =_{\text{Def}} \begin{cases} l_{\min} & \text{if } u_1(\alpha_t, \beta_t) > u_1(\alpha^e, \beta_t) \\ l_{\max} & \text{otherwise} \end{cases} \begin{matrix} \text{Winning} \\ \text{Losing} \end{matrix}$$
$$l_t^2 =_{\text{Def}} \begin{cases} l_{\min} & \text{if } u_2(\alpha_t, \beta_t) > u_2(\alpha_t, \beta^e) \\ l_{\max} & \text{otherwise} \end{cases} \begin{matrix} \text{Winning} \\ \text{Losing} \end{matrix}$$

where  $\alpha^e$  is a row strategy belonging to some NE, chosen by the row player.

# IGA-WoLF (Bowling *et al.*, 2001)

Bowling and Veloso modify IGA so as to ensure convergence in Case 2d.  
Idea: Win or Learn Fast (**WoLF**).

To this end, IGA-WoLF uses a variable learning rate:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \begin{bmatrix} l_t^1 \cdot \partial u_1 / \partial \alpha \\ l_t^2 \cdot \partial u_2 / \partial \beta \end{bmatrix}_t$$

where  $l_t^{\{1,2\}} \in \{l_{\min}, l_{\max}\}$  all positive.

$$l_t^1 =_{\text{Def}} \begin{cases} l_{\min} & \text{if } u_1(\alpha_t, \beta_t) > u_1(\alpha^e, \beta_t) \\ l_{\max} & \text{otherwise} \end{cases} \begin{matrix} \text{Winning} \\ \text{Losing} \end{matrix}$$
$$l_t^2 =_{\text{Def}} \begin{cases} l_{\min} & \text{if } u_2(\alpha_t, \beta_t) > u_2(\alpha_t, \beta^e) \\ l_{\max} & \text{otherwise} \end{cases} \begin{matrix} \text{Winning} \\ \text{Losing} \end{matrix}$$

where  $\alpha^e$  is a row strategy belonging to some NE, chosen by the row player. Similarly for  $\beta^e$  and column player.

# IGA-WoLF (Bowling *et al.*, 2001)

Bowling and Veloso modify IGA so as to ensure convergence in Case 2d.  
Idea: Win or Learn Fast (**WoLF**).

To this end, IGA-WoLF uses a variable learning rate:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix}_{t+1} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}_t + \eta \begin{bmatrix} l_t^1 \cdot \partial u_1 / \partial \alpha \\ l_t^2 \cdot \partial u_2 / \partial \beta \end{bmatrix}_t$$

where  $l_t^{\{1,2\}} \in \{l_{\min}, l_{\max}\}$  all positive.

$$l_t^1 =_{\text{Def}} \begin{cases} l_{\min} & \text{if } u_1(\alpha_t, \beta_t) > u_1(\alpha^e, \beta_t) \\ l_{\max} & \text{otherwise} \end{cases} \quad \begin{array}{l} \text{Winning} \\ \text{Losing} \end{array}$$
$$l_t^2 =_{\text{Def}} \begin{cases} l_{\min} & \text{if } u_2(\alpha_t, \beta_t) > u_2(\alpha_t, \beta^e) \\ l_{\max} & \text{otherwise} \end{cases} \quad \begin{array}{l} \text{Winning} \\ \text{Losing} \end{array}$$

where  $\alpha^e$  is a row strategy belonging to some NE, chosen by the row player. Similarly for  $\beta^e$  and column player. So  $(\alpha^e, \beta^e)$  need not be Nash!

## Case 2d: revolution around Nash equilibrium

**Lemma 1.** *With fixed  $l^1$  and  $l^2$ , the trajectory of the strategy pair  $(\alpha, \beta)$  is an *elliptic orbit* around  $(\alpha^*, \beta^*)$  with axes*

$$\begin{bmatrix} 0 \\ \sqrt{l^2|u|/l^1|u'|} \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

## Case 2d: revolution around Nash equilibrium

**Lemma 1.** *With fixed  $l^1$  and  $l^2$ , the trajectory of the strategy pair  $(\alpha, \beta)$  is an *elliptic orbit* around  $(\alpha^*, \beta^*)$  with axes*

$$\begin{bmatrix} 0 \\ \sqrt{l^2|u|/l^1|u'|} \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Remarks:

## Case 2d: revolution around Nash equilibrium

**Lemma 1.** With fixed  $l^1$  and  $l^2$ , the trajectory of the strategy pair  $(\alpha, \beta)$  is an *elliptic orbit* around  $(\alpha^*, \beta^*)$  with axes

$$\begin{bmatrix} 0 \\ \sqrt{l^2|u|/l^1|u'|} \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Remarks:

- For ellipses with center  $(\alpha^*, \beta^*)$  there are four possibilities, depending on  $u, u'$ , and whether  $\sqrt{l^2|u|/l^1|u'|} > 1$  or  $< 1$ .

## Case 2d: revolution around Nash equilibrium

**Lemma 1.** With fixed  $l^1$  and  $l^2$ , the trajectory of the strategy pair  $(\alpha, \beta)$  is an *elliptic orbit* around  $(\alpha^*, \beta^*)$  with axes

$$\begin{bmatrix} 0 \\ \sqrt{l^2|u|/l^1|u'|} \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Remarks:

■ For ellipses with center  $(\alpha^*, \beta^*)$  there are four possibilities, depending on  $u, u'$ , and whether  $\sqrt{l^2|u|/l^1|u'|} > 1$  or  $< 1$ .

1. Lies flat and axes  $< 1$ .



## Case 2d: revolution around Nash equilibrium

**Lemma 1.** With fixed  $l^1$  and  $l^2$ , the trajectory of the strategy pair  $(\alpha, \beta)$  is an *elliptic orbit* around  $(\alpha^*, \beta^*)$  with axes

$$\begin{bmatrix} 0 \\ \sqrt{l^2|u|/l^1|u'|} \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Remarks:

- For ellipses with center  $(\alpha^*, \beta^*)$  there are four possibilities, depending on  $u, u'$ , and whether  $\sqrt{l^2|u|/l^1|u'|} > 1$  or  $< 1$ .
  1. Lies flat and axes  $< 1$ .
  2. Stands and axes  $< 1$ .

## Case 2d: revolution around Nash equilibrium

**Lemma 1.** With fixed  $l^1$  and  $l^2$ , the trajectory of the strategy pair  $(\alpha, \beta)$  is an *elliptic orbit* around  $(\alpha^*, \beta^*)$  with axes

$$\begin{bmatrix} 0 \\ \sqrt{l^2|u|/l^1|u'|} \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Remarks:

- For ellipses with center  $(\alpha^*, \beta^*)$  there are four possibilities, depending on  $u, u'$ , and whether  $\sqrt{l^2|u|/l^1|u'|} > 1$  or  $< 1$ .
  1. Lies flat and axes  $< 1$ .
  2. Stands and axes  $< 1$ .
  3. Lies flat and axes  $> 1$ .

## Case 2d: revolution around Nash equilibrium

**Lemma 1.** With fixed  $l^1$  and  $l^2$ , the trajectory of the strategy pair  $(\alpha, \beta)$  is an *elliptic orbit* around  $(\alpha^*, \beta^*)$  with axes

$$\begin{bmatrix} 0 \\ \sqrt{l^2|u|/l^1|u'|} \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Remarks:

- For ellipses with center  $(\alpha^*, \beta^*)$  there are four possibilities, depending on  $u, u'$ , and whether  $\sqrt{l^2|u|/l^1|u'|} > 1$  or  $< 1$ .
  1. Lies flat and axes  $< 1$ .
  2. Stands and axes  $< 1$ .
  3. Lies flat and axes  $> 1$ .
  4. Stands and axes  $> 1$ .

## Case 2d: revolution around Nash equilibrium

**Lemma 1.** With fixed  $l^1$  and  $l^2$ , the trajectory of the strategy pair  $(\alpha, \beta)$  is an *elliptic orbit* around  $(\alpha^*, \beta^*)$  with axes

$$\begin{bmatrix} 0 \\ \sqrt{l^2|u|/l^1|u'|} \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Remarks:

- For ellipses with center  $(\alpha^*, \beta^*)$  there are four possibilities, depending on  $u, u'$ , and whether  $\sqrt{l^2|u|/l^1|u'|} > 1$  or  $< 1$ .
  1. Lies flat and axes  $< 1$ .
  2. Stands and axes  $< 1$ .
  3. Lies flat and axes  $> 1$ .
  4. Stands and axes  $> 1$ .
- Bowling *et al.* do not prove this result but refer to Sing *et al.*, who, on their turn refer to a work on differential equations by Reinhard (1987).

## Case 2d: revolution around Nash equilibrium

**Lemma 2.** *A player is “winning” if and only if that player’s strategy is moving away from the center.*

## Case 2d: revolution around Nash equilibrium

**Lemma 2.** *A player is “winning” if and only if that player’s strategy is moving away from the center.*

*Proof.* When play revolves around the center, there can only be one equilibrium.

## Case 2d: revolution around Nash equilibrium

**Lemma 2.** *A player is “winning” if and only if that player’s strategy is moving away from the center.*

*Proof.* When play revolves around the center, there can only be one equilibrium. So  $(\alpha^e, \beta^e)$  is in fact an equilibrium.

## Case 2d: revolution around Nash equilibrium

**Lemma 2.** *A player is “winning” if and only if that player’s strategy is moving away from the center.*

*Proof.* When play revolves around the center, there can only be one equilibrium. So  $(\alpha^e, \beta^e)$  is in fact an equilibrium.

Consider the row player, who wins if and only if

$$u_1(\alpha_t, \beta_t) - u_1(\alpha^e, \beta_t) > 0.$$

Simplifying and using  $\beta u - (r_{12} - r_{22}) = \partial u_1 / \partial \alpha$  yields

$$(\alpha - \alpha^e) \frac{\partial u_1}{\partial \alpha} > 0.$$



## Case 2d: revolution around Nash equilibrium

**Lemma 2.** *A player is “winning” if and only if that player’s strategy is moving away from the center.*

*Proof.* When play revolves around the center, there can only be one equilibrium. So  $(\alpha^e, \beta^e)$  is in fact an equilibrium.

Consider the row player, who wins if and only if

$$u_1(\alpha_t, \beta_t) - u_1(\alpha^e, \beta_t) > 0.$$

Simplifying and using  $\beta u - (r_{12} - r_{22}) = \partial u_1 / \partial \alpha$  yields

$$(\alpha - \alpha^e) \frac{\partial u_1}{\partial \alpha} > 0.$$

Thus, row player “wins” iff either  $\alpha > \alpha^e$  and  $\alpha$  increases, or else  $\alpha < \alpha^e$  and  $\alpha$  decreases. □

## Case 2d: revolution around Nash equilibrium

**Lemma 2.** *A player is “winning” if and only if that player’s strategy is moving away from the center.*

*Proof.* When play revolves around the center, there can only be one equilibrium. So  $(\alpha^e, \beta^e)$  is in fact an equilibrium.

Consider the row player, who wins if and only if

$$u_1(\alpha_t, \beta_t) - u_1(\alpha^e, \beta_t) > 0.$$

Simplifying and using  $\beta u - (r_{12} - r_{22}) = \partial u_1 / \partial \alpha$  yields

$$(\alpha - \alpha^e) \frac{\partial u_1}{\partial \alpha} > 0.$$

Thus, row player “wins” iff either  $\alpha > \alpha^e$  and  $\alpha$  increases, or else  $\alpha < \alpha^e$  and  $\alpha$  decreases. □

**Corollary.** *The learning rate is constant throughout any one quadrant.*

## Case 2d: revolution around Nash equilibrium

**Lemma 3.** *Let  $C$  be the center. For every initial strategy pair  $(\alpha, \beta)$  that is sufficiently close to  $C$ , the  $l_{\min} / l_{\max}$  dynamics will bring that pair to  $C$ .*

## Case 2d: revolution around Nash equilibrium

**Lemma 3.** *Let  $C$  be the center. For every initial strategy pair  $(\alpha, \beta)$  that is sufficiently close to  $C$ , the  $l_{\min} / l_{\max}$  dynamics will bring that pair to  $C$ .*

*Proof.*

Let  $(\alpha, \beta)$  be a strategy pair.  
According to Lemma 1, its trajectory forms an ellipse with center  $C$ .

## Case 2d: revolution around Nash equilibrium

**Lemma 3.** *Let  $C$  be the center. For every initial strategy pair  $(\alpha, \beta)$  that is sufficiently close to  $C$ , the  $l_{\min} / l_{\max}$  dynamics will bring that pair to  $C$ .*

*Proof.*

Let  $(\alpha, \beta)$  be a strategy pair. According to Lemma 1, its trajectory forms an ellipse with center  $C$ .

If  $(\alpha, \beta)$  is sufficiently close to  $C$ , this ellipse will entirely be in  $[0, 1]^2$  and its trajectory will not be disrupted.

## Case 2d: revolution around Nash equilibrium

**Lemma 3.** *Let  $C$  be the center. For every initial strategy pair  $(\alpha, \beta)$  that is sufficiently close to  $C$ , the  $l_{\min} / l_{\max}$  dynamics will bring that pair to  $C$ .*

*Proof.*

Let  $(\alpha, \beta)$  be a strategy pair. According to Lemma 1, its trajectory forms an ellipse with center  $C$ .

If  $(\alpha, \beta)$  is sufficiently close to  $C$ , this ellipse will entirely be in  $[0, 1]^2$  and its trajectory will not be disrupted.

There are two cases.

## Case 2d: revolution around Nash equilibrium

**Lemma 3.** *Let  $C$  be the center. For every initial strategy pair  $(\alpha, \beta)$  that is sufficiently close to  $C$ , the  $l_{\min} / l_{\max}$  dynamics will bring that pair to  $C$ .*

*Proof.*

Let  $(\alpha, \beta)$  be a strategy pair. According to Lemma 1, its trajectory forms an ellipse with center  $C$ .

If  $(\alpha, \beta)$  is sufficiently close to  $C$ , this ellipse will entirely be in  $[0, 1]^2$  and its trajectory will not be disrupted.

There are two cases.

1. *Strategy pair moves clockwise.*

## Case 2d: revolution around Nash equilibrium

**Lemma 3.** *Let  $C$  be the center. For every initial strategy pair  $(\alpha, \beta)$  that is sufficiently close to  $C$ , the  $l_{\min} / l_{\max}$  dynamics will bring that pair to  $C$ .*

*Proof.*

Let  $(\alpha, \beta)$  be a strategy pair. According to Lemma 1, its trajectory forms an ellipse with center  $C$ .

If  $(\alpha, \beta)$  is sufficiently close to  $C$ , this ellipse will entirely be in  $[0, 1]^2$  and its trajectory will not be disrupted.

There are two cases.

1. *Strategy pair moves clockwise.*

- (a) We will thus have to ensure that learning parameters are set such that ellipse that forms the trajectory “stands” when  $(\alpha, \beta)$  is in  $Q1$  and  $Q3$ .



## Case 2d: revolution around Nash equilibrium

**Lemma 3.** *Let  $C$  be the center. For every initial strategy pair  $(\alpha, \beta)$  that is sufficiently close to  $C$ , the  $l_{\min} / l_{\max}$  dynamics will bring that pair to  $C$ .*

*Proof.*

Let  $(\alpha, \beta)$  be a strategy pair. According to Lemma 1, its trajectory forms an ellipse with center  $C$ .

If  $(\alpha, \beta)$  is sufficiently close to  $C$ , this ellipse will entirely be in  $[0, 1]^2$  and its trajectory will not be disrupted.

There are two cases.

1. *Strategy pair moves clockwise.*

- (a) We will thus have to ensure that learning parameters are set such that ellipse that forms the trajectory “stands” when  $(\alpha, \beta)$  is in  $Q1$  and  $Q3$ .
- (b) Similarly, we will have to ensure that learning parameters are such that the ellipse “lies flat” when  $(\alpha, \beta)$  is in  $Q2$  and  $Q4$ .

## Case 2d: revolution around Nash equilibrium

**Lemma 3.** *Let  $C$  be the center. For every initial strategy pair  $(\alpha, \beta)$  that is sufficiently close to  $C$ , the  $l_{\min} / l_{\max}$  dynamics will bring that pair to  $C$ .*

*Proof.*

Let  $(\alpha, \beta)$  be a strategy pair. According to Lemma 1, its trajectory forms an ellipse with center  $C$ .

If  $(\alpha, \beta)$  is sufficiently close to  $C$ , this ellipse will entirely be in  $[0, 1]^2$  and its trajectory will not be disrupted.

There are two cases.

1. *Strategy pair moves clockwise.*

- (a) We will thus have to ensure that learning parameters are set such that ellipse that forms the trajectory “stands” when  $(\alpha, \beta)$  is in  $Q1$  and  $Q3$ .
- (b) Similarly, we will have to ensure that learning parameters are such that the ellipse “lies flat” when  $(\alpha, \beta)$  is in  $Q2$  and  $Q4$ .

2. *Strategy pair moves counter-clockwise.* Similar reasoning.

## Case 2d: revolution around Nash equilibrium

**Lemma 3.** *Let  $C$  be the center. For every initial strategy pair  $(\alpha, \beta)$  that is sufficiently close to  $C$ , the  $l_{\min} / l_{\max}$  dynamics will bring that pair to  $C$ .*

*Proof.*

Let  $(\alpha, \beta)$  be a strategy pair. According to Lemma 1, its trajectory forms an ellipse with center  $C$ .

If  $(\alpha, \beta)$  is sufficiently close to  $C$ , this ellipse will entirely be in  $[0, 1]^2$  and its trajectory will not be disrupted.

There are two cases.


1. *Strategy pair moves clockwise.*

- (a) We will thus have to ensure that learning parameters are set such that ellipse that forms the trajectory “stands” when  $(\alpha, \beta)$  is in  $Q1$  and  $Q3$ .
- (b) Similarly, we will have to ensure that learning parameters are such that the ellipse “lies flat” when  $(\alpha, \beta)$  is in  $Q2$  and  $Q4$ .


2. *Strategy pair moves counter-clockwise.* Similar reasoning.

(First a suggestive picture, then the rest of the proof.)


# Trajectory in different quadrants (clockwise)




# Trajectory in different quadrants (clockwise)



# Trajectory in different quadrants (clockwise)



# Compound trajectory



# Trajectory in different quadrants (clockwise)

**Claim.** *The learning parameters  $l_{\min}$  /  $l_{\max}$  alternate in such a way that the ellipse that forms the trajectory in clockwise movement “lies flat” when  $(\alpha, \beta)$  is in Q1 and Q3 of the ellipse and “stands” otherwise.*



# Trajectory in different quadrants (clockwise)

**Claim.** *The learning parameters  $l_{\min}$  /  $l_{\max}$  alternate in such a way that the ellipse that forms the trajectory in clockwise movement “lies flat” when  $(\alpha, \beta)$  is in Q1 and Q3 of the ellipse and “stands” otherwise.*

**Proof.**

# Trajectory in different quadrants (clockwise)

**Claim.** *The learning parameters  $l_{\min}$  /  $l_{\max}$  alternate in such a way that the ellipse that forms the trajectory in clockwise movement “lies flat” when  $(\alpha, \beta)$  is in Q1 and Q3 of the ellipse and “stands” otherwise.*

**Proof.**

Suppose movement is clockwise.

# Trajectory in different quadrants (clockwise)

**Claim.** *The learning parameters  $l_{\min}$  /  $l_{\max}$  alternate in such a way that the ellipse that forms the trajectory in clockwise movement “lies flat” when  $(\alpha, \beta)$  is in Q1 and Q3 of the ellipse and “stands” otherwise.*

**Proof.**

Suppose movement is clockwise.

1. Suppose  $(\alpha, \beta)$  is in Q1 (upper right) of the ellipse. Then row “wins” and col “loses”. By Lemma 2 horizontal velocity  $<$  vertical velocity, and the ellipse “stands”.

# Trajectory in different quadrants (clockwise)

**Claim.** *The learning parameters  $l_{\min}$  /  $l_{\max}$  alternate in such a way that the ellipse that forms the trajectory in clockwise movement “lies flat” when  $(\alpha, \beta)$  is in Q1 and Q3 of the ellipse and “stands” otherwise.*

**Proof.**

Suppose movement is clockwise.

1. Suppose  $(\alpha, \beta)$  is in Q1 (upper right) of the ellipse. Then row “wins” and col “loses”. By Lemma 2 horizontal velocity  $<$  vertical velocity, and the ellipse “stands”.
2. Suppose  $(\alpha, \beta)$  is in Q2 (lower right) of the ellipse. Then row “loses” and col “wins”. By Lemma 2 horizontal velocity  $>$  vertical velocity, and the ellipse “lies flat”.

# Trajectory in different quadrants (clockwise)

**Claim.** *The learning parameters  $l_{\min}$  /  $l_{\max}$  alternate in such a way that the ellipse that forms the trajectory in clockwise movement “lies flat” when  $(\alpha, \beta)$  is in Q1 and Q3 of the ellipse and “stands” otherwise.*

**Proof.**

Suppose movement is clockwise.

1. Suppose  $(\alpha, \beta)$  is in Q1 (upper right) of the ellipse. Then row “wins” and col “loses”. By Lemma 2 horizontal velocity  $<$  vertical velocity, and the ellipse “stands”.
2. Suppose  $(\alpha, \beta)$  is in Q2 (lower right) of the ellipse. Then row “loses” and col “wins”. By Lemma 2 horizontal velocity  $>$  vertical velocity, and the ellipse “lies flat”.

Clearly, the reasoning is similar when

- the strategy pair  $(\alpha, \beta)$  is in the other two quadrants

# Trajectory in different quadrants (clockwise)

**Claim.** *The learning parameters  $l_{\min}$  /  $l_{\max}$  alternate in such a way that the ellipse that forms the trajectory in clockwise movement “lies flat” when  $(\alpha, \beta)$  is in Q1 and Q3 of the ellipse and “stands” otherwise.*

**Proof.**

Suppose movement is clockwise.

1. Suppose  $(\alpha, \beta)$  is in Q1 (upper right) of the ellipse. Then row “wins” and col “loses”. By Lemma 2 horizontal velocity  $<$  vertical velocity, and the ellipse “stands”.
2. Suppose  $(\alpha, \beta)$  is in Q2 (lower right) of the ellipse. Then row “loses” and col “wins”. By Lemma 2 horizontal velocity  $>$  vertical velocity, and the ellipse “lies flat”.

Clearly, the reasoning is similar when

- the strategy pair  $(\alpha, \beta)$  is in the other two quadrants, or
- when movement is counter-clockwise.



# Part 4: Another solution

# Why not utilise Singh *et al.*'s result on emp. frequencies



# Why not utilise Singh *et al.*'s result on emp. frequencies

- **Theorem** (Singh, Kearns and Mansour, 2000) *If players follow IGA, where  $\eta \rightarrow 0$ , then their strategies will converge to a Nash equilibrium. If not, then at least their average payoffs will converge to the expected payoffs of a Nash equilibrium.*

# Why not utilise Singh *et al.*'s result on emp. frequencies

- **Theorem** (Singh, Kearns and Mansour, 2000) *If players follow IGA, where  $\eta \rightarrow 0$ , then their strategies will converge to a Nash equilibrium. If not, then at least their average payoffs will converge to the expected payoffs of a Nash equilibrium.*
- Idea: use average payoffs to correct the gradient.

# Why not utilise Singh *et al.*'s result on emp. frequencies


- **Theorem** (Singh, Kearns and Mansour, 2000) *If players follow IGA, where  $\eta \rightarrow 0$ , then their strategies will converge to a Nash equilibrium. If not, then at least their average payoffs will converge to the expected payoffs of a Nash equilibrium.*
- Idea: use average payoffs to correct the gradient.
- So gradient points slight more in direction of average payoffs.

# Why not utilise Singh *et al.*'s result on emp. frequencies

- **Theorem** (Singh, Kearns and Mansour, 2000) *If players follow IGA, where  $\eta \rightarrow 0$ , then their strategies will converge to a Nash equilibrium. If not, then at least their average payoffs will converge to the expected payoffs of a Nash equilibrium.*
- Idea: use average payoffs to correct the gradient.
- So gradient points slight more in direction of average payoffs.
- At least works empirically :-)

# Why not utilise Singh *et al.*'s result on emp. frequencies

- **Theorem** (Singh, Kearns and Mansour, 2000) *If players follow IGA, where  $\eta \rightarrow 0$ , then their strategies will converge to a Nash equilibrium. If not, then at least their average payoffs will converge to the expected payoffs of a Nash equilibrium.*
- Idea: use average payoffs to correct the gradient.
- So gradient points slight more in direction of average payoffs.
- At least works empirically :-)



# Literature

# Literature

- Original work on gradient ascent in general-sum games:

## ■ Original work on gradient ascent in general-sum games:

Singh, Kearns, and Mansour (2000). “Nash convergence of gradient ascent in general-sum games”. In: *Proc. of the Sixteenth Conf. on Uncertainty in Artificial Intelligence* (pp. 541-548).



# Literature

- Original work on gradient ascent in general-sum games:

Singh, Kearns, and Mansour (2000). “Nash convergence of gradient ascent in general-sum games”. In: *Proc. of the Sixteenth Conf. on Uncertainty in Artificial Intelligence* (pp. 541-548).

- Today’s presentation was mainly based on this conference publication:

# Literature

- Original work on gradient ascent in general-sum games:

Singh, Kearns, and Mansour (2000). “Nash convergence of gradient ascent in general-sum games”. In: *Proc. of the Sixteenth Conf. on Uncertainty in Artificial Intelligence* (pp. 541-548).

- Today’s presentation was mainly based on this conference publication:

Bowling and Veloso (2001). “Convergence of gradient ascent with a Variable Learning Rate”. In *Proc. of the Eighteenth Int. Conf. on Machine Learning (ICML)*, pp. 27-34, June 2001.

# Literature

- Original work on gradient ascent in general-sum games:

Singh, Kearns, and Mansour (2000). “Nash convergence of gradient ascent in general-sum games”. In: *Proc. of the Sixteenth Conf. on Uncertainty in Artificial Intelligence* (pp. 541-548).

- Today’s presentation was mainly based on this conference publication:

Bowling and Veloso (2001). “Convergence of gradient ascent with a Variable Learning Rate”. In *Proc. of the Eighteenth Int. Conf. on Machine Learning (ICML)*, pp. 27-34, June 2001.

Conference publication was elaborated, and published as a journal article:

# Literature

- Original work on gradient ascent in general-sum games:

Singh, Kearns, and Mansour (2000). “Nash convergence of gradient ascent in general-sum games”. In: *Proc. of the Sixteenth Conf. on Uncertainty in Artificial Intelligence* (pp. 541-548).

- Today’s presentation was mainly based on this conference publication:

Bowling and Veloso (2001). “Convergence of gradient ascent with a Variable Learning Rate”. In *Proc. of the Eighteenth Int. Conf. on Machine Learning (ICML)*, pp. 27-34, June 2001.

Conference publication was elaborated, and published as a journal article:

Bowling and Veloso (2002). “Multiagent Learning Using a Variable Learning Rate”. In: *Artificial Intelligence* **136**, pp. 215-250, 2002.

# What next?

# What next?

- With fictitious play, or gradient ascent, opponents are modelled by a single mixed strategy.

# What next?

- With fictitious play, or gradient ascent, opponents are modelled by a **single mixed strategy**.
- With Bayesian play, opponents are modelled by a **probability distribution over all opponent strategies**

$$\Delta \left[ \prod_{j \neq i} \Delta(X_j)^H \right].$$

- $\Delta(A)$  denotes the set of all probability distributions over  $A$ .
- $B^A$  denotes the set of all functions from  $A$  to  $B$ .
- $\prod_{j \neq i} A_j$  denotes the Cartesian product of  $\{A_j\}_{j \neq i}$ . In case of a finite product, this can be written as

$$\prod_{j \neq i} A_j = A_1 \times A_2 \times \cdots \times A_{i-1} \times A_{i+1} \times \cdots \times A_n.$$