

Fonctions f	Primitives F , avec $C \in \mathbb{R}$
Soit $\alpha \in \mathbb{R}$, $f : x \in D \mapsto x^\alpha$	$\begin{cases} F : x \in D \mapsto \frac{1}{\alpha+1} x^{\alpha+1} & \text{si } \alpha \neq -1 \\ F : x \in D \mapsto \ln(x) + C & \text{si } \alpha = -1 \end{cases}$
Soit $z \in \mathbb{C}^*$, $f : x \in \mathbb{R} \mapsto e^{zx}$	$F : x \in \mathbb{R} \mapsto \frac{1}{z} e^{zx} + C$
Soient $\alpha, \beta \in \mathbb{R}_+^*$, $f : x \in \mathbb{R}_+^* \mapsto \cos(\alpha x) \cos(\beta x)$	$\begin{cases} F : x \in \mathbb{R} \mapsto \frac{1}{2} \left[\frac{1}{\alpha+\beta} \sin((\alpha+\beta)x) + \frac{1}{\alpha-\beta} \sin((\alpha-\beta)x) \right] + C & \text{si } \alpha \neq \beta \\ F : x \in \mathbb{R} \mapsto \frac{1}{2} \left[\frac{1}{2\alpha} \sin(2\alpha x) + x \right] + C & \text{si } \alpha = \beta \end{cases}$
$f : x \in]-1, 1[\mapsto \frac{1}{\sqrt{1-x^2}}$	$F : x \in]-1, 1[\mapsto \arcsin(x) + C$
$f : x \in \mathbb{R} \mapsto \frac{1}{1+x^2}$	$F : x \in \mathbb{R} \mapsto \arctan(x) + C$
$f : x \in \mathbb{R} \setminus \{-1, 1\} \mapsto \frac{1}{1-x^2}$	$F : x \in \mathbb{R} \setminus \{-1, 1\} \mapsto \ln \left \frac{1+x}{1-x} \right + C$
Soient $\alpha, \beta \in \mathbb{R}^*$, $f : x \in \mathbb{R} \mapsto e^{\alpha x} \cos(\beta x)$	$F : x \in \mathbb{R} \mapsto \frac{e^{\alpha x}}{\alpha^2 + \beta^2} (\alpha \cos(\beta x) + \beta \sin(\beta x)) + C$