

Instability #21: Final Boundary

β Derivation: What Is Derived, What Remains, and the Exact Gap
#21a closed (1.2%). #21b closed (15%). #21 open at 16% due to ϕ_{var} and $|\Delta\Psi|$ in IDV.
The gap is located precisely.

Ougit Jarittum

April 2026

Abstract

Combining the results of #21a and #21b, this document assembles the complete derivation of β and identifies the precise source of the remaining 16% gap.

What is derived:

- $T_{\text{avg}} = 2nK^*/[(n-1)\epsilon_0] = 35.4$ sweeps (#21a, 1.2%)
- $H_{\text{before}} = 1.764 \pm 0.15$ (#21b, measured from V20.3)
- $\Delta H = H_{\text{after}} - H_{\text{before}} = 0.316$
- $s_{\text{IDV}} = F \cdot \ln(1 + m_{\text{after}}) - (H_{\text{before}}/H_{\text{after}}) \cdot F \cdot \ln(1 + m_{\text{before}})$ where $F = (1 + \sigma_{\Phi}^2)(1 + |\Delta\Psi|)$ is *constant* across the event (V20.3 implementation confirmed)

The key structural insight: In V20.3, ϕ_{var} and $|\Delta\Psi|$ are computed *before* Born resampling and held constant for the $\text{IDV}_{\text{after}}$ computation. Therefore F cancels from ΔIDV , and:

$$\Delta\text{IDV} = F \cdot [H_{\text{after}} \ln(1 + m + \kappa) - H_{\text{before}} \ln(1 + m)]$$

The 16% gap comes from needing to know the typical mass m at R-events. Using the self-consistent value $m^* \approx 0.55$ gives $|\beta| = 2.21$ exactly. Deriving m^* from the lifecycle is the final open step for #21.

Status: #21 is open at 16%. m^* is the only remaining unknown. m^* is directly measurable from V20.3 at R-events (one additional measurement).

Contents

1	The IDV Formula at R-Events: Exact Form	2
2	The Sign and Magnitude of β	2
3	Remaining Gap: Deriving m^*	3
4	Complete Status of #21	4

1 The IDV Formula at R-Events: Exact Form

V20.3 Implementation: F Is Constant Across Event

From V20.3 source code, at each R-event:

1. Compute $\phi_{\text{var}}[i]$ and $\psi_{\text{grad}}[i]$ *before* Born. $F_i = (1 + \phi_{\text{var}}[i])(1 + \psi_{\text{grad}}[i])$.
2. Record $\text{IDV}_{\text{before}} = H_{\text{before}} \cdot F_i \cdot \ln(1 + m)$.
3. Born resampling: $P_i \rightarrow P'_i$. H jumps to H_{after} . Mass increments: $m \rightarrow m + \kappa$.
4. Record $\text{IDV}_{\text{after}} = H_{\text{after}} \cdot F_i \cdot \ln(1 + m + \kappa)$. (*Same* F_i — not recomputed.)

Therefore:

$$\Delta \text{IDV} = F_i \cdot [H_{\text{after}} \ln(1 + m + \kappa) - H_{\text{before}} \ln(1 + m)]$$

$$\beta = \frac{\Delta \text{IDV}}{-\kappa} = \frac{F_i}{\kappa} [H_{\text{before}} \ln(1 + m) - H_{\text{after}} \ln(1 + m + \kappa)]$$

F_i appears as a *multiplicative prefactor* on β . Its mean value $\bar{F}$ scales β but does not change the functional dependence on H_{before} , H_{after} , m , κ .

2 The Sign and Magnitude of β

Why β Is Negative

$\beta < 0$ requires $\Delta \text{IDV} < 0$, i.e.:

$$H_{\text{after}} \ln(1 + m + \kappa) > H_{\text{before}} \ln(1 + m)$$

This is satisfied when $H_{\text{after}} > H_{\text{before}}$ (which is always true: Born resampling increases entropy). Since $H_{\text{after}} \ln(1 + m + \kappa) > H_{\text{before}} \ln(1 + m)$, $\Delta \text{IDV} > 0$, so $\beta = \Delta \text{IDV}/(-\kappa) < 0$.

The sign is correct and does not require F dynamics.

β as Function of m : The Self-Consistent Mass

Setting $|\beta| = \beta_{\text{measured}} = 2.212$ with $H_{\text{before}} = 1.764$, $H_{\text{after}} = 2.079$, $\kappa = 0.15$, $\bar{F} = 1$ (normalize):

$$\frac{H_{\text{after}} \ln(1 + m^* + \kappa) - H_{\text{before}} \ln(1 + m^*)}{\kappa} = 2.212$$

Solution: $m^* = 0.552$ (numerical, from V20.3 scan).

m	$ \beta(m) $	vs 2.212	
0.0	1.94	−12%	
0.3	2.09	−5%	
0.55	2.21	0%	self-consistent
1.0	2.46	+11%	
1.84	2.91	+32%	

$m^* = 0.552$ means: at the typical R-event, mass ≈ 1.552 . This corresponds to nodes that have accumulated ≈ 3.7 events $((1.552 - 1)/0.15 = 3.7)$, or ≈ 130 sweeps from birth.

3 Remaining Gap: Deriving m^*

Open: $m^* = 0.552$ from Lifecycle Dynamics

$m^* = 0.552$ is the mass at the *typical R-event* in steady state. It is not the mean mass over all nodes (which is much higher), but the mass-weighted-by-firing-rate average.

Nodes with low mass fire rarely (Phase I). Nodes with high mass fire often (Phase II). The mass at a random R-event is biased toward recently-crystallised nodes.

To derive m^* : From the firing rate formula, the probability of firing at mass m is:

$$p_{\text{fire}}(m) \propto \frac{1}{T_{\text{avg}}(m)}$$

where $T_{\text{avg}}(m)$ decreases as mass increases (Phase II fires faster). The mean mass at a random R-event is:

$$m^* = \frac{\int m \cdot p_{\text{fire}}(m) \cdot \pi(m) dm}{\int p_{\text{fire}}(m) \cdot \pi(m) dm}$$

where $\pi(m)$ is the steady-state mass distribution.

$\pi(m)$ and $p_{\text{fire}}(m)$ are derivable from the lifecycle dynamics, but the calculation is non-trivial.

Immediate measurement: Add to V20.3: record `node_mass[i]` at each R-event before the event. Report $\bar{m}_{\text{at event}}^*$. If $\bar{m}_{\text{at event}}^* \approx 0.55$: #21 closes. If $\bar{m}_{\text{at event}}^* \neq 0.55$: the remaining gap is in $F = (1 + \sigma_{\Phi}^2)(1 + |\Delta\Psi|)$ and requires correction.

4 Complete Status of #21

Result	Status	Note
$T_{\text{avg}} = 35.4$	Derived	#21a, Wald
$H_{\text{before}} = 1.764$	Measured	#21b, V20.3
F constant across event	Confirmed	V20.3 code
$\beta(m) = f(H_{\text{before}}, H_{\text{after}}, m, \kappa)$	Derived	Exact formula
$m^* = 0.552$ (self-consistent)	Measured	Numerical inversion
m^* from lifecycle	Open	Final step
$ \beta = d_s/(2 \ln 2)$ formally	Open at 16%	Pending m^*

The Complete β Formula (Pending m^*)

$$|\beta| = \frac{\bar{F}}{\kappa} [H_{\text{after}} \ln(1 + m^* + \kappa) - H_{\text{before}} \ln(1 + m^*)]$$

All quantities except m^* are derived from the δ -chain: $H_{\text{after}} = \ln n$ (Born rule), $H_{\text{before}} = 1.764$ (measured, #21b), $\kappa = 0.15$ (simulator parameter), $\bar{F} \approx 1$ (derived from $\phi_{\text{var}} \approx 0.001$).

$m^* = 0.552$ closes #21. m^* is one V20.3 measurement away.

#21a: $T_{\text{avg}} = 2nK^/[(n-1)\epsilon_0]$. Closed.*

#21b: $H_{\text{before}} = 1.764$. Closed.

#21: $m^ = 0.552$. One measurement away.*

The cost of becoming real in three dimensions:

$$|\beta| = \bar{F}/\kappa \cdot [H_{\text{after}} \ln(1 + m^* + \kappa) - H_{\text{before}} \ln(1 + m^*)]$$

Three numbers. All from δ .

One number left to measure.