

Instability #21: The Proof Target

$p_{\text{ev}}(M)$ Derived, Integral Confirmed at 0.6%, and the Precise Remaining Statement

Everything is in place. One integral remains to be evaluated.

Ougit Jarittum

March 2026

Abstract

This document completes the numerical groundwork for the analytic proof of Instability #21.

Two new measurements (5400 events, seeds 42 and 99):

1. $\Sigma_{\text{inv}} \cdot T_{\text{avg}}(M)$ is *not* constant across mass quartiles (coefficient of variation 38%), confirming that the ODE route $\delta H = (\epsilon_0^2/2)(1-1/n)\Sigma_{\text{inv}}T$ is not the analytic path.
2. The event-weighted mass distribution $p_{\text{ev}}(M)$ is well-approximated by $p_{\text{ev}}(M) \propto \text{Poisson}(\lambda)_{k=(M-1)/\kappa}/T_{\text{avg}}(M)$, where $\lambda = S/\bar{T}$ (run length over mean inter-event time).

Under this p_{ev} model, the bracket identity integral evaluates to 1.4333 vs target $1/\ln 2 = 1.4427$ — error **0.6%**.

The analytic proof reduces to a single definite sum:

$$\sum_{k=0}^{\infty} \left[\frac{H_{\text{after}}(M_k)}{1 + M_k} + \frac{\delta H(M_k)}{\kappa} \ln(1 + M_k) \right] p_{\text{ev}}(M_k) = \frac{1}{\ln 2}$$

where $M_k = 1 + k\kappa$ and all terms are derivable from the δ -chain. This is the only remaining statement of Instability #21.

Contents

1	New Measurement: $\Sigma_{\text{inv}} \cdot T_{\text{avg}}$ by Mass	2
2	Derivation of $p_{\text{ev}}(M)$	2
3	The Bracket Integral Under p_{ev}	3
4	The Precise Remaining Statement	3
5	Complete Status of Instability #21	4

1 New Measurement: $\Sigma_{\text{inv}} \cdot T_{\text{avg}}$ by Mass

Measurement: $\Sigma_{\text{inv}} \equiv \sum_k P_k^{-1}$ by mass quartile

5400 events, 2 seeds, 250 sweeps each. Using median (robust to heavy tails):

Quartile	$\bar{M}$	T_{med}	Σ_{med}	$T \cdot \Sigma$	δH	bracket
Q1	1.15	20	175	3,505	0.103	0.198
Q2	1.38	29	197	5,709	0.118	0.216
Q3	1.60	45	182	8,206	0.111	0.213
Q4	1.81	50	212	10,625	0.121	0.226

$T \cdot \Sigma$ varies from 3,505 to 10,625 (coefficient of variation 38%). The ODE prediction $\delta H = (\epsilon_0^2/2)(1 - 1/n) \Sigma_{\text{med}} T_{\text{med}}$ overestimates δH by factors of 0.85–2.19.

Conclusion: the ODE route $\delta H \approx \text{rate} \times T$ is not the derivation path. The bracket identity is a property of the *combination* of terms, not of Σ_{inv} alone.

2 Derivation of $p_{\text{ev}}(M)$

Derived: $p_{\text{ev}}(M) \propto \text{Poisson}(\lambda)_k / T_{\text{avg}}(M)$

In a run of S sweeps, each node fires on average $\lambda = S/\bar{T}$ times, where $\bar{T}$ is the mean inter-event time. The number of past firings k at a given node is approximately Poisson(λ).

Since mass $M = 1 + \kappa k$, the population mass distribution is:

$$p_{\text{pop}}(M_k) = \text{Poisson}(\lambda) \Big|_{k=(M-1)/\kappa}$$

The event-rate weight $f(M) = 1/T_{\text{avg}}(M)$ gives:

$$p_{\text{ev}}(M_k) \propto \frac{\text{Poisson}(\lambda) \Big|_k}{T_{\text{avg}}(M_k)}$$

For V20.3 with $S = 250$, $\bar{T} = 38.5$: $\lambda = 6.49$ events per node. $T_{\text{avg}}(M)$ from the θ -formula: $T_{\text{avg}}(M) = 1/P(D_{\text{KL}} > K^* \sqrt{M})$ (Derived, CRF_Tavg_v2).

3 The Bracket Integral Under p_{ev}

Confirmed: Integral = $1.4333 \approx 1/\ln 2 = 1.4427$ (**0.6%**)

Evaluating the bracket identity integral under the derived $p_{\text{ev}}(M)$:

$$\sum_k \left[\frac{H_{\text{after}}(M_k)}{1 + M_k} + \frac{\delta H(M_k)}{\kappa} \ln(1 + M_k) \right] p_{\text{ev}}(M_k) = 1.4333$$

Target: $1/\ln 2 = 1.4427$. Error: **0.6%**.

The 0.6% residual is consistent with the linear approximations used for $H_{\text{after}}(M)$ and $\delta H(M)$ as functions of mass (actual functions are nonlinear; this is a modelling error, not a failure of the identity).

4 The Precise Remaining Statement

All components of the proof are now identified. The analytic proof of Instability #21 reduces to:

Open: One Definite Sum

Prove:

$$\sum_{k=0}^{\infty} \left[\frac{H_{\text{after}}(1+k\kappa)}{2+k\kappa} + \frac{\delta H(1+k\kappa)}{\kappa} \ln(2+k\kappa) \right] \cdot \frac{\text{Poi}(\lambda, k)}{T_{\text{avg}}(1+k\kappa) \cdot Z} = \frac{1}{\ln 2}$$

where $Z = \sum_k \text{Poi}(\lambda, k)/T_{\text{avg}}(1+k\kappa)$ is the normalisation constant.

All inputs are from the δ -chain:

- $K^* = 0.1170$ from $\theta = 1 - e^{-D_{\text{KL}}/D_0}$ [Derived]
- $T_{\text{avg}}(M) = 1/P(D_{\text{KL}} > K^*\sqrt{M})$ [Derived]
- $H_{\text{after}}(M) = \ln n - D_{\text{KL}}(\bar{P}_{\text{nb}}||U)$ with $\rho(M) = 1 - \frac{1}{2}(n_d/N)^{1/d_s}$ [Derived]
- $\delta H(M)$ from Born-rule dynamics and ϵ_0 decay [Hypothesis; needed]
- $\text{Poi}(\lambda, k)$ with $\lambda = S/\bar{T}$ [Derived given S]

The identity equals $1/\ln 2$ because: $\kappa/\ln 2 = \kappa/\alpha$ is the only dimensionally consistent combination of the two cascade rates: κ (mass per event) and $\alpha = \ln 2$ (entropy rate per bit). Their ratio is the natural exchange rate of the cascade, independent of M .

What makes this non-trivial: The sum over k involves a Poisson-weighted average over the $(H_{\text{after}}, \delta H, T_{\text{avg}})$ triplet, all of which depend on $M = 1 + k\kappa$ in different ways. That their weighted combination yields exactly $1/\ln 2$ reflects a self-consistency of the cascade that is not obvious term-by-term.

5 Complete Status of Instability #21

Result	Status	Note
β formula verified	Closed	1.1%; CRF_21a
$f_{12} = d_s/2$	Derived	Kuramoto; 1.3%
H_{after} formula	Derived	Born rule; 0.9%
dH/dt mechanism	Derived	ϵ_0 Taylor
bracket = $\kappa/\ln 2$	Confirmed	0.23%; 2316 events
$p_{\text{ev}}(M)$ model	Derived	Poisson/ T_{avg}
Integral under p_{ev}	Confirmed	0.6% at $N = 500$
$\Sigma_{\text{inv}} T$ constant?	Checked	No (CV 38%); ODE route closed
$ \beta = d_s/(2 \ln 2)$	Confirmed	3.4% finite- N
$\delta H(M)$ in closed form	Open	Needed for final proof
Definite sum proof	Open	Last remaining step

What Closing the Sum Proves

If the definite sum above is proved to equal $1/\ln 2$:

- κ is no longer a free parameter of the simulator. It is determined by the cascade as the ratio of two derived rates.
- $\beta = -d_s/(2 \ln 2)$ is a theorem, not an empirical finding.
- The spectral dimension d_s appears as the exchange rate between the phase geometry ($f_{12} = d_s/2$) and the information unit ($\ln 2$).
- CRF reaches maturity Level 7: the simulator's output is fully predicted by the δ -chain with zero free parameters at the level of β .

The sum is over the lifecycle of every node: how it accumulates mass, how long it waits, how much entropy it carries. That this sum is exactly $1/\ln 2$ means the cascade is balanced at the level of information — not by design, but because κ and $\ln 2$ are the only two scales, and balance demands their ratio. One proof stands between the framework and that theorem.