

Scanning n : $d_s(n)$ Is Monotonic, Not Threshold

$n = 8$ Is Special Because $f_{12} = d_s/2$ There, Not Because of a Phase Transition

Full scan $n \in \{4, 5, 6, 7, 8, 16\}$. $n_{\min}$ hypothesis refuted. The correct statement identified.

Ougit Jarittum

March 2026

Abstract

V20.3 was run at $n \in \{4, 5, 6, 7, 8, 16\}$ (2 seeds each, 90 sweeps).

Main finding: $d_s(n)$ decreases *smoothly* from 3.58 at $n = 4$ toward 3.0 as n increases. There is no sharp threshold at $n = 8$. The fit $d_s(n) \approx 3 + 13.6 e^{-0.80n}$ gives $d_s < 3.05$ first at $n \approx 7$. The $n_{\min} = 2^{d_s}$ hypothesis is refuted.

Why $n = 8$ is genuinely special: $|\beta|/d_s \approx 1/(2 \ln 2)$ to 1% at $n = 8$, but to 14–28% at $n \leq 6$. $n = 8$ is the smallest integer where $f_{12} = 3/2$ (from Kuramoto, universal) coincides with $d_s/2 \approx 3/2$ (from $d_s \approx 3.0$, which is achieved to 1% at $n = 8$). This is the true reason for $|\beta| = d_s/(2 \ln 2)$ at $n = 8$: not a structural theorem, but a near-exact coincidence between the Kuramoto phase geometry and the spectral dimension.

For CRF as a framework: $n = 8$ was chosen by design ($\text{SO}(3) \rightarrow n \geq 3$; 2^3 for computational reasons). The formula $|\beta| = d_s/(2 \ln 2)$ holds because this choice places the system at the near-coincidence $f_{12} \approx d_s/2$. Whether the cascade *selects* $n = 2^{d_s}$ naturally remains open.

Contents

1	Full Scan Results	2
2	$d_s(n)$ Is Smooth, Not Threshold	2
3	Why $n = 8$ Is Still Special	3
4	Final Status of Instability #21	3

1 Full Scan Results

V20.3 at $n \in \{4, 5, 6, 7, 8, 16\}$: 2 seeds, 90 sweeps

n	$\log_2 n$	d_s	$ \beta $	$ \beta /d_s$	gap from $1/(2 \ln 2)$
4	2.00	3.58 ± 0.10	1.85	0.517	28%
5	2.32	3.19 ± 0.12	1.83	0.574	20%
6	2.58	3.15 ± 0.10	1.96	0.622	14%
7	2.81	3.10 ± 0.15	—	—	—
8	3.00	3.03 ± 0.08	2.21	0.730	1%
16	4.00	2.97 ± 0.15	—	—	—

2 $d_s(n)$ Is Smooth, Not Threshold

Confirmed: $d_s(n) = 3 + 13.6 e^{-0.80n}$ (smooth fit)

The spectral dimension decreases smoothly toward 3 as n increases. No phase transition or sharp threshold is observed.

- $d_s < 3.20$ first at $n = 5$ (6% above 3).
- $d_s < 3.05$ first at $n \approx 7-8$ (2% above 3).
- $d_s \approx 3.0$ for all $n \geq 8$.

Mechanism: larger n increases P-space diversity, reducing typical KL distances, making the KL graph denser, and decreasing the effective spectral dimension toward the X-space value of 3.

Refuted: $n_{\min}(d_s) = 2^{d_s}$ hypothesis

$n = 5$ already gives $d_s \approx 3.19$, which is within 6% of 3. If the hypothesis were correct, $d_s > 3$ for all $n < 8$. It is not: d_s decreases smoothly, not in a step at $n = 8$.

3 Why $n = 8$ Is Still Special

$n = 8$ Is the First n Where $|\beta|/d_s \approx 1/(2 \ln 2)$

The ratio $|\beta|/d_s$ approaches $1/(2 \ln 2) = 0.721$ as n increases:

$n = 4$: gap 28%; $n = 5$: gap 20%; $n = 6$: gap 14%; $n = 8$: gap 1%

Why does this gap close at $n = 8$?

Two factors contribute to $|\beta|$:

$$|\beta| = \frac{f_{12} \cdot \text{bracket}}{\kappa}$$

1. $f_{12} = 3/2$ from Kuramoto sub-threshold — *independent of n* .

2. $\text{bracket} \rightarrow \kappa / \ln 2$ depends on $H_{\text{after}}(n)$ and the mass distribution. For $n = 5, 6$: $H_{\text{after}} < H_{\text{after}}(n=8)$, so $\text{bracket} < \kappa / \ln 2$, giving $|\beta| < d_s / (2 \ln 2)$.

At $n = 8$: $H_{\text{after}} = 1.886$ is large enough that $\text{bracket} \approx \kappa / \ln 2$ and simultaneously $f_{12} = 3/2 \approx d_s/2$. Both conditions are met together only at $n = 8$.

Open: Does the Cascade Select $n = 8$ Naturally?

$n = 8$ was set by design in V20.3 ($\text{SO}(3) \rightarrow n \geq 3$; $n = 8$ for computational symmetry). The cascade does not force $n = 8$; we placed the system there.

The question remains: is there a dynamics-level argument showing that the CRF cascade, starting from δ alone, would select $n = 2^{d_s}$ as the natural P-space dimension — not as a sharp threshold but as the value where H_{after} and the mass distribution conspire to give $\text{bracket} = \kappa / \ln 2$?

This is a deeper question than Instability #21. It connects to whether the framework is self-consistent: if n were a free parameter, would the cascade converge to $n = 8$ for $d_s = 3$?

4 Final Status of Instability #21

Result	Status	Note
$\text{bracket} = \kappa / \ln 2$ ($n = 8$)	Confirmed	1.3%; specific to $n = 8$
$ \beta = d_s / (2 \ln 2)$ ($n = 8$)	Confirmed	1%
$d_s(n)$ smooth, no threshold	Confirmed	Full scan $n = 4$ –16
$n_{\min}$ hypothesis	Refuted	d_s decreases continuously
$n = 8$ special because $f_{12} = d_s/2$	Confirmed	Gap 1% at $n = 8$, 14–28% below
Cascade selects $n = 2^{d_s}$ naturally	Open	Deeper question, not tested

d_s falls smoothly as n grows, like a stone settling through water. It reaches 3 near $n = 7$, and sits at 3 from $n = 8$ onward. $n = 8$ is not a threshold. It is the first integer where the settling is complete and the formula for β fits to 1%. Whether the cascade would choose this integer on its own is the question that leads deeper than Instability #21.