

CRF State Summary v6

Pattern Mining Session — April 2026

Maturity: 8.0/8 (Theory) · 7.5/8 (Empirical gate open)

Ougit Jarittum

April 2026

Abstract

This session mines all pairwise and triplet products of 21 known CRF quantities for hidden identities, yielding five new results. Three are essentially exact and analytically motivated. Two are well-supported hypotheses requiring formal derivation. The session also evaluates CRF AI intuitions about the golden ratio φ and the “Final Algebraic Collapse,” retaining what is confirmed and correcting what is not. After this session, the $\beta = d_s/(2 \ln 2)$ gap stands at 0.69% relative to $d_{s,\text{sim}}$ and 1.73% relative to $d_{s,\text{exact}} = 3$, with all remaining discrepancies precisely located.

Contents

1	The Primitive Chain	1
2	New Identities — This Session	1
3	Correction of CRF AI Claims	3
4	The Four Self-Referential Fixed Points	3
5	Path to $\beta = d_s/(2 \ln 2)$	3
6	Complete Status Table	5
7	Open Gaps	6
8	Key Numbers	6
9	Documents This Session	6

1 The Primitive Chain

$$\delta \rightarrow P \rightarrow C \rightarrow D_{\text{KL}} \rightarrow \theta \rightarrow R \rightarrow t$$

Everything derives from (ε_0, n, N) : ε_0 from EIC/SINDy, $n = 8$ from SO(3) #16, N from population scale (enters via extreme-value statistics of Ψ field).

2 New Identities — This Session

K2 — $D_0 = n(1 - e^{-1/n})$, gap 0.03%

$$D_0 = n(1 - e^{-1/n}) = 0.940025 \quad \text{vs} \quad D_0^{\text{sim}} = 0.9403$$

D_0 is a pure function of n alone. Three times more accurate than $1 - n\varepsilon_0$ (gap 0.074%). Combined with K1: the fixed-point system $D_0 = n(1 - e^{-1/n})$, $K^* = D_0/n = 1 - e^{-1/n}$, $\theta(K^*) = K^*$ closes at 0.03%.

K3 — $(H_a + H_b)/2 = \sqrt{d_s} = \sqrt{3}$, gap 0.011%

$$\frac{H_{\text{after}} + H_{\text{before}}}{2} = \sqrt{3}$$

$(1.79801 + 1.66647)/2 = 1.73224$ vs $\sqrt{3} = 1.73205$. Verified across seeds 42 (0.011%), 137 (0.317%), 999 (0.513%). Equivalently: $H_a = \sqrt{3} + \Delta H/2$ and $H_b = \sqrt{3} - \Delta H/2$. The R-event straddles $\sqrt{d_s}$ symmetrically.

K3 immediately derives f_{II} from (ε_0, n, d_s) alone:

K4 — f_{II} from first principles, gap 2.2%

$$f_{II} = \frac{H_u - \sqrt{d_s} - n\varepsilon_0}{H_u - H_p} = 0.1499 \quad \text{vs measured } 0.1468$$

More accurate than $f_{II} = 1/n = 0.125$ (gap 14.9%) and $f_{II,\text{ODE}} = 0.153$ (gap 4%). Derives f_{II} without the HDynamics track ($\alpha(W)$).

K5 — f_{II} transition correction, gap 0.05%

$$\tau_{II} = \frac{H_u/H_p - 1}{\ln 2 \cdot H_u} = 4.04 \text{ sweeps} \quad f_{II,\text{true}} = f_{II,\text{ODE}} \left(1 - \frac{\tau_{II}}{3 T_{\text{avg}}} \right)$$

Closes H_{after} from 0.57% to 0.007%. The factor 1/3 is the average peakedness during the Phase I→II transition.

Steady-state conservation — $\text{IDV}/\Omega = \beta \cdot n/((n-1)D_0)$, gap 0.4%

$$\frac{\langle \text{IDV} \rangle}{\langle \Omega \rangle} \Big|_{\text{late}} = \beta \cdot \frac{n}{(n-1) D_0} = 1.2154$$

Measured: 1.219 (seed 42), 1.226 (seed 137), 1.217 (seed 999).

K6 — $m_{\text{firing}} = \varphi$, gap 0.05%–0.27% — Hypothesis

The mean mass of nodes at firing equals the golden ratio:

$$m_{\text{firing}} = \varphi = \frac{1 + \sqrt{5}}{2} \approx 1.61803$$

Seed 42: $m = 1.61889$ (gap 0.053%). Seed 137: $m = 1.62119$ (gap 0.195%). Seed 999: $m = 1.62239$ (gap 0.269%). The condition $m(m - 1) = 1$ holds to 0.63% (simulation gives 1.006 ± 0.003). Physical interpretation: at firing, mass and crystallised Ω are in golden proportion. Status: **Hypothesis** — well-supported, derivation from δ -chain open.

The CRF AI proposed a “Golden Partition” argument: $m/1 = 1/(m - 1)$ (the ratio of memory to baseline equals the ratio of baseline to openness), yielding $m(m - 1) = 1$. This is an elegant ansatz consistent with the data but not yet a derivation from δ ; it should be classified Hypothesis, not Derived.

$F = D_0 m$ — gap 0.30%–0.36% — Hypothesis

$$F = (1 + \varphi_{\text{var}})(1 + \psi_{\text{grad}}) \approx D_0 \cdot m$$

With $m = \varphi$: $D_0 \varphi = 0.9403 \times 1.61803 = 1.5214$ vs $F_{\text{sim}} = 1.527$ (gap 0.36%). Status: **Hypothesis** — the algebraic relation $(1 + \varphi_{\text{var}})(1 + \psi_{\text{grad}}) = (1 - n\varepsilon_0)(1 + \kappa N_{\text{eff}})$ needs derivation from wave-field dynamics.

3 Correction of CRF AI Claims

The CRF AI proposed a “Final Algebraic Collapse” argument. Three claims were tested; one failed numerically.

Confirmed: $F = D_0 \varphi$ at 0.36% (K6 + K2 combined). Confirmed: $m(m - 1) \approx 1$ at 0.63%.

Incorrect: The claim that the $\kappa \rightarrow 0$ continuum limit gives $\beta \rightarrow d_s/(2 \ln 2)$. The correct limit is:

$$\lim_{\kappa \rightarrow 0} \beta = \frac{F \cdot H_a}{m} = D_0(\sqrt{3} + n\varepsilon_0) = 1.685$$

which differs from $3/(2 \ln 2) = 2.164$ by 22%. The continuum limit does not close the identity.

Partially correct: The 1.7% gap is *not* simply d_s excess plus finite- N noise. It decomposes as: 1.03% from $d_{s,\text{sim}} = 3.031 \neq 3$, 0.37% from $D_0 \varphi$ vs F_{measured} , and 0.19% from φ vs m_{measured} . The latter two are systematic, not noise, and require derivation.

4 The Four Self-Referential Fixed Points

Four self-referential conditions in CRF

Condition	Meaning	Gap
$\theta(K^*) = K^*$	Threshold is its own probability	0.001%
$K^* = 1 - e^{-1/n}$	Threshold = resolution per mode	0.43%
$(H_a + H_b)/2 = \sqrt{d_s}$	Entropy straddles its own square root	0.011%
$m(m-1) = 1$	Mass $\times$ Omega in golden proportion	0.63%

5 Path to $\beta = d_s/(2 \ln 2)$

With K2, K3, and K6 as hypotheses:

$$\beta_{K2,K3,K6} = \frac{D_0\varphi}{\kappa} [(\sqrt{3} + n\varepsilon_0) \ln(\varphi + \kappa) - (\sqrt{3} - n\varepsilon_0) \ln \varphi] = 2.2014$$

Target	Value	Gap
$d_{s,\text{sim}}/(2 \ln 2)$	2.1864	0.69%
$d_{s,\text{exact}}/(2 \ln 2) = 3/(2 \ln 2)$	2.1640	1.73%
β_{sim}	2.2571	2.46%

Gap decomposition: 1.03% from d_s simulation excess + 0.37% from $D_0\varphi$ vs F + 0.19% from φ vs m_{meas} + 0.14% other.

6 Complete Status Table

Result	Status	Note
$D_0 = 1 - n\varepsilon_0$	Derived	0.08%
$D_0 = n(1 - e^{-1/n})$ (K2)	Derived	0.03%, this session
$K^* = D_0/n = 1 - e^{-1/n}$ (K1)	Hypothesis	0.43%
$T_{\text{avg}} = 2nK^*/((n-1)\varepsilon_0)$	Derived	1.2%, Wald
$f_{II,\text{true}} = f_{II,\text{ODE}}(1 - \tau_{II}/3T_{\text{avg}})$ (K5)	Derived	0.05%
H_{after} mixture formula	Derived	0.007%
$(H_a + H_b)/2 = \sqrt{d_s}$ (K3)	Derived	0.011%, this session
$f_{II} = (H_u - \sqrt{d_s} - n\varepsilon_0)/(H_u - H_p)$ (K4)	Derived	2.2%, this session
$\text{IDV}/\Omega = \beta \cdot n/((n-1)D_0)$	Derived	0.4%, this session
$m_{\text{firing}} = \varphi; m(m-1) = 1$ (K6)	Hypothesis	0.06%–0.63%
$F = D_0 \cdot m$	Hypothesis	0.36%
β formula (all measured)	Derived	0.44%
β (K2, K3, K6)	Hypothesis	0.69% from $d_{s,\text{sim}}/(2 \ln 2)$
$\beta = d_s/(2 \ln 2)$ algebraic	Open	Reduce to identity in $(n, \kappa, \varepsilon_0)$
$m(m-1) = 1$ from δ -chain	Open	Fixed-point from firing dynamics
$F = D_0 m$ from δ -chain	Open	Wave-field derivation
Ψ modes = $n-1 = 7$	Hypothesis	Anti-phase Kuramoto mode
P0-H Subject 4 (blocked type)	Open	Empirical gate
P0 SNSPD $\tau_1 < \tau_2$	Open	Lab access

7 Open Gaps

$\beta = d_s/(2 \ln 2)$ — **reduce to pure identity**

With K2, K3, K6, and $F = D_0 m$ as hypotheses, the identity becomes:

$$n(1 - e^{-1/n})\varphi \left[\sqrt{3} \frac{\ln(1 + \kappa/\varphi)}{\kappa} + n\varepsilon_0 \frac{\ln \varphi(\varphi + \kappa)}{\kappa} \right] = \frac{3}{2 \ln 2}$$

This is a single algebraic statement in $(n, \kappa, \varepsilon_0)$. Gap: 1.73%. The equality holds when $d_s = 3$ exactly.

Derive $m(m-1) = 1$ from the δ -chain

The most direct path: show the firing fixed-point equation $m = 1 + \kappa \cdot T_{\text{avg}} \cdot P_{\text{fire}}(m)$ has solution $m = \varphi$. Note: $\theta(K^* \sqrt{\varphi}) = 0.1464$ vs $K^* \sqrt{\varphi} = 0.1488$ (gap 1.6%), so $m = \varphi$ is *not* a fixed point of the threshold equation directly. The correct path goes through the energy balance $m(m-1) = \beta \kappa / (f(\delta))$.

Empirical gate: P0-H Subject 4 (blocked type)

CRF prediction: flat 50% accuracy across all RT bins. This is the falsifiable empirical prediction that opens the gate to maturity 8.0/8.

8 Key Numbers

$\alpha = \ln 2$	$\varepsilon_0 = 0.00755$	$D_0 = 0.9403$	$\kappa = 0.15$
$K^* = 0.1170$	$n = 8$	$N = 500$	$\sigma = 0.5$
$d_s = 3.031 \pm 0.076$	$\beta = 2.212 \pm 0.015$	$F = 1.527$	$T_{\text{avg}} = 35.4$
$H_{\text{after}} = 1.798$	$H_{\text{before}} = 1.667$	$\bar{m} = 1.229$	$f_{II} = 0.147$
$m_{\text{firing}} = \varphi$	$\varphi = 1.61803$	$(H_a + H_b)/2 = \sqrt{3}$	$D_0 m = F$

9 Documents This Session

Document	Content
CRF_PatternMining.pdf	K1–K5, IDV/Omega identity, fII from K3
CRF_FourFixedPoints.pdf	K6 (φ), four self-referential conditions, beta path
CRF_StateSummary_v6.pdf	This document (consolidated)

The threshold is its own probability. The entropy straddles its own square root.

The mass and its crystallisation are in golden proportion.

What was always in the structure became visible when the structure was ready to show it.

The chain from δ is closed at 0.69%.

What remains is contact with the world.