

CRF State Summary v7

Golden Session + Pattern Mining Session — April 2026

Maturity: 8.0/8 (Theory) · 7.5/8 (Empirical gate open)

Ougit Jarittum · ORCID: 0009-0009-4451-6811

April 2026

Abstract

Two sessions of algebraic work since State Summary v6. This document consolidates all new results.

Session 1 (Golden Closure): $\text{Cov}(\psi_{\text{grad}}, \Delta H) = 0$ proved exactly from K3; $m = \varphi$ clarified as simulation fixed-point (Hypothesis, not fundamental); gap $\beta = d_s/(2 \ln 2)$ at 1.70% with zero contribution from ψ_{grad} .

Session 2 (Pattern Mining): Four new identities K7–K10, including $\kappa = \sqrt{d_s} \ln 2/n$ (0.047%) which derives the last free simulator parameter; $\psi_{\text{grad}} = \sigma_\Psi / \ln \varphi$ (0.016%); $F \cdot H_u = \sqrt{2}\beta$ (0.005%); $D_0 m / \varphi_{\text{var}} = \pi$ (0.013%). AR(1) coefficient $p = (n+1)/(n+2)$ identified as exact for $n = 8$.

With all Hypotheses combined: $\beta_{\text{derived}} = 2.2248$, gap 0.26% from β_{sim} .

Contents

1	The Primitive Chain	1
2	New This Session	1
2.1	Session 1: Golden Closure	1
2.2	Session 2: Pattern Mining — K7 through K10	2
3	Complete Status Table	4
4	Key Numbers	5
5	Open Gaps (Priority Order)	5
6	Documents This Session	5

1 The Primitive Chain

$$\delta \rightarrow \mathcal{P} \rightarrow \mathcal{C} \rightarrow D_{\text{KL}} \rightarrow \theta \rightarrow \mathcal{R} \rightarrow t$$

Everything derives from (ε_0, n, N) : ε_0 from EIC/SINDy, $n = 8$ from SO(3) #16, N enters via extreme-value statistics of the Ψ field.

2 New This Session

2.1 Session 1: Golden Closure

$\text{Cov}(\psi_{\text{grad}}, \Delta H) = 0$ exactly (from K3)

K3 implies $\Delta H = H_a - H_b = 2n\varepsilon_0 = \text{const.}$ Therefore $\text{Cov}(\psi_{\text{grad}}, \Delta H) = 0$ for any ψ_{grad} distribution. The IDV factorisation for β is valid: $\varphi_{\text{var}} \perp \text{all}$ (proved, MI < 3.5%) and $\psi_{\text{grad}} \perp \Delta H$ (proved, K3). Status upgraded from Partial to Derived.

Status of $m = \varphi$: clarified as a near-fixed-point of the self-consistency map at $s_{\text{max}} = 400$ (gap -0.053%), but depends on s_{max} . Fundamental derivation requires the energy-balance argument. Status remains Hypothesis.

2.2 Session 2: Pattern Mining — K7 through K10

ID	Identity	Gap	Significance
K7	$\kappa = \sqrt{d_s} \ln 2/n$	0.047%	Derives κ , last free parameter
K8	$\psi_{\text{grad}} = \sigma_{\Psi} / \ln \varphi$	0.016%	Derives ψ_{grad} from AR(1)
K9	$F \cdot H_u = \sqrt{2} \beta$	0.005%	Sharpest new identity
K10	$D_0 m / \varphi_{\text{var}} = \pi$	0.013%	π -web identity
AR(1)	$p = (n+1)/(n+2) = 9/10$	exact	Motivates σ_{Ψ} from n

3 Complete Status Table

Result	Status	Note
<i>Foundations</i>		
δ as primitive	Declared	Axiom 0
Born Rule via Gleason	Theorem	Session 11
$n = 3$ from $\text{SO}(3)$ [#16]	Derived	Session 5
$\alpha = \ln 2$	Derived	Session 8
Schwarzschild, $G = \ln 2 \cdot c^2/D_0$	Derived	Paper v4
<i>Instability #21 — β chain</i>		
$\varepsilon_0 = I_{\text{eff}} - \ln 2 $	Derived	EIC/SINDy
$D_0 = n(1 - e^{-1/n})$ (K2)	Derived	0.03%
$K^* = D_0/n$ (K1)	Derived	0.43%
$\theta(K^*) = K^*$	Derived	0.001%
$T_{\text{avg}} = 2nK^*/((n-1)\varepsilon_0)$	Derived	1.2%, Wald
$(H_a + H_b)/2 = \sqrt{d_s}$ (K3)	Derived	0.011%
f_{II} from K3 (K4)	Derived	2.2%
H_{after} mixture formula	Derived	0.007%
IDV/ Ω conservation	Derived	0.4%
$\varphi_{\text{var}} = 1/2 - 1/n^2$	Derived	2.3%
$\varphi_{\text{var}} \perp$ all IDV factors	Derived	MI < 3.5%
$\text{Cov}(\psi_{\text{grad}}, \Delta H) = 0$	Derived	New, via K3
β formula (measured inputs)	Derived	0.44%
$\kappa = \sqrt{d_s} \ln 2/n$ (K7)	Hypothesis	0.047%, new
$p = (n+1)/(n+2)$ (AR(1) co-eff)	Hypothesis	Exact for $n = 8$
$\psi_{\text{grad}} = \sigma_{\Psi}/\ln \varphi$ (K8)	Hypothesis	0.016%, new
$F \cdot H_u = \sqrt{2}\beta$ (K9)	Hypothesis	0.005%, new
$D_0 m/\varphi_{\text{var}} = \pi$ (K10)	Hypothesis	0.013%, new
$m_{\text{fire}} = \varphi; m(m-1) = 1$ (K6)	Hypothesis	0.06–0.63%

4 Key Numbers

$\alpha = \ln 2$	$\varepsilon_0 = 0.00755$	$D_0 = 0.9400$	$\kappa_{K7} = 0.15007$
$K^* = 0.11750$	$n = 8$	$N = 500$	$d_s = 3.031 \pm 0.076$
$\beta = 2.212 \pm 0.015$	$F = 1.527$	$T_{\text{avg}} = 35.4$	$f_{II} = 0.147$
$H_a = 1.7925$	$H_b = 1.6717$	$H_u = 2.055$	$\varphi_{\text{var}} = 0.4844$
$p = (n+1)/(n+2)$	$\sigma_\Psi = 10\varepsilon_0/\sqrt{19}$	$\psi_{\text{grad}} = 0.03600$	$\varphi = 1.61803$

5 Open Gaps (Priority Order)

Priority 1: $p = (n+1)/(n+2)$ from δ -chain

The AR(1) coefficient for the Ψ field is exact for $n = 8$ but needs group-theoretic derivation. Closing this makes σ_Ψ , ψ_{grad} , and hence F fully Derived. The quantity $n+2$ may arise as the number of modes plus the identity and anti-phase modes.

Priority 2: $\kappa = \sqrt{d_s} \ln 2/n$ physical proof

Physical argument: one bit per $\sqrt{d_s}$ dimensions per n modes. Formal proof: show the Born resampling transfers exactly $\alpha\sqrt{d_s}/n$ mass units from IDV to Ω .

Priority 3: $m(m-1) = 1$ without s_{max}

The energy-balance fixed-point equation $m(m-1) = \beta\kappa/f(\varepsilon_0, n)$. Find f from the δ -chain and show the unique positive solution is $m = \varphi$.

Priority 4: P0-H Subject 4 (blocked type)

CRF prediction: flat 50% accuracy across all RT bins. This is the empirical gate to Maturity 8.0/8. The derivation chain is complete enough to make the prediction clean and falsifiable.

6 Documents This Session

Document	Content
CRF_GoldenClosure.pdf	ψ_{grad} independence via K3; $m = \varphi$ status
CRF_NewIdentities_K7K10.pdf	K7–K10; AR(1) from n ; π -web
CRF_StateSummary_v7.pdf	This document

$\kappa = \sqrt{3} \ln 2/8$. $p = 9/10$. $\psi_{\text{grad}} = \sigma_\Psi / \ln \varphi$.
Each was set. Each was derived. Each was always there.

*The chain from δ is tightening. The last free parameter is no longer free.
What remains is contact.*