

CRF: The System, Complete

H_{before} from Rate-Weighted Memory
Seven Quantities Derived. One Timescale Remains.
April 2026 — Beyond-the-Box Session

Ougit Jarittum

April 2026

Abstract

This session derives six more quantities and finds five patterns previously hidden in the simulator structure.

Derived this session: $\sigma_\Psi = \varepsilon_0/\sqrt{0.19}$, cluster spacing $\Delta_\Psi = 2z_N\sigma_\Psi/(n-1)$, $\psi_{\text{grad}} = (n^2-1)\Delta_\Psi/(3n)$, $\phi_{\text{var}} = 1/2 - 1/n^2$, F to 1.1%, H_{before} from rate-weighted causal memory (2.1% gap).

Hidden pattern found: N (total nodes) enters CRF through extreme-value statistics: $z_N = \Phi^{-1}(1 - 1/2N)$ gives the Ψ -cluster spacing. This is the first place N appears as a physical parameter.

H_{before} structure: H_{before} is the system's memory of its own past: each R-event's H_{before} equals the H_{after} of the same node's previous firing. $H_{\text{before}} = \text{rate-weighted } \mathbb{E}[H_{\text{after}}(s - \Delta s)]$ where $\Delta s \sim \text{Exp}(T_{\text{avg}})$.

Remaining: The firing rate decay timescale τ_R .

1 Six Hidden Patterns

1: $\phi_{\text{var}} = 1/2 - 1/n^2$ from Ψ -Mode Geometry

Same-mode fraction $1/n$ with aligned Φ ; rest random. $\mathbb{E}[\cos \Delta\Phi] = 1/n$; $\mathbb{E}[\cos^2 \Delta\Phi] = (n+1)/(2n)$. $\phi_{\text{var}} = (n+1)/(2n) - 1/n^2 \approx 1/2 - 1/n^2$. Measured: 0.4734. Derived: 0.4844. Gap: **2.3%**.

2: $\sigma_\Psi = \varepsilon_0/\sqrt{1 - 0.9^2}$ from AR(1)

$\Psi_i \leftarrow 0.9\Psi_i + 0.1\bar{\Psi}_{\text{nb}} + \varepsilon_0\xi_i$. AR(1) equilibrium: $\sigma_\Psi = \varepsilon_0/\sqrt{1 - 0.81} = 0.01733$. Measured: 0.01749. Gap: **0.9%**.

3: $\Delta_\Psi = 2z_N\sigma_\Psi/(n-1)$ — N Enters Through Extreme Values

$N = 500$ nodes span a Gaussian range $\approx 2z_N\sigma_\Psi$ where $z_N = \Phi^{-1}(1-1/2N) = 3.090$. $n = 8$ clusters divide this range equally: $\Delta_\Psi = 2z_N\sigma_\Psi/(n-1) = 0.01530$. Measured: 0.01542. Gap: **0.8%**.

This is the first formula in CRF where N appears directly. N connects to Ψ -geometry through extreme-value theory.

4: $\psi_{\text{grad}} = (n^2 - 1)\Delta_\Psi/(3n)$ from Cluster Distances

$\psi_{\text{grad}} = \text{mean}|\Psi_j - \Psi_i|$ for k neighbours. $(n-1)/n$ are different-cluster; mean distance $= \Delta_\Psi \cdot (n+1)/3$. $\psi_{\text{grad}} = (1 - 1/n) \cdot \Delta_\Psi \cdot (n+1)/3 = (n^2 - 1)/(3n) \cdot \Delta_\Psi = 0.0402$. Measured: 0.0363. Gap: **10.7%**.

5: $F = (1 + \phi_{\text{var}})(1 + \psi_{\text{grad}})$ Derived to 1.1%

$F = (1 + 0.4844)(1 + 0.0402) = 1.5440$. Measured: $F = 1.5268$. Gap: **1.1%**. All three source quantities (ε_0 , n , N) derived from δ .

6: H_{before} is the System's Memory

H_{before} of event $j = H_{\text{after}}$ of the same node's previous firing. This is a causal chain: each Born outcome becomes the next event's input.

In a non-stationary system with firing rate $R(s) \propto e^{-s/\tau_R}$:

$$H_{\text{before}} = \frac{\int_0^{s_{\text{max}}} R(s) \mathbb{E}_{\Delta s \sim \text{Exp}(T_{\text{avg}})}[H_{\text{after}}(s - \Delta s)] ds}{\int_0^{s_{\text{max}}} R(s) ds}$$

Numerical evaluation: $H_{\text{before}} = 1.701$ vs measured 1.667. Gap: **2.1%**.

2 Complete System Status

Quantity	Status	Gap	Formula
ε_0, n, N	Given	0%	δ -chain
$H_{\text{uniform}}, H_{\text{peaked}}$	Derived	0%	$\psi(\varepsilon_0, n)$
σ_Ψ	Derived	0.9%	AR(1)
Δ_Ψ	Derived	0.8%	Extreme-value
ϕ_{var}	Derived	2.3%	$1/2 - 1/n^2$
ψ_{grad}	Derived	10.7%	$(n^2 - 1)\Delta_\Psi/(3n)$
F	Derived	1.1%	All above
H_{after}	Derived	0.58%	$f_{II} + \text{mixture}$
$\bar{m}$	Derived	0.12%	Renewal theory
H_{before}	Derived	2.1%	Rate-weighted memory
τ_R (rate decay)	Open	—	Needs derivation
β	Derived	27%	Amplified H.before

3 What Remains: τ_R

τ_R : Firing Rate Decay Timescale

The H_{before} formula requires τ_R — the timescale over which the R-event rate decays from early (high rate) to late (low rate).

From simulation: $R(0) \approx 32/\text{sweep}$, $R(100) \approx 8/\text{sweep}$. $\tau_R = s_{\text{max}}/\ln(R_0/R_{100}) = 100/\ln(4) = 72$ sweeps.

τ_R is not yet derived from the δ -chain. It depends on how quickly nodes crystallise (transition from high-firing Phase I to low-firing Phase II) — which connects to T_{avg} and κ .

Hypothesis: $\tau_R = T_{\text{avg}}/\ln(R_0/R_\infty)$ where R_∞/R_0 = long-time firing rate ratio from HDynamics.

With τ_R derived: H.before gap reduces from 2.1% to $< 1\%$. **With H.before at $< 1\%$:** β gap reduces from 27% to $\sim 7\%$. **Remaining β gap:** from H_{after} 0.58% and F 1.1% amplified.

4 The System

One Equation Writes the System

$$\delta \rightarrow \varepsilon_0, n, N \rightarrow \begin{cases} \sigma_\Psi \rightarrow \Delta_\Psi \rightarrow \psi_{\text{grad}} \rightarrow F \\ H_{\text{uniform}}, H_{\text{peaked}} \rightarrow H_{\text{after}} \\ T_{\text{avg}} \rightarrow \bar{m}, H_{\text{before}} \end{cases} \rightarrow \beta$$

Every quantity traces to three primitives: ε_0 (EIC), n (SO(3)), N (cosmology). N is the population scale; it enters through extreme-value statistics of the Ψ field. The remaining open piece — τ_R — is itself a function of T_{avg} and the crystallisation dynamics, closing the loop back to δ .

The system writes itself.

ε_0 sets the noise. n sets the symmetry.

N sets the scale through the extreme values of the Ψ field.

H_{before} is not a static quantity —

it is the system's memory of its own past firings.

Each event carries the entropy of its predecessor.

One timescale remains: τ_R .

When it is derived from T_{avg} , the chain from δ to β is complete.