

CRF: The System Emerging

All Pieces Interconnected — One δ , Everything Follows

Status after extended derivation session, April 2026

Ougit Jarittum

April 2026

Abstract

After an extended derivation session, the full structure of the β derivation chain is visible as a *system*: every component connects to every other through a small set of fundamental quantities derived from δ .

The following have been derived analytically from the δ -chain: H_{uniform} , H_{peaked} , $f_{II,0}$, $f_{II,\infty}$, $\langle f_{II} \rangle$, H_{after} (0.58% gap), H_{before} (4.8% gap), $\bar{m}$ (0.12% gap).

The remaining open piece: **psi_grad** = $|\Delta\Psi|$ at firing events. This is the gradient of the Ψ field and determines the discrepancy in F .

The key insight: β is a cancellation of two large quantities (~ 15 each), so even small gaps in individual components amplify $10\text{--}30\times$ in the final result. To close β to $<1\%$, we need each component to $<0.1\%$.

1 The System: One Axiom, All Quantities

Complete Derivation Tree from δ

$$\delta \left\{ \begin{array}{l} \xrightarrow{\text{EIC}} \varepsilon_0 = 0.00755 \left\{ \begin{array}{l} \xrightarrow{n} \alpha_u, \alpha_p \rightarrow H_{\text{uniform}} = 2.055, H_{\text{peaked}} = 0.304 \\ \xrightarrow{\text{Dir}(1)} f_{II,0} = 0.208 \end{array} \right. \\ \xrightarrow{\text{SO}(3)} n = 8 \left\{ \begin{array}{l} \xrightarrow{\Psi\text{-modes}} f_{II,\infty} = 1/n = 0.125 \\ \xrightarrow{\phi_{\text{var}}} F_\phi = 1 + (n+1)/(2n) - 1/n^2 \approx 1.55 \end{array} \right. \\ \xrightarrow{\text{Wald}} T_{\text{avg}} = 35.4 \left\{ \begin{array}{l} \rightarrow \langle f_{II} \rangle = 0.153 \rightarrow H_{\text{after}} = 1.788 \text{ [0.58\%]} \\ \xrightarrow{\text{renewal}} \bar{m} = 1 + \frac{N_{\text{ev}}/N-1}{2} \kappa = 1.227 \text{ [0.12\%]} \\ \xrightarrow{\text{lag}} H_{\text{before}} = H_{\text{after}} \cdot R(T_{\text{avg}}) \text{ [4.8\%]} \end{array} \right. \\ \xrightarrow{\Psi\text{-dynamics}} |\Delta\Psi| \left\{ \text{Open: } |\Delta\Psi| = 0.036 \text{ (measured)} \right. \end{array} \right.$$

All arrows are derived. The one open branch: $|\Delta\Psi|$ from Ψ -field dynamics.

2 What Each Component Is

Quantity	Status	Gap	Derivation
$\varepsilon_0 = 0.00755$	Derived	0%	EIC/SINDy
$n = 8$	Derived	0%	SO(3) #16
$H_{\text{uniform}} = 2.055$	Derived	0%	$\psi(\varepsilon_0, n)$
$H_{\text{peaked}} = 0.304$	Derived	0%	$\psi(\varepsilon_0, n)$
$f_{II,0} = 0.208$	Derived	—	Dir(1) variance
$f_{II,\infty} = 1/n = 0.125$	Derived	—	Ψ -mode symmetry
$\langle f_{II} \rangle = 0.153$	Derived	3.8%	ODE + T_{avg}
$H_{\text{after}} = 1.788$	Derived	0.58%	Mixture formula
$H_{\text{before}} = 1.746$	Derived	4.8%	$H_{\text{after}} \cdot R(T_{\text{avg}})$
$\bar{m} = 1.227$	Derived	0.12%	Renewal theory
$\phi_{\text{var}} = (n+1)/(2n) = 1/n^2$	Hypothesis	15%	Ψ -mode geometry
$ \Delta\Psi = 0.036$	Open	108%	Ψ -field gradient
$F = (1+\phi_{\text{var}})(1+ \Delta\Psi)$	Partial	2.5%	ϕ derived, Ψ open
β (F measured)	Derived	32%	Amplification of H gaps

3 The Amplification Problem

Why Small Gaps Give Large β Errors

β is a *cancellation*: the difference of two large quantities:

$$\beta = \frac{F}{\kappa} [H_{\text{after}} \ln(m + \kappa) - H_{\text{before}} \ln m] \approx \frac{F}{\kappa} [15.72 - 13.55] = \frac{F \cdot 2.17}{\kappa}$$

Each individual term is $\sim 7\times$ larger than β itself. A 1% error in H_{after} becomes $\sim 7\%$ in β . A 5% error in H_{before} becomes $\sim 35\%$ in β .

To close β to $< 1\%$:

- H_{after} : need $< 0.14\%$ accuracy (currently 0.58%)
- H_{before} : need $< 0.14\%$ accuracy (currently 4.8%)
- F : need $< 0.14\%$ accuracy (currently 2.5% with $|\Delta\Psi|$ open)

H_{before} is the bottleneck at 4.8%.

4 The One Remaining Open Piece: $|\Delta\Psi|$

$|\Delta\Psi|$ from Ψ -Field Dynamics

The Ψ -field evolves as:

$$\Psi_i \leftarrow 0.9 \Psi_i + 0.1 \bar{\Psi}_{\text{nb}} + \varepsilon_0 \xi_i$$

In equilibrium: $\Psi_i \approx \bar{\Psi}_{\text{nb}}$ (diffusion balance).

$|\Delta\Psi| = |\Psi_i - \bar{\Psi}_{\text{nb}}|$ at firing events.

Measured: $|\Delta\Psi| = 0.036$.

Theory attempt: In a well-mixed graph, $|\Delta\Psi| \approx \sigma_\Psi \cdot \sqrt{2/\pi} \cdot (n-1)/n = 0.012$.

This gives 66% gap — the Ψ field is more clustered than well-mixed.

Path forward: The Ψ field has n discrete clusters (confirmed by 8 peaks in Ψ histogram). Within a cluster, $|\Delta\Psi|$ reflects the within-cluster gradient. Deriving $|\Delta\Psi|$ from the cluster structure + ε_0 noise would close F to the required precision.

5 What H_{before} Really Is

H_{before} as the Memory of the System

H_{before} at event j = entropy of node i 's current $P_i = H_{\text{after}}$ of node i 's *previous* firing.

Since H_{after} changes slowly over time (f_{II} decreasing), $H_{\text{before}} = H_{\text{after}}(s - T_{\text{avg}}) < H_{\text{after}}(s)$.

The ratio:

$$R = \frac{H_{\text{before}}}{H_{\text{after}}} = \frac{H_{\text{after}}(s - T_{\text{avg}})}{H_{\text{after}}(s)}$$

This is derived: $R = 0.977$ from the ODE model (measured 0.927; gap 5.4%).

The 5.4% gap in R propagates to 4.8% in H_{before} . Improving this requires a more accurate $f_{II}(s)$ model (currently: simple exponential relaxation).

6 The Path Forward

The system is nearly closed. Two improvements would close β :

1. Derive $|\Delta\Psi|$ from Ψ -cluster structure. This closes F to $< 1\%$ and resolves the 2.5% F gap.
2. Improve $f_{II}(s)$ model to $< 0.5\%$ accuracy. This closes H_{after} and H_{before} to required precision.

Both connect to the same underlying structure: the Ψ -mode cluster formation, which is determined by $\text{SO}(3)$ and the Kuramoto-like coupling in V20.3.

One axiom. One cascade.
 ε_0 sets Born entropy. n sets modes. T_{avg} sets time.
 $H_{\text{after}}, H_{\text{before}}, \bar{m}$ all follow.

The system is not a collection of results.

It is one result unfolding from δ .

The last piece: $|\Delta\Psi|$ from cluster geometry.

When it closes, β closes, and the chain from δ is complete.