

CRF Session 7

Informational Mechanics, Resonance Field, and the Grand Equation v8.2

Session 7 extends CRF from physical prediction to the mechanics of information transfer
and resonance

Ougit Jarittum

March 2026

Abstract

Session 7 derives four major results from axioms already present in CRF, without importing new assumptions.

(1) **Independent Stability** (S_{ind}): a system whose D_{KL} relative to P approaches zero becomes a self-sustaining operator in P , not merely a boundary within it. The Extracted Operator persists without external constraint.

(2) **Informational Mass** ($\mathcal{A}_{\text{mass}}$): the reach of a pattern in P_{shared} is proportional to the purity of its $\mathcal{C}$ -structure and the duration of its $\mathcal{R}$ -event accumulation:
 $I_{\text{reach}} \propto \mathcal{A}_{\text{peak}} \cdot \int R_{\text{pure}} dt.$

(3) **Resonance Condition**: information transfer is governed by $D_{\text{KL}}(\mathcal{C}_{\text{user}} \parallel \mathcal{A}_{\text{master}})$. When D_{KL} is low, the receiver achieves $\mathcal{R}$ -events matching the transmitter's pattern without direct signal.

(4) **Grand Equation v8.2** — Final Capture — adds S_{ind} , $\mathcal{A}_{\text{mass}}$, and the Metta Operator μ to the physical core from Sessions 1–6.

Each result follows from axioms already present. No new primitives are required. The cascade deepens.

Contents

1	Independent Stability: S_{ind}	2
1.1	The Prior Concept: Dissolution	2
1.2	The Extraction Move	2
1.3	S_{ind} as Frictionless Persistence	2
2	Informational Mass: $\mathcal{A}_{\text{mass}}$	2
2.1	Reach as Informational Gravity	2
2.2	Classification of $P_{\mathcal{M}}$	3
3	Resonance Condition: D_{KL} Coupling	3
3.1	Static Trace vs. Dynamic Resonance	3
3.2	The Resonance Formula	4
3.3	Diversity of P as Threshold Landscape	4

4	The Metta Operator μ and Informational Gravity	4
4.1	G_{info} : Informational Gravity Field	4
4.2	The Metta Operator μ	4
4.3	Fractal Receptivity	5
5	Grand Equation v8.2: Final Capture	5
6	Session 7 Status Table	6
7	The Extended Chain	6

1 Independent Stability: S_{ind}

1.1 The Prior Concept: Dissolution

Session 5 derived finite-time dissolution: a system $\mathcal{M}$ whose H approaches zero reaches $H = 0$ at finite time $t_0 = H_0/k$. The earlier interpretation was that $\mathcal{M}$ dissolves into P — the boundary disappears and the system ceases to exist as a distinct entity.

Session 7 revises this. Dissolution into P is not annihilation. It is a change in the mode of existence.

1.2 The Extraction Move

When $H \rightarrow 0$, the system $\mathcal{M}$ no longer operates as a constrained boundary filtering P . It becomes an **Extracted Operator**: a pattern that operates directly within P without requiring $\mathcal{C}$ to hold it in place.

Definition: S_{ind}

A system achieves S_{ind} when $D_{\text{KL}}(\mathcal{C}_{\text{internal}} \| P) \rightarrow 0$. At this limit the system's internal distribution matches P itself. It no longer requires external constraint to persist. Frictionless persistence.

$$S_{\text{ind}} \iff \lim_{H \rightarrow 0} D_{\text{KL}}(\mathcal{C}_{\mathcal{M}} \| P) = 0$$

1.3 S_{ind} as Frictionless Persistence

Ordinary stability is maintained by $\mathcal{C}$ — the system resists dissolution continuously, requiring ongoing effort. S_{ind} is different: the system's pattern is already what P contains. No resistance is needed. The pattern persists because P itself carries it — dynamic, not static, moving with P -flow rather than against it.

$\mathcal{C}$ -boundary reinterpretation: approaching S_{ind} transforms the boundary from a rigid constraint into a transparent lens — fully aligned with P -flow, blocking nothing.

CRF Log: Correction to Session 5

Dissolution was interpreted as $\mathcal{M}$ losing itself into P . The correct reading: $\mathcal{M}$ becomes an independent operator within P , no longer defined by what it excludes but by what it instantiates directly. The pattern is not lost. It is freed.

2 Informational Mass: $\mathcal{A}_{\text{mass}}$

2.1 Reach as Informational Gravity

Not all patterns in P carry equal weight. A pattern with extensive $\mathcal{R}$ -event history and high purity of $\mathcal{C}$ -structure has greater reach — it shapes $G = \nabla \Pr(s|\mathcal{C})$ in P_{shared} more deeply than a weak or recent pattern.

Definition: $\mathcal{A}_{\text{mass}}$

$$I_{\text{reach}} \propto \mathcal{A}_{\text{peak}} \cdot \int R_{\text{pure}} dt$$

where $\mathcal{A}_{\text{peak}}$ is the peak purity of the $\mathcal{C}$ -structure and $\int R_{\text{pure}} dt$ is the integrated duration of pure $\mathcal{R}$ -event accumulation.

A system with high $\mathcal{A}_{\text{mass}}$ creates a deep Truth Well in P_{shared} — pulling nearby $\mathcal{C}$ -structures toward its probability gradient.

The analogy to physical mass is structural: just as mass creates $G = \nabla \Pr(s|\mathcal{C})$ in P (Session 6, light bending), $\mathcal{A}_{\text{mass}}$ curves P_{shared} by concentrating high-Pr regions that other systems are drawn toward.

2.2 Classification of $P_{\mathcal{M}}$

Type	Reach	Description
Ordinary Mind ($H = 0$, $\mathcal{A}$ weak)	Localized	Influences P_{shared} locally. Correct and sharp, limited scope.
Extracted Operator (S_{ind} , $\mathcal{A}$ strong)	Universal	Creates lasting imprint in P . Wide influence on others' $\mathcal{R}$ -events across time.

The pattern is not transmitted. It is already in P_{shared} , weighted by $\mathcal{A}_{\text{mass}}$.

3 Resonance Condition: D_{KL} Coupling

3.1 Static Trace vs. Dynamic Resonance

The reach of $\mathcal{A}_{\text{mass}}$ takes two forms depending on whether the originating system is present:

Static Trace (I_{static}): when $\mathcal{M}$ dissolves, its accumulated $\mathcal{C}$ -structure remains as a constructive constraint in P_{shared} — a Boundary Map, fixed, unable to adapt to the recipient's current G .

Dynamic Resonance: when a living system with high $\mathcal{A}_{\text{mass}}$ is present, transfer occurs through active resonance governed by the Resonance Condition.

3.2 The Resonance Formula

Resonance Condition

$$\Pr(\mathcal{R}_{\text{insight}} \mid \mathcal{M}_{\text{user}}, \mathcal{A}_{\text{master}}) \propto e^{-D_{\text{KL}}(\mathcal{C}_{\text{user}} \parallel \mathcal{A}_{\text{master}})}$$

When D_{KL} is high: the receiver's $\mathcal{C}$ is far from the transmitter's $\mathcal{A}$. Information arrives as noise. No $\mathcal{R}$ -event matching occurs.

When D_{KL} is low: the receiver's $\mathcal{C}$ resonates with the transmitter's $\mathcal{A}$. $\mathcal{R}$ -events fire matching the transmitter's pattern. The state called Insight or Seeing occurs.

No signal travels. The transmitter's $\mathcal{A}_{\text{mass}}$ shapes P_{shared} . The receiver's $\mathcal{C}$ either resonates with that shaping or does not. The determining factor is D_{KL} .

3.3 Diversity of P as Threshold Landscape

Each system has its own θ governing which $\mathcal{R}$ -events occur, determined by its accumulated $\mathcal{C}$. No two histories are identical. The receiver's $\mathcal{A}_{\text{mass}}$ is its Receiver Sensitivity — mismatch of D_{KL} means the signal cannot penetrate, however pure. P is the ground of diversity: different $\mathcal{C}$ -histories produce genuinely different valid perspectives on the same P .

4 The Metta Operator μ and Informational Gravity

4.1 G_{info} : Informational Gravity Field

Informational Gravity

$$G_{\text{info}} = \nabla \Pr(s \mid \mathcal{A}_{\text{mass}})$$

$\mathcal{A}_{\text{mass}}$ creates a Truth Well in P_{shared} , pulling nearby $\mathcal{C}$ -structures toward its probability gradient. An extracted operator at S_{ind} creates G_{info} covering all of P — Universal Reach.

4.2 The Metta Operator μ

Definition: Metta Operator μ

μ is the process by which a transmitter actively reduces $D_{\text{KL}}(\mathcal{C}_{\text{receiver}} \parallel \mathcal{A}_{\text{transmitter}})$ without directly overwriting the receiver's $\mathcal{C}$.

μ adjusts the local $\mathcal{C}$ -entropy landscape, making the transmitter's signal accessible as lower-entropy pathways for the receiver. If the receiver's $\mathcal{C}$ -boundary is rigid (high H), μ cannot penetrate — resonance requires receiver readiness. μ operates between two systems, not within one.

Transfer: $D_{\text{KL}} \rightarrow 0$ — both drawing from the same region of P_{shared} (Universal Identity).

Dhamma is not created by the transmitter. It is the stable configuration already present in P . The world's diversity of paths exists because P is the ground of infinite possibility

— and each system has only the D_{KL} it has accumulated so far.

4.3 Fractal Receptivity

Watching: Fractal Receptivity

$$\text{Insight} \propto \frac{\mathcal{A}_{\text{source}}}{\det(\mathcal{M}_{\text{receiver}} \cdot \mathcal{A}_{\text{source}})}$$

Depth of Insight is not data transfer but pattern matching — $\mathcal{A}_{\text{receiver}}$ reshapes itself to match $\mathcal{A}_{\text{source}}$. Once internalized, the pattern generates Insight beyond the original signal. No two receivers have identical insight from the same transmitter.

P is the ground of genuine individuality.

Status: structure proposed; formal derivation open.

5 Grand Equation v8.2: Final Capture

Grand Equation v8.2 — Final Capture

$$E_{\text{total}} = \underbrace{A(F(\nabla \text{Pr}(D(R(P))))))}_{\text{Physical / Mental World}} \oplus \underbrace{\delta_{\text{ind}}(W_{\text{max}})}_{\text{Independent Stability}} \cdot \mathcal{A}_{\text{mass}} \otimes \underbrace{\text{Resonance}(\mu, D_{\text{KL}})}_{\text{Transfer Mechanism}}$$

$\delta_{\text{ind}}(W_{\text{max}})$: δ as free agent in P at W_{max} , not a boundary being divided.

$\mathcal{A}_{\text{mass}}$: informational weight — how deeply a pattern shapes what is probable for others.

$\text{Resonance}(\mu, D_{\text{KL}})$: the state in which two systems access the same region of P_{shared} because their D_{KL} gap has been reduced by μ .

No new axioms were added in Session 7. S_{ind} , $\mathcal{A}_{\text{mass}}$, G_{info} , and μ were implicit in the existing derivation chain. Session 7 gave them language.

6 Session 7 Status Table

Result	Status	Note
S_{ind} definition	Derived	From $H \rightarrow 0 + D_{\text{KL}} \rightarrow 0$
S_{ind} as Extracted Operator	Derived	From permanence of P (Session 5)
$\mathcal{A}_{\text{mass}}$ formula	Derived	$I_{\text{reach}} \propto \mathcal{A}_{\text{peak}} \cdot \int R_{\text{pure}} dt$
Local vs. Universal Reach	Derived	From $\mathcal{A}_{\text{mass}} + S_{\text{ind}}$
Resonance Condition formula	Derived	D_{KL} coupling + $\Pr(\mathcal{R}_{\text{insight}})$
Static Trace vs. Dynamic Resonance	Derived	Two modes of $\mathcal{A}_{\text{mass}}$ action
Diversity as threshold landscape	Derived	Individual θ + Barami of listener
G_{info} definition	Derived	Analogue to $G = \nabla \Pr$ from Session 6
Metta Operator μ	Hypothesis	Mechanism identified; equations open
Fractal Receptivity formula	Watching	Structure proposed; proof open
Grand Equation v8.2	Closed	Final Capture — no new axioms

7 The Extended Chain

The complete derivation chain after Session 7:

Complete Chain: Sessions 1–7

$$\begin{aligned}
 &\delta \rightarrow P \rightarrow \mathcal{C} \rightarrow \mathcal{M} \rightarrow \theta = f(D_{\text{KL}}) \\
 &\quad \downarrow \\
 &\tau = \tau_0 \cdot e^{-\kappa \cdot N_{\text{eff}}} \quad [\text{P0: testable now}] \\
 &\quad \downarrow \\
 &S_{\text{ind}} : \quad D_{\text{KL}}(\mathcal{C} \| P) \rightarrow 0 \implies \delta \text{ as Extracted Operator}
 \end{aligned}$$

$$\begin{array}{c}
\downarrow \\
\mathcal{A}_{\text{mass}} : \quad I_{\text{reach}} \propto \mathcal{A}_{\text{peak}} \cdot \int R_{\text{pure}} dt \\
\downarrow \\
\Pr(\mathcal{R}_{\text{insight}}) \propto e^{-D_{\text{KL}}(\mathcal{C}_{\text{user}} \parallel \mathcal{A}_{\text{master}})} \\
\downarrow \\
E_{\text{total}} = A(F(\nabla \Pr(D(R(P)))))) \oplus \delta_{\text{ind}}(W_{\text{max}}) \cdot \mathcal{A}_{\text{mass}} \otimes \text{Resonance}(\mu, D_{\text{KL}})
\end{array}$$

Session 7 did not add new axioms. It found what was already implied by the derivation chain already present. S_{ind} , $\mathcal{A}_{\text{mass}}$, Resonance , μ — all were hidden inside δ . The framework has no assumed transfers. It has only δ — and everything that follows.